



**TOXICOLOGICAL REVIEW
OF
TRICHLOROETHYLENE
CHAPTER 7**

(CAS No. 79-01-6)

**In Support of Summary Information on the
Integrated Risk Information System (IRIS)**

September 2011

7. REFERENCES

- 't Mannetje, A; Steenland, K; Attfield, M; Boffetta, P; Checkoway, H; DeKlerk, N; Koskela, R. (2002). Exposure-response analysis and risk assessment for silica and silicosis mortality in a pooled analysis of six cohorts. *Occup Environ Med* 59: 723-728.
- 't Mannetje, A; Dryson, E; Walls, C; McLean, D; McKenzie, F; Maule, M; Cheng, S; Cunningham, C; Kromhout, H; Boffetta, P; Blair, A; Pearce, N. (2008). High risk occupations for non-Hodgkin's lymphoma in New Zealand: case-control study. *Occup Environ Med* 65: 354-363. <http://dx.doi.org/10.1136/oem.2007.035014>.
- Abbas, R; Seckel, C; Kidney, J; Fisher, J. (1996). Pharmacokinetic analysis of chloral hydrate and its metabolism in B6C3F1 mice. *Drug Metab Dispos* 24: 1340-1346.
- Abbas, R and Fisher, J. (1997). A physiologically based pharmacokinetic model for trichloroethylene and its metabolites, chloral hydrate, trichloroacetate, dichloroacetate, trichloroethanol, and trichloroethanol glucuronide in B6C3F1 mice. *Toxicol Appl Pharmacol* 147: 15-30. <http://dx.doi.org/10.1006/taap.1997.8190>.
- Abbas, R; Seckel, C; MacMahon, K; Fisher, J. (1997). Determination of kinetic rate constants for chloral hydrate, trichloroethanol, trichloroacetic acid and dichloroacetic acid: A physiologically based modeling approach. *Toxicologist* 36: 32-33.
- Abbate, C; Giorgianni, C; Munao, F; Brecciaroli, R. (1993). Neurotoxicity induced by exposure to toluene: An electrophysiologic study. *Int Arch Occup Environ Health* 64: 389-392. <http://dx.doi.org/10.1007/BF00517943>.
- Abraham, D; Patel, P; Cooper, A. (1995a). Isolation from rat kidney of a cytosolic high molecular weight cysteine-S-conjugate beta-lyase with activity toward leukotriene E4. *J Biol Chem* 270: 180-188. <http://dx.doi.org/10.1074/jbc.270.1.180>.
- Abraham, D; Thomas, R; Cooper, A. (1995b). Glutamine transaminase K is not a major cysteine S-conjugate beta-lyase of rat kidney mitochondria: Evidence that a high-molecular weight enzyme fulfills this role. *Mol Pharmacol* 48: 855-860.
- Adachi, H; Kubota, I; Okamura, N; Iwata, H; Tsujimoto, M; Nakazato, H; Nishihara, T; Noguchi, T. (1989). Purification and characterization of human microsomal dipeptidase. *J Biochem* 105: 957-961.
- Adami, H; Chow, W; Nyrén, O; Berne, C; Linet, M; Ekbom, A; Wolk, A; McLaughlin, J; Fraumeni, J. (1996). Excess risk of primary liver cancer in patients with diabetes mellitus. *J Natl Cancer Inst* 88: 1472-1477.
- Adams, C; Keil, D; Meyers, K; Eudaly, A; Smythe, J; Eudaly, J; Gilkeson, G; Peden-Adams, M. (2003). Lifetime exposure to trichloroethylene (TCE) modulates immune function. *Toxicologist* 72: 375.
- Adams, L and Lindor, K. (2007). Nonalcoholic fatty liver disease. *Ann Epidemiol* 17: 863-869. <http://dx.doi.org/10.1016/j.annepidem.2007.05.013>.
- Adgate, J; Church, T; Ryan, A; Ramachandran, G; Fredrickson, A; Stock, T; Morandi, M; Sexton, K. (2004a). Outdoor, indoor, and personal exposure to VOCs in children. *Environ Health Perspect* 112: 1386-1392. <http://dx.doi.org/10.1289/ehp.7107>.
- Adgate, J; Eberly, L; Stroebel, C; Pellizzari, E; Sexton, K. (2004b). Personal, indoor, and outdoor VOC exposures in a probability sample of children. *J Expo Anal Environ Epidemiol* 14: S4-S13. <http://dx.doi.org/10.1038/sj.jea.7500353>.
- ADHS. (Arizona Department of Health Services). (1990). Incidence study of childhood cancer in Maricopa County: 1965-1986. Phoenix, AZ.
- ADHS. (Arizona Department of Health Services). (1995). Update of the incidence of childhood cancers and testicular cancer in southwest Tucson: 1987-1991. Phoenix, AZ.
- Adler, I. (1993). Synopsis of the in vivo results obtained with the 10 known or suspected aneugens tested in the CEC collaborative study. *DNA Repair* 287: 131-137. [http://dx.doi.org/10.1016/0027-5107\(93\)90152-6](http://dx.doi.org/10.1016/0027-5107(93)90152-6).

- Adli, M and Baldwin, A. (2006). IKK-i/IKKepsilon controls constitutive, cancer cell-associated NF-kappaB activity via regulation of Ser-536 p65/RelA phosphorylation. *J Biol Chem* 281: 26976-26984. <http://dx.doi.org/10.1074/jbc.M603133200>.
- Aickin, M; Chapin, C; Flood, T; Englender, S; Caldwell, G. (1992). Assessment of the spatial occurrence of childhood leukaemia mortality using standardized rate ratios with a simple linear Poisson model. *Int J Epidemiol* 21: 649-655. <http://dx.doi.org/10.1093/ije/21.4.649>.
- Aickin, M. (2004). Bayes without priors. *J Clin Epidemiol* 57: 4-13. [http://dx.doi.org/10.1016/S0895-4356\(03\)00251-8](http://dx.doi.org/10.1016/S0895-4356(03)00251-8).
- Aikata, H; Takaishi, H; Kawakami, Y; Takahashi, S; Kitamoto, M; Nakanishi, T; Nakamura, Y; Shimamoto, F; Kajiyama, G; Ide, T. (2000). Telomere reduction in human liver tissues with age and chronic inflammation. *Exp Cell Res* 256: 578-582. <http://dx.doi.org/10.1006/excr.2000.4862>.
- Akira, S and Takeda, K. (2004). Toll-like receptor signalling. *Nat Rev Immunol* 4: 499-511. <http://dx.doi.org/10.1038/nri1391>.
- Akiyama, T; Nicol, C; Fievet, C; Staels, B; Ward, J; Auwerx, J; Lee, S; Gonzalez, F; Peters, J. (2001). Peroxisome proliferator-activated receptor-alpha regulates lipid homeostasis, but is not associated with obesity: Studies with congenic mouse lines. *J Biol Chem* 276: 39088-39093. <http://dx.doi.org/10.1074/jbc.M107073200>.
- Albanese, R; Banks, H; Evans, M; Potter, L. (2002). Physiologically based pharmacokinetic models for the transport of trichloroethylene in adipose tissue. *Bull Math Biol* 64: 97-131. <http://dx.doi.org/10.1006/bulm.2001.0268>.
- Albee, R; Nitschke, K; Mattsson, J; Stebbins, K. (1997). Dichloroacetylene: Effects on the rat trigeminal nerve somatosensory evoked potential. *Neurotoxicol Teratol* 19: 27-37.
- Albee, R; Spencer, P; Johnson, K; Bradley, G; Marable, B; Wilmer, J; Mattsson, J. (2006). Lack of trigeminal nerve toxicity in rats exposed to trichloroethylene vapor for 13 weeks. *Int J Toxicol* 25: 531-540. <http://dx.doi.org/10.1080/10915810600972678>.
- Alberati-Giani, D; Malherbe, P; Kohler, C; Lang, G; Kiefer, V; Lahm, H; Cesura, A. (1995). Cloning and characterization of a soluble kynurenine aminotransferase from rat brain: Identity with kidney cysteine conjugate [beta]-lyase. *J Neurochem* 64: 1448-1455. <http://dx.doi.org/10.1046/j.1471-4159.1995.64041448.x>.
- Albers, J; Wald, J; Werner, R; Franzblau, A; Berent, S. (1999). Absence of polyneuropathy among workers previously diagnosed with solvent-induced toxic encephalopathy. *J Occup Environ Med* 41: 500-509.
- Albert, D; Albert, A; Vernace, M; Sebastian, J; Hsia, E. (2005). Analysis of a cluster of cases of Wegener granulomatosis. *J Clin Rheumatol* 11: 188-193.
- Albertini, S. (1990). Analysis of nine known or suspected spindle poisons for mitotic chromosome malsegregation using *Saccharomyces cerevisiae* D61.M. *Mutagenesis* 5: 453-459.
- Alexander, D; Mink, P; Mandel, J; Kelsh, M. (2006). A meta-analysis of occupational trichloroethylene exposure and multiple myeloma or leukaemia. *Occup Med (Lond)* 56: 485-493. <http://dx.doi.org/10.1093/occmed/kql083>.
- Alexander, D; Kelsh, M; Mink, P; Mandel, J; Basu, R; Weingart, M. (2007a). A meta-analysis of occupational trichloroethylene exposure and liver cancer. *Int Arch Occup Environ Health* 81: 127-143. <http://dx.doi.org/10.1007/s00420-007-0201-4>.
- Alexander, D; Mink, P; Adami, H; Chang, E; Cole, P; Mandel, J; Trichopoulos, D. (2007b). The non-Hodgkin lymphomas: A review of the epidemiologic literature. *Int J Cancer* 120: 1-39. <http://dx.doi.org/10.1002/ijc.22719>.
- Alimov, A; Kost-Alimova, M; Liu, J; Li, C; Bergerheim, U; Imreh, S; Klein, G; Zabarovsky, E. (2000). Combined LOH/CGH analysis proves the existence of interstitial 3p deletions in renal cell carcinoma. *Oncogene* 19: 1392-1399. <http://dx.doi.org/10.1038/sj.onc.1203449>.
- Allen, B; Shipp, A; Crump, K; Kilian, B; Hogg, M; Tudor, J; Keller, B. (1987). Investigation of cancer risk assessment methods (3 volumes plus summary report). Washington, D.C.: U.S. Environmental Protection Agency.

- Allen, B and Fisher, J. (1993). Pharmacokinetic modeling of trichloroethylene and trichloroacetic acid in humans. Risk Anal 13: 71-86.
- Allen, D; Pearse, G; Haseman, J; Maronpot, R. (2004). Prediction of rodent carcinogenesis: An evaluation of prechronic liver lesions as forecasters of liver tumors in NTP carcinogenicity studies. Toxicol Pathol 32: 393-401. <http://dx.doi.org/10.1080/01926230490440934>.
- Allen, J; Collins, B; Evansky, P. (1994). Spermatid micronucleus analyses of trichloroethylene and chloral hydrate effects in mice. Mutat Res Lett 323: 81-88. [http://dx.doi.org/10.1016/0165-7992\(94\)90049-3](http://dx.doi.org/10.1016/0165-7992(94)90049-3).
- Altmann, L; Welge, P; Mensing, T; Lilienthal, H; Voss, B; Wilhelm, M. (2002). Chronic exposure to trichloroethylene affects neuronal plasticity in rat hippocampal slices. Environ Toxicol Pharmacol 12: 157-167.
- Altuntas, T; Park, S; Kharasch, E. (2004). Sulfoxidation of cysteine and mercapturic acid conjugates of the sevoflurane degradation product fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether (compound A). Chem Res Toxicol 17: 435-445. <http://dx.doi.org/10.1021/tx034254k>.
- Amacher, D and Zelljadt, I. (1983). The morphological transformation of Syrian hamster embryo cells by chemicals reportedly nonmutagenic to *Salmonella typhimurium*. Carcinogenesis 4: 291-296.
- American Cancer Society. (2008). What are the risk factors for cervical cancer? pp. 3-7. Atlanta, GA. http://www.cancer.org/docroot/CRI/content/CRI_2_4_2X_What_are_the_risk_factors_for_cervical_cancer_8.asp.
- Amet, Y; Berthou, F; Fournier, G; Dreano, Y; Bardou, L; Cledes, J; Menez, J. (1997). Cytochrome P450 4A and 2E1 expression in human kidney microsomes. Biochem Pharmacol 53: 765-771. [http://dx.doi.org/10.1016/S0006-2952\(96\)00821-0](http://dx.doi.org/10.1016/S0006-2952(96)00821-0).
- Ammini, C; Fernandez-Canon, J; Shroads, A; Cornett, R; Cheung, J; James, M; Henderson, G; Grompe, M; Stacpoole, P. (2003). Pharmacologic or genetic ablation of maleylacetoacetate isomerase increases levels of toxic tyrosine catabolites in rodents. Biochem Pharmacol 66: 2029-2038. <http://dx.doi.org/10.1016/j.bcp.2003.07.002>.
- Andelman, J. (1985). Human exposures to volatile halogenated organic chemicals in indoor and outdoor air. Environ Health Perspect 62: 313-318.
- Anders, M and Jakobson, I. (1985). Biotransformation of halogenated solvents. Scand J Work Environ Health 11: 23-32.
- Andersen, M; Clewell, H, III; Gargas, M; Smith, F; Reitz, R. (1987a). Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. Toxicol Appl Pharmacol 87: 185-205. [http://dx.doi.org/10.1016/0041-008X\(87\)90281-X](http://dx.doi.org/10.1016/0041-008X(87)90281-X).
- Andersen, M; Gargas, M; Clewell, H, III; Severyn, K. (1987b). Quantitative evaluation of the metabolic interactions between trichloroethylene and 1,1-dichloroethylene in vivo using gas uptake methods. Toxicol Appl Pharmacol 89: 149-157.
- Anderson, B; McKee, A; Holford, N. (1997). Size, myths and the clinical pharmacokinetics of analgesia in paediatric patients. Clin Pharmacokinet 33: 313-327.
- Anderson, S; Howroyd, P; Liu, J; Qian, X; Bahnemann, R; Swanson, C; Kwak, M; Kensler, T; Corton, J. (2004). The transcriptional response to a peroxisome proliferator-activated receptor alpha agonist includes increased expression of proteome maintenance genes. J Biol Chem 279: 52390-52398. <http://dx.doi.org/10.1074/jbc.M409347200>.
- Andersson, L; Haraldsson, B; Johansson, C; Barregard, L. (2008). Methodological issues on the use of urinary alpha-1-microglobuline in epidemiological studies. Nephrol Dial Transplant 23: 1252-1256. <http://dx.doi.org/10.1093/ndt/gfm729>.
- Anna, C; Maronpot, R; Pereira, M; Foley, J; Malarkey, D; Anderson, M. (1994). Ras proto-oncogene activation in dichloroacetic acid-, trichloroethylene- and tetrachloroethylene-induced liver tumors in B6C3F1 mice. Carcinogenesis 15: 2255-2261. <http://dx.doi.org/10.1093/carcin/15.10.2255>.
- Anti, M; Marra, G; Rapaccini, G; Rumi, C; Bussa, S; Fadda, G; Vecchio, F; Valenti, A; Percesepe, A; Pompili, M. (1994). DNA ploidy pattern in human chronic liver diseases and hepatic nodular lesions. Flow cytometric analysis on echo-guided needle liver biopsy. Cancer 73: 281-288.

- Antti-Poika, M. (1982). Overall prognosis of patients with diagnosed chronic organic solvent intoxication. Int Arch Occup Environ Health 51: 127-138.
<http://dx.doi.org/10.1007/BF00378157>.
- Anttila, A; Pukkala, E; Sallmen, M; Hernberg, S; Hemminki, K. (1995). Cancer incidence among Finnish workers exposed to halogenated hydrocarbons. J Occup Environ Med 37: 797-806.
- Anway, M; Cupp, A; Uzumcu, M; Skinner, M. (2005). Epigenetic transgenerational actions of endocrine disruptors and male fertility. Science 308: 1466-1469. <http://dx.doi.org/10.1126/science.1108190>.
- Araki, S. (1978). The effects of water restriction and water loading on urinary excretion of lead, delta-aminolevulinic acid and coproporphyrin. Br J Ind Med 35: 312-317.
- Aranyi, C; O'Shea, W; Graham, J; Miller, F. (1986). The effects of inhalation of organic chemical air contaminants on murine lung host defenses. Fundam Appl Toxicol 6: 713-720.
[http://dx.doi.org/10.1016/0272-0590\(86\)90184-3](http://dx.doi.org/10.1016/0272-0590(86)90184-3).
- Aratani, J; Suzuki, H; Hashimoto, K. (1993). Measurement of vibratory perception threshold (VPT) in workers exposed to organic solvents. Environ Res 61: 357-361.
<http://dx.doi.org/10.1006/enrs.1993.1080>.
- Arif, A and Shah, S. (2007). Association between personal exposure to volatile organic compounds and asthma among US adult population. Int Arch Occup Environ Health 80: 711-719.
<http://dx.doi.org/10.1007/s00420-007-0183-2>.
- Arito, H; Takahashi, M; Ishikawa, T. (1994). Effect of subchronic inhalation exposure to low-level trichloroethylene on heart rate and wakefulness-sleep in freely moving rats. Sangyo Eiseigaku Zasshi 36: 1-8.
- Aronson, K; Siemiatycki, J; Dewar, R; Gérin, M. (1996). Occupational risk factors for prostate cancer: Results from a case-control study in Montréal, Québec, Canada. Am J Epidemiol 143: 363-373.
- Arp, E; Wolf, P; Checkoway, H. (1983). Lymphocytic leukemia and exposures to benzene and other solvents in the rubber industry. J Occup Med 25: 598-602.
- Aryal, B; Khuder, S; Schaub, E. (2001). Meta-analysis of systemic sclerosis and exposure to solvents. Am J Ind Med 40: 271-274. <http://dx.doi.org/10.1002/ajim.1098>.
- Asal, N; Geyer, J; Risser, D; Lee, E; Kadaman, S; Cherng, N. (1988a). Risk factors in renal cell carcinoma. II. Medical history, occupation, multivariate analysis, and conclusions. Cancer Detect Prev 13: 263-279.
- Asal, N; Risser, D; Kadaman, S; Geyer, J; Lee, E; Cherng, N. (1988b). Risk factors in renal cell carcinoma: I. Methodology, demographics, tobacco, beverage use, and obesity. Cancer Detect Prev 11: 359-377.
- Ashby, J; Brady, A; Elcombe, C; Elliott, B; Ishmael, J; Odum, J; Tugwood, J; Kettle, S; Purchase, I. (1994). Mechanistically-based human hazard assessment of peroxisome proliferator-induced hepatocarcinogenesis. Hum Exp Toxicol 13: S1-S117.
<http://dx.doi.org/10.1177/096032719401300201>.
- Ashley, D; Bonin, M; Cardinali, F; McCraw, J; Wooten, J. (1994). Blood concentrations of volatile organic compounds in a nonoccupationally exposed US population and in groups with suspected exposure. Clin Chem 40: 1401-1404.
- Astrand, I. (1975). Uptake of solvents in the blood and tissues of man. A review. Scand J Work Environ Health 1: 199-218.
- Astrand, I and Ovrum, P. (1976). Exposure to trichloroethylene I. Uptake and distribution in man. Scand J Work Environ Health 4: 199-211.
- Astrand, I. (1982). Work load and uptake of solvents in tissues of man. 141 - 152.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1990). Toxicological profile for cis-, trans-1,2-dichloroethene. Atlanta, GA: U.S. Department of Health and Human Services.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1995). Toxicological profile for 1,1,1-trichloroethane. Atlanta, GA: U.S. Department of Health and Human Services.
<http://www.worldcat.org/title/toxicological-profile-for-111-trichloroethane/oclc/33599544>.

- ATSDR. (Agency for Toxic Substances and Disease Registry). (1996a). Toxicological profile for 1,2-dichloroethene. Atlanta, GA: U.S. Department of Health and Human Services.
<http://www.atsdr.cdc.gov/ToxProfiles/tp.asp?id=464&tid=82>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1996b). Biennial report to congress (1991 and 1992). Atlanta, GA: U.S. Department of Health and Human Services.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1997a). Toxicological profile for tetrachloroethylene. Atlanta, GA: U.S. Department of Health and Humans Services.
<http://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=265&tid=48>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1997b). Toxicological profile for trichloroethylene (update). (PB98-101165). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
<http://www.atsdr.cdc.gov/toxprofiles/tp19-p.pdf>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1997c). Toxicological profile for trichloroethylene (TCE). Atlanta, GA: U.S. Department of Health and Human Services.
<http://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=173&tid=30>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1998a). Volatile organic compounds in drinking water and adverse pregnancy outcomes: United States Marine Corps Base, Camp Lejeune, North Carolina. <http://www.atsdr.cdc.gov/hs/lejeune/>
- ATSDR. (Agency for Toxic Substances and Disease Registry). (1998b). Health consultation: Evaluation of health impacts from potential past exposure to tetrachloroethylene in the Natick public water supply. Atlanta, GA: Agency for Toxic Substances & Disease Registry.
<http://www.atsdr.cdc.gov/hac/pha/pha.asp?docid=1270&pg=0>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2000). Public Health Assessment for Tucson International Airport Area (a/k/a El Vado Residential Properties) Tucson, Pima County, Arizona, Region 9. CERCLIS No. AZD980737530. pp. 44p. Office of Environmental Health, Environmental Health Consultation Services: U.S. Department of Health and Humans Services.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2001). Final Report: Evaluation of priority health conditions in a community with historical contamination by trichloroethylene. Atlanta, GA: U.S. Department of Health and Human Services.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2002). Impact of trichloroethylene exposure on oral motor, speech, and hearing in children. Atlanta, GA: U.S. Department of Health and Human Services. <http://www.ntis.gov/search/product.aspx?ABBR=PB2004100016>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2003a). Progress report: Survey of specific childhood cancers and birth defects among children whose mothers were pregnant while living at U.S. Marine Corps Base Camp Lejeune, North Carolina, 1968-1985.
<http://www.atsdr.cdc.gov/sites/lejeune/lejeunehealthsurvey.pdf>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2003b). Health consultation: View-Master factory supply well (a.k.a. Mattel Portland Operations) Beaverton, Washington County, Oregon. Atlanta, GA: U.S. Department of Health and Human Services.
<http://www.atsdr.cdc.gov/hac/pha/pha.asp?docid=293&pg=0>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2004a). Feasibility investigation of worker exposure to trichloroethylene at the View-Master factory in Beaverton, Oregon. Atlanta, GA: U.S. Department of Health and Human Services.
<http://www.ntis.gov/search/product.aspx?ABBR=PB2005102173>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2004b). Interaction profile: 1,1,1-trichloroethane, 1,1-dichloroethane, trichloroethylene, and tetrachloroethylene. Atlanta, GA.
<http://www.atsdr.cdc.gov/interactionprofiles/ip02.html>.
- ATSDR. (Agency for Toxic Substances and Disease Registry). (2006a). Health statistics review: Cancer and birth outcome analysis: Endicott area investigation: Endicott area, town of Union, Broome County, New York. Atlanta, GA: U.S. Department of Health and Humans Services.

<http://www.atsdr.cdc.gov/HAC/pha/EndicottAreaInvestigation/EndicottHealthStatsReviewHC052606.pdf>.

ATSDR. (Agency for Toxic Substances and Disease Registry). (2006b). Health consultation: Public health implications of exposures to low-level volatile organic compounds in public drinking water: Endicott area investigation, Broome County, New York. Atlanta, GA: U.S. Department of Health and Human Services.

<http://www.atsdr.cdc.gov/HAC/pha/EndicottAreaInvestigation113006/EndicottAreaInvestigationHC113006.pdf>.

ATSDR. (Agency for Toxic Substances and Disease Registry). (2007a). Analyses of groundwater flow, contaminant fate and transport and distribution of drinking water at Tarawa Terrace and vicinity, U.S. Marine Corps Base Camp Lejeune, North Carolina: Historical reconstruction and present-day conditions. Atlanta, GA: U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/sites/lejeune/tarawaterrace.html>.

ATSDR. (2007b) The verbatim transcript of the Meeting of the Camp Lejeune Community Assistance Panel held at the ATSDR, 1825 Century Boulevard, Atlanta, Georgia, on December 6, 2007 Meeting of the Camp Lejeune Community Assistance Panel. Atlanta, GA: Author.

ATSDR. (Agency for Toxic Substances and Disease Registry). (2008a). Toxicological profile for 1,1,2,2 tetrachloroethane. Atlanta, GA: U.S. Department of Health and Humans Services.

<http://www.atsdr.cdc.gov/ToxProfiles/tp.asp?id=801&tid=156>.

ATSDR. (Agency for Toxic Substances and Disease Registry). (2008b). Health consultation: Health statistics review follow-up: Cancer and birth outcome analysis: Endicott area investigation, Endicott area, Town of Union, Broome County, New York. Atlanta, GA: U.S. Department of Health and Human Services.

<http://www.atsdr.cdc.gov/hac/pha//EndicottAreaInvestigationFollowUp/EndicottAreaHC051508.pdf>.

ATSDR. (Agency for Toxic Substances and Disease Registry). (2009). Study on birth defects and childhood cancers. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Services. <http://www.atsdr.cdc.gov/sites/lejeune/update.html>.

ATSDR. (Agency for Toxic Substances and Disease Registry). (2010). The verbatim transcript of the Meeting of the Camp Lejeune Community Assistance Panel held at the Chamblee Building 106, Conference Room B, Atlanta, Georgia, on Dec. 9, 2010. Atlanta, GA.

http://www.atsdr.cdc.gov/sites/lejeune/docs/CAPtranscript_12_10.pdf.

Attallah, A; Tabll, A; Salem, S; El-Sadany, M; Ibrahim, T; Osman, S; El-Dosoky, I. (1999). DNA ploidy of liver biopsies from patients with liver cirrhosis and hepatocellular carcinoma: A flow cytometric analysis. *Cancer Lett* 142: 65-69.

Auperin, A; Benhamou, S; Ory-Paoletti, C; Flamant, R. (1994). Occupational risk factors for renal cell carcinoma: A case-control study. *Occup Environ Med* 51: 426-428.

<http://dx.doi.org/10.1136/oem.51.6.426>.

Austin, E; Parrish, J; Kinder, D; Bull, R. (1996). Lipid peroxidation and formation of 8-hydroxydeoxyguanosine from acute doses of halogenated acetic acids. *Fundam Appl Toxicol* 31: 77-82. <http://dx.doi.org/10.1006/faat.1996.0078>.

Austin, H; Delzell, E; Gruffman, S; Levine, R; Morrison, A; Stolley, P; Cole, P. (1987). Case-control study of hepatocellular carcinoma, occupation, and chemical exposures. *J Occup Med* 29: 665-669.

Auwerx, J; Schoonjans, K; Fruchart, J; Staels, B. (1996). Transcriptional control of triglyceride metabolism: Fibrates and fatty acids change the expression of the LPL and apo C-III genes by activating the nuclear receptor PPAR. *Atherosclerosis* 124: S29-S37.

[http://dx.doi.org/10.1016/0021-9150\(96\)05854-6](http://dx.doi.org/10.1016/0021-9150(96)05854-6).

Axelson, O; Seldén, A; Andersson, K; Hogstedt, C. (1994). Updated and expanded Swedish cohort study on trichloroethylene and cancer risk. *J Occup Med* 36: 556-562.

- Baden, J; Kelley, M; Mazze, R; Simmon, V. (1979). Mutagenicity of inhalation anaesthetics: trichloroethylene, divinyl ether, nitrous oxide and cyclopropane. Br J Anaesth 51: 417-421.
- Badger, T; Ronis, M; Seitz, H; Albano, E; Ingelman-Sundberg, M; Lieber, C. (2003). Alcohol metabolism: Role in toxicity and carcinogenesis. Alcohol Clin Exp Res 27: 336-347.
<http://dx.doi.org/10.1097/01.ALC.0000052583.87673.37>.
- Bai, C-L and Stacey, N. (1993). Mechanism of trichloroethylene-induced elevation of individual serum bile acids: II. In vitro and in vivo interference by trichloroethylene with bile acid transport in isolated rat hepatocytes. Toxicol Appl Pharmacol 121: 296-302.
<http://dx.doi.org/10.1006/taap.1993.1157>.
- Bai, C; Canfield, P; Stacey, N. (1992a). Effects of hexachloro-1,3-butadiene and 1,1,2,2-tetrachloroethylene on individual serum bile acids. Toxicol Ind Health 8: 191-203.
- Bai, C; Canfield, P; Stacey, N. (1992b). Individual serum bile acids as early indicators of carbon tetrachloride- and chloroform-induced liver injury. Toxicology 75: 221-234.
[http://dx.doi.org/10.1016/0300-483X\(92\)90004-X](http://dx.doi.org/10.1016/0300-483X(92)90004-X).
- Bailer, A and Piegorsch, W. (1997). Statistics for environmental biology and toxicology. London, England: Chapman & Hall/CRC Press.
- Bailly, M; Bain, C; Favrot, M; Ozturk, M. (1995). Somatic mutations of von Hippel-Lindau (VHL) tumor-suppressor gene in European kidney cancers. Int J Cancer 63: 660-664.
- Baker, V; Harries, H; Waring, J; Duggan, C; Ni, H; Jolly, R; Yoon, L; De Souza, A; Schmid, J; Brown, R; Ulrich, R; Rockett, J. (2004). Clofibrate-induced gene expression changes in rat liver: A cross-laboratory analysis using membrane cDNA arrays. Environ Health Perspect 112: 428-438.
<http://dx.doi.org/10.1289/ehp.6677>.
- Bakke, B; Stewart, P; Waters, M. (2007). Uses of and exposure to trichloroethylene in U.S. industry: A systematic literature review. J Occup Environ Hyg 4: 375-390.
<http://dx.doi.org/10.1080/15459620701301763>.
- Balkwill, F and Coussens, L. (2004). Cancer: An inflammatory link. Nature 431: 405-406.
<http://dx.doi.org/10.1038/431405a>.
- Balkwill, F; Charles, K; Mantovani, A. (2005). Smoldering and polarized inflammation in the initiation and promotion of malignant disease. Cancer Cell 7: 211-217.
<http://dx.doi.org/10.1016/j.ccr.2005.02.013>.
- Ballestar, E and Esteller, M. (2002). The impact of chromatin in human cancer: Linking DNA methylation to gene silencing. Carcinogenesis 23: 1103-1109.
<http://dx.doi.org/10.1093/carcin/23.7.1103>.
- Balluz, L; Phalen, R; Ortega, L; Rosales, C; Brock, J; Barr, D; Kieszak, S. (2001). Investigation of systemic lupus erythematosus in Nogales, Arizona. Am J Epidemiol 154: 1029-1036.
- Band, P; Le, N; Fang, R; Deschamps, M; Gallagher, R; Yang, P. (2000). Identification of occupational cancer risks in British Columbia: A population-based case-control study of 995 incident breast cancer cases by menopausal status, controlling for confounding factors. J Occup Environ Med 42: 284-310.
- Banerjee, S and Van Duuren, B. (1978). Covalent binding of the carcinogen trichloroethylene to hepatic microsomal proteins and to exogenous DNA in vitro. Cancer Res 38: 776-780.
- Banks, P. (1992). Changes in diagnosis of non-Hodgkin's lymphomas over time. Cancer Res 52: 5453s-5455s.
- Bannasch, P. (1996). Pathogenesis of hepatocellular carcinoma: Sequential cellular, molecular, and metabolic changes. Prog Liver Dis 14: 161-197.
- Bannasch, P. (2001). Hormonal and hormone-like effects eliciting hepatocarcinogenesis. Folia Histochem Cytophiol 39: 28-29.
- Bannasch, P; Nehrbass, D; Kopp-Schneider, A. (2001). Significance of hepatic preneoplasia for cancer chemoprevention. In AB Miller, H Bartsch, P Bofetta, L Dragsted & H Vainio (Eds.), Biomarkers in cancer chemoprevention (Vol. 154, pp. 223-240). Lyon, France: International Agency for Research on Cancer.

- Bannasch, P; Haertel, T; Su, Q. (2003). Significance of hepatic preneoplasia in risk identification and early detection of neoplasia. *Toxicol Pathol* 31: 134-139.
<http://dx.doi.org/10.1080/01926230390173923>.
- Barak, Y; Sadovsky, Y; Shalom-Barak, T. (2008). PPAR Signaling in Placental Development and Function. *PPAR Research* 2008: 142082. <http://dx.doi.org/10.1155/2008/142082>.
- Baraona, E; Shoichet, L; Navder, K; Lieber, C. (2002a). Mediation by nitric oxide of the stimulatory effects of ethanol on blood flow. *Life Sci* 70: 2987-2995. [http://dx.doi.org/10.1016/S0024-3205\(02\)01572-2](http://dx.doi.org/10.1016/S0024-3205(02)01572-2).
- Baraona, E; Zeballos, G; Shoichet, L; Mak, K; Lieber, C. (2002b). Ethanol consumption increases nitric oxide production in rats, and its peroxynitrite-mediated toxicity is attenuated by polyenylphosphatidylcholine. *Alcohol Clin Exp Res* 26: 883-889.
<http://dx.doi.org/10.1111/j.1530-0277.2002.tb02618.x>.
- Bardodej, Z and Vyskocil, J. (1956). The problem of trichloroethylene in occupational medicine: trichloroethylene metabolism and its effect on the nervous system evaluated as a means of hygienic control. *Arch Environ Occup Health* 13: 581-592.
- Barrass, N; Price, R; Lake, B; Orton, T. (1993). Comparison of the acute and chronic mitogenic effects of the peroxisome proliferators methylclofenapate and clofibrate acid in rat liver. *Carcinogenesis* 14: 1451-1456. <http://dx.doi.org/10.1093/carcin/14.7.1451>.
- Barret, L; Arsac, P; Vincent, M; Faure, J; Garrel, S; Reymond, F. (1982). Evoked trigeminal nerve potential in chronic trichloroethylene intoxication. *Clin Toxicol* 19: 419-423.
<http://dx.doi.org/10.3109/15563658208992496>.
- Barret, L; Faure, J; Guillard, B; Chomat, D; Didier, B; Debru, J. (1984). Trichloroethylene occupational exposure: Elements for better prevention. *Int Arch Occup Environ Health* 53: 283-289.
<http://dx.doi.org/10.1007/BF00380666>.
- Barret, L; Garrel, S; Danel, V; Debru, J. (1987). Chronic trichloroethylene intoxication: A new approach by trigeminal-evoked potentials? *Arch Environ Occup Health* 42: 297-302.
<http://dx.doi.org/10.1080/00039896.1987.9935824>.
- Barret, L; Torch, S; Usson, Y; Gonthier, B; Saxod, R. (1991). A morphometric evaluation of the effects of trichloroethylene and dichloroacetylene on the rat mental nerve. Preliminary results. *Neurosci Lett* 131: 141-144. [http://dx.doi.org/10.1016/0304-3940\(91\)90598-N](http://dx.doi.org/10.1016/0304-3940(91)90598-N).
- Barret, L; Torch, S; Leray, C; Sarlièvre, L; Saxod, R. (1992). Morphometric and biochemical studies in trigeminal nerve of rat after trichloroethylene or dichloroacetylene oral administration. *Neurotoxicology* 13: 601-614.
- Barter, Z; Bayliss, M; Beaune, P; Boobis, A; Carlile, D; Edwards, R; Houston, J; Lake, B; Lipscomb, J; Pelkonen, O; Tucker, G; Rostami-Hodjegan, A. (2007). Scaling factors for the extrapolation of in vivo metabolic drug clearance from in vitro data: Reaching a consensus on values of human microsomal protein and hepatocellularity per gram of liver. *Curr Drug Metab* 8: 33-45.
- Barton, H; Creech, J; Godin, C; Randall, G; Seckel, C. (1995). Chloroethylene mixtures: Pharmacokinetic modeling and in vitro metabolism of vinyl chloride, trichloroethylene, and trans-1,2-dichloroethylene in rat. *Toxicol Appl Pharmacol* 130: 237-247.
<http://dx.doi.org/10.1006/taap.1995.1029>.
- Barton, H and Das, S. (1996). Alternatives for a risk assessment on chronic noncancer effects from oral exposure to trichloroethylene. *Regul Toxicol Pharmacol* 24: 269-285.
<http://dx.doi.org/10.1006/rtpb.1996.0140>.
- Barton, H; Flemming, C; Lipscomb, J. (1996). Evaluating human variability in chemical risk assessment: hazard identification and dose-response assessment for noncancer oral toxicity of trichloroethylene. *Toxicology* 111: 271-287.
- Barton, H; Bull, R; Schultz, I; Andersen, M. (1999). Dichloroacetate (DCA) dosimetry: Interpreting DCA-induced liver cancer dose response and the potential for DCA to contribute to trichloroethylene-induced liver cancer. *Toxicol Lett* 106: 9-21.

- Barton, H and Clewell, H, III. (2000). Evaluating noncancer effects of trichloroethylene: Dosimetry, mode of action, and risk assessment. Environ Health Perspect 108: 323-334.
- Barton, H; Chiu, W; Setzer, R; Andersen, M; Bailer, A; Bois, F; DeWoskin, R; Hays, S; Johanson, G; Jones, N; Loizou, G; Macphail, R; Portier, C; Spendiff, M; Tan, Y. (2007). Characterizing uncertainty and variability in physiologically-based pharmacokinetic (PBPK) models: State of the science and needs for research and implementation. Toxicol Sci 99: 395-402.
<http://dx.doi.org/10.1093/toxsci/kfm100>.
- Bartonicek, V. (1962). Metabolism and excretion of trichloroethylene after inhalation by human subjects. Occup Environ Med 19: 134-141. <http://dx.doi.org/10.1136/oem.19.2.134>.
- Bartonicek, V and Teisinger, J. (1962). Effect of tetraethyl thiuram disulphide (disulfiram) on metabolism of trichloroethylene in man. Occup Environ Med 19: 216-221.
<http://dx.doi.org/10.1136/oem.19.3.216>.
- Bartosiewicz, M; Penn, S; Buckpitt, A. (2001). Applications of gene arrays in environmental toxicology: Fingerprints of gene regulation associated with cadmium chloride, benzo(a)pyrene, and trichloroethylene. Environ Health Perspect 109: 71-74.
- Bartsch, H; Malaveille, C; Barbin, A; Planche, G. (1979). Mutagenic and alkylating metabolites of haloethylenes, chlorobutadienes and dichlorobutenes produced by rodent or human liver tissues: Evidence for oxirane formation by P450-linked microsomal mono-oxygenases. Arch Toxicol 41: 249-277. <http://dx.doi.org/10.1007/BF00296896>.
- Beaudreuil, S; Lasfargues, G; Laueriere, L; El Ghoul, Z; Fourquet, F; Longuet, C; Halimi, J; Nivet, H; Buchler, M. (2005). Occupational exposure in ANCA-positive patients: A case-control study. Kidney Int 67: 1961-1966. <http://dx.doi.org/10.1111/j.1523-1755.2005.00295.x>.
- Beaumont, J and Breslow, N. (1981). Power considerations in epidemiologic studies of vinyl chloride workers. Am J Epidemiol 114: 725-734.
- Becker, N; Deeg, E; Nieters, A. (2004). Population-based study of lymphoma in Germany: Rationale, study design and first results. Leuk Res 28: 713-724.
<http://dx.doi.org/10.1016/j.leukres.2003.11.010>.
- Beckstead, M; Weiner, J; Eger, E; Gong, D; Mihic, S. (2000). Glycine and [gamma]-aminobutyric acid(A) receptor function is enhanced by inhaled drugs of abuse. Mol Pharmacol 57: 1199-1205.
- Beekman, J; Boess, F; Hildebrand, H; Kalkuhl, A; Suter, L. (2006). Gene expression analysis of the hepatotoxicant methapyrilene in primary rat hepatocytes: An interlaboratory study. Environ Health Perspect 114: 92-99.
- Begg, C and Mazumdar, M. (1994). Operating characteristics of a rank correlation test for publication bias. Biometrics 50: 1088-1101.
- BEIR. ([Committee on the] Biological Effects of Ionizing Radiation). (1988). Health risks of radon and other internally deposited alpha-emitters: BEIR IV. (PB88-225594). Washington, DC: National Academy Press. <http://www.nap.edu/openbook.php?isbn=0309037972>.
- Beland, F. (1999). NTP technical report on the toxicity and metabolism studies of chloral hydrate (CAS No. 302-17-0). Administered by gavage to F344/N rats and B6C3F1 mice. In Toxicity Report Series. U.S. Dept. of Health and Human Services, National Institutes of Health: Rockville, MD.
- Benedetti, M; Whomsley, R; Canning, M. (2007). Drug metabolism in the paediatric population and in the elderly. Drug Discov Today 12: 599-610. <http://dx.doi.org/10.1016/j.drudis.2007.06.011>.
- Benichou, J; Chow, W; McLaughlin, J; Mandel, J; Fraumeni, J. (1998). Population attributable risk of renal cell cancer in Minnesota. Am J Epidemiol 148: 424-430.
- Benson, R (2004) [Correspondence from Robert Benson (U. S. EPA) to Dr. P. D. Johnson (University of Arizona)].
- Beppu, K. (1968). Transmission of the anesthetic agents through the placenta in painless delivery and their effects on newborn infants. Keio J Med 17: 81-107.
- Berger, J and Daxenbichler, G. (2002). DNA methylation of nuclear receptor genes - Possible role in malignancy. J Steroid Biochem Mol Biol 80: 1-11. [http://dx.doi.org/10.1016/S0960-0760\(01\)00179-0](http://dx.doi.org/10.1016/S0960-0760(01)00179-0).

- Berger, T and Horner, C. (2003). In vivo exposure of female rats to toxicants may affect oocyte quality. *Reprod Toxicol* 17: 273-281. [http://dx.doi.org/10.1016/S0890-6238\(03\)00009-1](http://dx.doi.org/10.1016/S0890-6238(03)00009-1).
- Bergeron, A; Jorquera, R; Tanguay, R. (2003). [Hereditary tyrosinemia: An endoplasmic reticulum stress disorder?]. *Med Sci (Paris)* 19: 976-980. <http://dx.doi.org/10.1051/medsci/20031910976>.
- Bergman, K. (1983). Interactions of trichloroethylene with DNA in vitro and with RNA and DNA of various mouse tissues in vivo. *Arch Toxicol* 54: 181-193.
- Berman, E; Schlicht, M; Moser, V; MacPhail, R. (1995). A multidisciplinary approach to toxicological screening: I. Systemic toxicity. *J Toxicol Environ Health* 45: 127-143. <http://dx.doi.org/10.1080/15287399509531986>.
- Bernad, P; Newell, S; Spyker, D. (1987). Neurotoxicity and behavior abnormalities in a cohort chronically exposed to trichloroethylene. *Vet Hum Toxicol* 29: 475.
- Bernauer, U; Birner, G; Dekant, W; Henschler, D. (1996). Biotransformation of trichloroethene: Dose-dependent excretion of 2,2,2-trichloro-metabolites and mercapturic acids in rats and humans after inhalation. *Arch Toxicol* 70: 338-346. <http://dx.doi.org/10.1007/s002040050283>.
- Bernillon, P and Bois, F. (2000). Statistical issues in toxicokinetic modeling: A bayesian perspective. *Environ Health Perspect* 108: 883-893.
- Béroud, C; Joly, D; Gallou, C; Staroz, F; Orfanelli, M; Junien, C. (1998). Software and database for the analysis of mutations in the VHL gene. *Nucleic Acids Res* 26: 256-258.
- Besson, H; Brennan, P; Becker, N; Nieters, A; De Sanjosé, S; Font, R; Maynadié, M; Foretova, L; Cocco, P; Staines, A; Vornanen, M; Boffetta, P. (2006). Tobacco smoking, alcohol drinking and non-Hodgkin's lymphoma: A European multicenter case-control study (Epilymph). *Int J Cancer* 119: 901-908. <http://dx.doi.org/10.1002/ijc.21913>.
- Best, D and Coleman, W. (2007). Treatment with 2-AAF blocks the small hepatocyte-like progenitor cell response in retrorsine-exposed rats. *J Hepatol* 46: 1055-1063. <http://dx.doi.org/10.1016/j.jhep.2007.01.040>.
- Beyer, R; Fry, R; Lasarev, M; McConnachie, L; Meira, L; Palmer, V; Powell, C; Ross, P; Bammler, T; Bradford, B; Cranson, A; Cunningham, M; Fannin, R; Higgins, G; Hurban, P; Kayton, R; Kerr, K; Kosyk, O; Lobenhofer, E; Sieber, S; Vliet, P; Weis, B; Wolfinger, R; Woods, C; Freedman, J; Linney, E; Kaufmann, W; Kavanagh, T; Paules, R; Rusyn, I; Samson, L; Spencer, P; Suk, W; Tennant, R; Zarbl, H; Consortium, MotTR. (2007). Multicenter study of acetaminophen hepatotoxicity reveals the importance of biological endpoints in genomic analyses. *Toxicol Sci* 99: 326-337. <http://dx.doi.org/10.1093/toxsci/kfm150>.
- Bhunya, S and Behera, B. (1987). Relative genotoxicity of trichloroacetic acid (TCA) as revealed by different cytogenetic assays: Bone marrow chromosome aberration, micronucleus and sperm-head abnormality in the mouse. *Mutat Res Genet Toxicol Environ Mutagen* 188: 215-221. [http://dx.doi.org/10.1016/0165-1218\(87\)90092-9](http://dx.doi.org/10.1016/0165-1218(87)90092-9).
- Bhunya, S and Jena, G. (1996). The evaluation of clastogenic potential of trichloroacetic acid (TCA) in chick in vivo test system. *Mutat Res Genet Toxicol Environ Mutagen* 367: 253-259. [http://dx.doi.org/10.1016/S0165-1218\(96\)90085-3](http://dx.doi.org/10.1016/S0165-1218(96)90085-3).
- Bignami, M; Conti, G; Conti, L; Crebelli, R; Misuraca, F; Puglia, A; Randazzo, R; Sciandrello, G; Carere, A. (1980). Mutagenicity of halogenated aliphatic hydrocarbons in *Salmonella typhimurium*, *Streptomyces coelicolor* and *Aspergillus nidulans*. *Chem Biol Interact* 30: 9-23. [http://dx.doi.org/10.1016/0009-2797\(80\)90110-6](http://dx.doi.org/10.1016/0009-2797(80)90110-6).
- Binaschi, S and Cantu, L. (1983). Vestibular function in solvent exposure: Clinical criteria. *Adv Biosci* 46: 205-210.
- Bingham, E; Cohrssen, B; Powell, C. (2001). *Patty's toxicology* (5 ed. Vol. 1-9). Hoboken, NJ: John Wiley & Sons.
- Birner, G; Vamvakas, S; Dekant, W; Henschler, D. (1993). Nephrotoxic and genotoxic N-acetyl-S-dichlorovinyl-L-cysteine is a urinary metabolite after occupational 1,1,2-trichloroethene exposure in humans: Implications for the risk of trichloroethene exposure. *Environ Health Perspect* 99: 281-284.

- Birner, G; Richling, C; Henschler, D; Anders, M; Dekant, W. (1994). Metabolism of tetrachloroethene in rats: Identification of N[epsilon]-(dichloroacetyl)-L-lysine and N[epsilon]-(trichloroacetyl)-L-lysine as protein adducts. *Chem Res Toxicol* 7: 724-732. <http://dx.doi.org/10.1021/tx00042a003>.
- Birner, G; Bernauer, U; Werner, M; Dekant, W. (1997). Biotransformation, excretion and nephrotoxicity of haloalkene-derived cysteine S-conjugates. *Arch Toxicol* 72: 1-8. <http://dx.doi.org/10.1007/s002040050461>.
- Birner, G; Werner, M; Rosner, E; Mehler, C; Dekant, W. (1998). Biotransformation, excretion, and nephrotoxicity of the hexachlorobutadiene metabolite (E)-N-acetyl-S-(1,2,3,4, 4-pentachlorobutadienyl)-L-cysteine sulfoxide. *Chem Res Toxicol* 11: 750-757. <http://dx.doi.org/10.1021/tx970216n>.
- Bjornsson, H; Sigurdsson, M; Fallin, M; Irizarry, R; Aspelund, T; Cui, H; Yu, W; Rongione, M; Ekström, T; Harris, T; Launer, L; Eiriksdottir, G; Leppert, M; Sapienza, C; Gudnason, V; Feinberg, A. (2008). Intra-individual change over time in DNA methylation with familial clustering. *JAMA* 299: 2877-2883. <http://dx.doi.org/10.1001/jama.299.24.2877>.
- Blackburn, A; Tzeng, H; Anders, M; Board, P. (2000). Discovery of a functional polymorphism in human glutathione transferase zeta by expressed sequence tag database analysis. *Pharmacogenetics* 10: 49-57.
- Blackburn, A; Coggan, M; Tzeng, H; Lantum, H; Polekhina, G; Parker, M; Anders, M; Board, P. (2001). GSTZ1d: A new allele of glutathione transferase zeta and maleylacetoacetate isomerase. *Pharmacogenetics* 11: 671-678.
- Blackshaw, A. (1970). Histochemical localization of testicular enzyme. In AD Johnson, WR Gomes & NL VanDemark (Eds.), *The testis* (pp. 73-173). New York, NY: Academic Press.
- Blain, L; Lachapelle, P; Molotchnikoff, S. (1994). Electoretinal responses are modified by chronic exposure to trichloroethylene. *Neurotoxicology* 15: 627-631.
- Blair, A; Haas, T; Prosser, R; Morrissette, M; Blackman, K; Grauman, D; Dusen, P; Moran, F. (1989). Mortality among United States Coast Guard marine inspectors. *Arch Environ Health* 44: 150-156. <http://dx.doi.org/10.1080/00039896.1989.9935879>.
- Blair, A; Linos, A; Stewart, P; Burmeister, L; Gibson, R; Everett, G; Schuman, L; Cantor, K. (1993). Evaluation of risks for non-Hodgkin's lymphoma by occupation and industry exposures from a case-control study. *Am J Ind Med* 23: 301-312. <http://dx.doi.org/10.1002/ajim.4700230207>.
- Blair, A; Hartge, P; Stewart, P; McAdams, M; Lubin, J. (1998). Mortality and cancer incidence of aircraft maintenance workers exposed to trichloroethylene and other organic solvents and chemicals: Extended follow-up. *Occup Environ Med* 55: 161-171. <http://dx.doi.org/10.1136/oem.55.3.161>.
- Blake, J; Compton, K; Schmidt, W; Orrego, H. (1988). Accuracy of death certificates in the diagnosis of alcoholic liver cirrhosis. *Alcohol Clin Exp Res* 12: 168-172.
- Blake, M; Castro, L; Leeder, J; Kearns, G. (2005). Ontogeny of drug metabolizing enzymes in the neonate. *Semin Fetal Neonatal Med* 10: 123-138. <http://dx.doi.org/10.1016/j.siny.2004.11.001>.
- Blatter, B; Roeleveld, N; Zielhuis, G; Verbeek, A. (1997). Assessment of occupational exposure in a population based case-control study: Comparing postal questionnaires with personal interviews. *Occup Environ Med* 54: 54-59.
- Blechacz, B and Gores, G. (2008). Cholangiocarcinoma. *Clin Liver Dis* 12: 131-150, ix. <http://dx.doi.org/10.1016/j.cld.2007.11.003>.
- Bloemen, L and Tomenson, J. (1995). Increased incidence of renal cell tumours in a cohort of cardboard workers exposed to trichloroethylene. *Arch Toxicol* 70: 129-130. <http://dx.doi.org/10.1007/BF02733675>.
- Bloemen, L; Monster, A; Kezic, S; Commandeur, J; Veulemans, H; Vermeulen, N; Wilmer, J. (2001). Study on the cytochrome P-450- and glutathione-dependent biotransformation of trichloroethylene in humans. *Int Arch Occup Environ Health* 74: 102-108. <http://dx.doi.org/10.1007/s004200000198>.
- Blossom, S; Pumford, N; Gilbert, K. (2004). Activation and attenuation of apoptosis of CD4+ T cells following in vivo exposure to two common environmental toxicants, trichloroacetaldehyde

- hydrate and trichloroacetic acid. *J Autoimmun* 23: 211-220. <http://dx.doi.org/10.1016/j.jaut.2004.06.007>.
- Blossom, S and Gilbert, K. (2006). Exposure to a metabolite of the environmental toxicant, trichloroethylene, attenuates CD4+ T cell activation-induced cell death by metalloproteinase-dependent FasL shedding. *Toxicol Sci* 92: 103-114. <http://dx.doi.org/10.1093/toxsci/kfj212>.
- Blossom, S and Doss, J. (2007). Trichloroethylene alters central and peripheral immune function in autoimmune-prone MRL(++) mice following continuous developmental and early life exposure. *J Immunotoxicol* 4: 129-141. <http://dx.doi.org/10.1080/15476910701337035>.
- Blossom, S; Doss, J; Gilbert, K. (2007). Chronic exposure to a trichloroethylene metabolite in autoimmune-prone MRL+/+ mice promotes immune modulation and alopecia. *Toxicol Sci* 95: 401-411. <http://dx.doi.org/10.1093/toxsci/kfl149>.
- Blossom, S; Doss, J; Hennings, L; Jernigan, S; Melnyk, S; James, S. (2008). Developmental exposure to trichloroethylene promotes CD4+ T cell differentiation and hyperactivity in association with oxidative stress and neurobehavioral deficits in MRL+/+ mice. *Toxicol Appl Pharmacol* 231: 344-353. <http://dx.doi.org/10.1016/j.taap.2008.05.009>.
- Board, P; Chelvanayagam, G; Jermiin, L; Tetlow, N; Tzeng, H; Anders, M; Blackburn, A. (2001). Identification of novel glutathione transferases and polymorphic variants by expressed sequence tag database analysis. *Drug Metab Dispos* 29: 544-547.
- Boerrigter, M. (2004). Mutagenicity of the peroxisome proliferators clofibrate, Wyeth 14,643 and di-2-ethylhexyl phthalate in the lacZ plasmid-based transgenic mouse mutation assay. *Journal of Carcinogenesis* 3: 7. <http://dx.doi.org/10.1186/1477-3163-3-7>.
- Boffetta, P and de Vocht, F. (2007). Occupation and the risk of non-Hodgkin lymphoma. *Cancer Epidemiol Biomarkers Prev* 16: 369-372. <http://dx.doi.org/10.1158/1055-9965.EPI-06-1055>.
- Bogen, K. (1990). Uncertainty in environmental health risk assessment. New York: Garland Publishing Company.
- Bogen, K; Colston, B, Jr; Machicao, L. (1992). Dermal absorption of dilute aqueous chloroform, trichloroethylene, and tetrachloroethylene in hairless guinea pigs. *Toxicol Sci* 18: 30-39. [http://dx.doi.org/10.1016/0272-0590\(92\)90192-K](http://dx.doi.org/10.1016/0272-0590(92)90192-K).
- Boice, J; Cohen, S; Mumma, M; Dupree Ellis, E; Eckerman, K; Leggett, R; Boecker, B; Brill, A; Henderson, B. (2006a). Mortality among radiation workers at Rocketdyne (Atomics International), 1948-1999. *Radiat Res* 166: 98-115. <http://dx.doi.org/10.1667/RR3582.1>.
- Boice, J; Marano, D; Cohen, S; Mumma, M; Blot, W; Brill, A; Fryzek, J; Henderson, B; McLaughlin, J. (2006b). Mortality among Rocketdyne workers who tested rocket engines, 1948-1999. *J Occup Environ Med* 48: 1070-1092. <http://dx.doi.org/10.1097/01.jom.0000240661.33413.b5>.
- Boice, J, Jr; Marano, D; Fryzek, J; Sadler, C; McLaughlin, J. (1999). Mortality among aircraft manufacturing workers. *Occup Environ Med* 56: 581-597. <http://dx.doi.org/10.1136/oem.56.9.581>.
- Bois, F. (2000a). Statistical analysis of Fisher et al. PBPK model of trichloroethylene kinetics. *Environ Health Perspect* 108: 275-282.
- Bois, F. (2000b). Statistical analysis of Clewell et al. PBPK model of trichloroethylene kinetics. *Environ Health Perspect* 108: 307-316.
- Bolm-Audorff, U; Ahrens, W; Jockel, K-H; Molik, B; Greiser, E; Timm, J. (1988). Experience with supplementary questionnaires in a lung cancer case-referent study. In D Henon & M Goldberg (Eds.), *Proceedings of the workshop on methodology of assessment of occupational exposures in the context of epidemiological detection of cancer risks* (pp. 117-168). Paris, France: Commission of the European Communities.
- Bolt, H and Filser, J. (1977). Irreversible binding of chlorinated ethylenes to macromolecules. *Environ Health Perspect* 21: 107-112.
- Bolt, H; Lammert, M; Selinski, S; Brüning, T. (2004). Urinary alpha1-microglobulin excretion as biomarker of renal toxicity in trichloroethylene-exposed persons. *Int Arch Occup Environ Health* 77: 186-190. <http://dx.doi.org/10.1007/s00420-003-0500-3>.

- Bombail, V; Moggs, J; Orphanides, G. (2004). Perturbation of epigenetic status by toxicants. *Toxicol Lett* 149: 51-58. <http://dx.doi.org/10.1016/j.toxlet.2004.01.003>.
- Bonatti, S; Cavalieri, Z; Viaggi, S; Abbondandolo, A. (1992). The analysis of 10 potential spindle poisons for their ability to induce CREST-positive micronuclei in human diploid fibroblasts. *Mutagenesis* 7: 111-114. <http://dx.doi.org/10.1093/mutage/7.2.111>.
- Boorman, G. (1999). Drinking water disinfection byproducts: Review and approach to toxicity evaluation. *Environ Health Perspect* 107: 207-217.
- Borzio, M; Fargion, S; Borzio, F; Fracanzani, A; Croce, A; Stroffolini, T; Oldani, S; Cotichini, R; Roncalli, M. (2003). Impact of large regenerative, low grade and high grade dysplastic nodules in hepatocellular carcinoma development. *J Hepatol* 39: 208-214. [http://dx.doi.org/10.1016/S0168-8278\(03\)00190-9](http://dx.doi.org/10.1016/S0168-8278(03)00190-9).
- Boublik, T; Fried, V; Hala, E. (1984). The vapour pressures of pure substances: selected values of the temperature dependence of the vapour pressures of some pure substances in the normal and low pressure region. Amsterdam, The Netherlands: Elsevier.
- Bove, F; Fulcomer, M; Klotz, J; Esmar, J; Dufficy, E; Savrin, J. (1995). Public drinking water contamination and birth outcomes. *Am J Epidemiol* 141: 850-862.
- Bove, F. (1996). Public drinking water contamination and birthweight, prematurity, fetal deaths, and birth defects. *Toxicol Ind Health* 12: 255-266.
- Bovenzi, M; Barbone, F; Pisa, F; Betta, A; Romeo, L; Tonello, A; Biasi, D; Caramaschi, P. (2004). A case-control study of occupational exposures and systemic sclerosis. *Int Arch Occup Environ Health* 77: 10-16. <http://dx.doi.org/10.1007/s00420-003-0462-5>.
- Bovo, R; Aimoni, C; Martini, A. (2006). Immune-mediated inner ear disease. *Acta Otolaryngol* 126: 1012-1021. <http://dx.doi.org/10.1080/00016480600606723>.
- Bowler, R; Mergler, D; Huel, G; Harrison, R; Cone, J. (1991). Neuropsychological impairment among former microelectronics workers. *Neurotoxicology* 12: 87-103.
- Box, G; Hunter, W; Hunter, J. (1978). Statistics for experimenters: an introduction to design, data analysis, and model building. New York, NY: Wiley.
- Boyault, S; Rickman, D; de Reyniés, A; Balabaud, C; Rebouissou, S; Jeannot, E; Herault, A; Saric, J; Belghiti, J; Franco, D; Bioulac-Sage, P; Laurent-Puig, P; Zucman-Rossi, J. (2007). Transcriptome classification of HCC is related to gene alterations and to new therapeutic targets. *Hepatology* 45: 42-52. <http://dx.doi.org/10.1002/hep.21467>.
- Boyer, A; Finch, W; Runyan, R. (2000). Trichloroethylene inhibits development of embryonic heart valve precursors in vitro. *Toxicol Sci* 53: 109-117.
- Boyes, W; Bushnell, P; Crofton, K; Evans, M; Simmons, J. (2000). Neurotoxic and pharmacokinetic responses to trichloroethylene as a function of exposure scenario. *Environ Health Perspect* 108: 317-322.
- Boyes, W; Bercegeay, M; Ali, J; Krantz, T; McGee, J; Evans, M; Raymer, J; Bushnell, P; Simmons, J. (2003). Dose-based duration adjustments for the effects of inhaled trichloroethylene on rat visual function. *Toxicol Sci* 76: 121-130. <http://dx.doi.org/10.1093/toxsci/kfg213>.
- Boyes, W; Bercegeay, M; Krantz, T; Evans, M; Benignus, V; Simmons, J. (2005a). Momentary brain concentration of trichloroethylene predicts the effects on rat visual function. *Toxicol Sci* 87: 187-196. <http://dx.doi.org/10.1093/toxsci/kfi242>.
- Boyes, W; Simmons, J; Eklund, C; Benignus, V; Janssen, P; Bushnell, P. (2005b). Applications of dosimetry modeling to assessment of neurotoxic risk. *Environ Toxicol Pharmacol* 19: 599 - 605.
- Boyle, C and Brann, E. (1992). Proxy respondents and the validity of occupational and other exposure data. The Selected Cancers Cooperative Study Group. *Am J Epidemiol* 136: 712-721.
- Bradford, B; Kono, H; Isayama, F; Kosyk, O; Wheeler, M; Akiyama, T; Bleye, L; Krausz, K; Gonzalez, F; Koop, D; Rusyn, I. (2005). Cytochrome P450 CYP2E1, but not nicotinamide adenine dinucleotide phosphate oxidase, is required for ethanol-induced oxidative DNA damage in rodent liver. *Hepatology* 41: 336-344. <http://dx.doi.org/10.1002/hep.20532>.

- Braissant, O and Wahli, W. (1998). Differential expression of peroxisome proliferator-activated receptor-alpha, -beta, and -gamma during rat embryonic development. *Endocrinology* 139: 2748-2754.
- Branda, M and Wands, J. (2006). Signal transduction cascades and hepatitis B and C related hepatocellular carcinoma. *Hepatology* 43: 891-902. <http://dx.doi.org/10.1002/hep.21196>.
- Brauch, H; Weirich, G; Hornauer, M; Störkel, S; Wöhl, T; Brüning, T. (1999). Trichloroethylene exposure and specific somatic mutations in patients with renal cell carcinoma. *J Natl Cancer Inst* 91: 854-861.
- Brauch, H; Weirich, G; Klein, B; Rabstein, S; Bolt, H; Brüning, T. (2004). VHL mutations in renal cell cancer: does occupational exposure to trichloroethylene make a difference. *Toxicol Lett* 151: 301-310. <http://dx.doi.org/10.1016/j.toxlet.2003.12.074>.
- Brautbar, N and Williams, J, II. (2002). Industrial solvents and liver toxicity: Risk assessment, risk factors and mechanisms. *Int J Hyg Environ Health* 205: 479-491. <http://dx.doi.org/10.1078/1438-4639-00175>.
- Brenner, H and Gefeller, O. (1993). Use of the positive predictive value to correct for disease misclassification in epidemiologic studies. *Am J Epidemiol* 138: 1007-1015.
- Breslow, N and Day, N. (1980). Statistical methods in cancer research: Volume 1 - the analysis of case-control studies (Vol. 32). Lyon, France: International Agency for Research on Cancer.
- Breslow, NE and Day, NE (Eds.). (1987). Statistical methods in cancer research. Volume 2 - The design and analysis of cohort studies (Vol. 82). Lyon, France: International Agency for Research on Cancer.
- Bringmann, G; Friedrich, H; Feineis, D. (1992). Trichloroharmanes as potential endogenously formed inducers of Morbus Parkinson: synthesis, analytics, and first in vivo-investigations. *J Neural Transm Suppl* 38: 15-26.
- Bringmann, G; God, R; Feineis, D; Wesemann, W; Riederer, P; Rausch, W; Reichmann, H; Sontag, K. (1995). The TaClo concept: 1-trichloromethyl-1,2,3,4-tetrahydro-beta-carboline (TaClo), a new toxin for dopaminergic neurons. *J Neural Transm Suppl* 46: 235-244.
- Briving, C; Jacobson, I; Hamberger, A; Kjellstrand, P; Haglid, K; Rosengren, L. (1986). Chronic effects of perchloroethylene and trichloroethylene on the gerbil brain amino acids and glutathione. *Neurotoxicology* 7: 101-108.
- Brodsky, W and Uryvaeva, I. (1977). Cell polyploidy: Its relation to tissue growth and function. *Int Rev Cytol* 50: 275-332. [http://dx.doi.org/10.1016/S0074-7696\(08\)60100-X](http://dx.doi.org/10.1016/S0074-7696(08)60100-X).
- Bronley-DeLancey, A; McMillan, D; McMillan, J; Jollow, D; Mohr, L; Hoel, D. (2006). Application of cryopreserved human hepatocytes in trichloroethylene risk assessment: Relative disposition of chloral hydrate to trichloroacetate and trichloroethanol. *Environ Health Perspect* 114: 1237-1242.
- Bronzetti, G; Corsi, C; Nieri, R. (1980). [Use of the D7 strain of *S. cerevisiae* in the determination of environmental risk. 1. Genetic effects of trichloroethylene]. *Boll Soc Ital Biol Sper* 56: 1315-1321.
- Bronzetti, G; Galli, A; Corsi, C; Cundari, E; Del Carratore, R; Nieri, R; Paolini, M. (1984). Genetic and biochemical investigation on chloral hydrate in vitro and in vivo. *Mutat Res* 141: 19-22. [http://dx.doi.org/10.1016/0165-7992\(84\)90031-9](http://dx.doi.org/10.1016/0165-7992(84)90031-9).
- Bross, G; DiFrancesco, D; Desmond, M. (1983). The effects of low dosages of trichloroethylene on chick development. *Toxicology* 28: 283-294.
- Brown, H; Bishop, D; Rowan, C. (1984). The role of skin absorption as a route of exposure for volatile organic compounds (VOCs) in drinking water. *Am J Public Health* 74: 479-484.
- Brown, R; Delp, M; Lindstedt, S; Rhomberg, L; Beliles, R. (1997). Physiological parameter values for physiologically based pharmacokinetic models. *Toxicol Ind Health* 13: 407-484. <http://dx.doi.org/10.1177/074823379701300401>.
- Brownson, R. (1988). A case-control study of renal cell carcinoma in relation to occupation, smoking, and alcohol consumption. *Arch Environ Health* 43: 238-241.
- Bruckner J, unpublished dataset

- Brüning, T; Lammert, M; Kempkes, M; Thier, R; Golka, K; Bolt, H. (1997a). Influence of polymorphisms of GSTM1 and GSTT1 for risk of renal cell cancer in workers with long-term high occupational exposure to trichloroethene. *Arch Toxicol* 71: 596-599.
- Brüning, T; Weirich, G; Hornauer, M; Höfler, H; Brauch, H. (1997b). Renal cell carcinomas in trichloroethene (TCE) exposed persons are associated with somatic mutations in the von Hippel-Lindau (VHL) tumour suppressor gene. *Arch Toxicol* 71: 332-335.
<http://dx.doi.org/10.1007/s002040050394>.
- Brüning, T; Vamvakas, S; Makropoulos, V; Birner, G. (1998). Acute intoxication with trichloroethene: Clinical symptoms, toxicokinetics, metabolism, and development of biochemical parameters for renal damage. *Toxicol Sci* 41: 157-165. <http://dx.doi.org/10.1006/toxs.1997.2401>.
- Brüning, T; Mann, H; Melzer, H; Sundberg, A; Bolt, H. (1999a). Pathological excretion patterns of urinary proteins in renal cell cancer patients exposed to trichloroethylene. *Occup Med (Lond)* 49: 299-305.
- Brüning, T; Sundberg, A; Birner, G; Lammert, M; Bolt, H; Appelkvist, E; Nilsson, R; Dallner, G. (1999b). Glutathione transferase alpha as a marker for tubular damage after trichloroethylene exposure. *Arch Toxicol* 73: 246-254.
- Brüning, T; Pesch, B; Wiesenhütter, B; Rabstein, S; Lammert, M; Baumüller, A; Bolt, H. (2003). Renal cell cancer risk and occupational exposure to trichloroethylene: Results of a consecutive case-control study in Arnsberg, Germany. *Am J Ind Med* 43: 274-285.
<http://dx.doi.org/10.1002/ajim.10185>.
- Brzezinski, M; Boutelet-Bochan, H; Person, R; Fantel, A; Juchau, M. (1999). Catalytic activity and quantitation of cytochrome P-450 2E1 in prenatal human brain. *J Pharmacol Exp Ther* 289: 1648-1653.
- Buben, J and O'Flaherty, E. (1985). Delineation of the role of metabolism in the hepatotoxicity of trichloroethylene and perchloroethylene: A dose-effect study. *Toxicol Appl Pharmacol* 78: 105-122.
- Buckpitt, A; Chang, A-M; Weir, A; Van Winkle, L; Duan, X; Philpot, R; Plopper, C. (1995). Relationship of cytochrome P450 activity to Clara cell cytotoxicity IV Metabolism of naphthalene and naphthalene oxide in microdissected airways from mice, rats, and hamsters. *Mol Pharmacol* 47: 74-81.
- Budavari, S. (Ed.). (1996). *The Merck Index: An Encyclopedia of Chemicals, Drugs, and Biologicals* (12th ed.). Rahway, NJ: Merck & Co., Inc.
- Bull, R; Sanchez, I; Nelson, M; Larson, J; Lansing, A. (1990). Liver tumor induction in B6C3F1 mice by dichloroacetate and trichloroacetate. *Toxicology* 63: 341-359. [http://dx.doi.org/10.1016/0300-483X\(90\)90195-M](http://dx.doi.org/10.1016/0300-483X(90)90195-M).
- Bull, R. (2000). Mode of action of liver tumor induction by trichloroethylene and its metabolites, trichloroacetate and dichloroacetate. *Environ Health Perspect* 108: 241-259.
- Bull, R; Orner, G; Cheng, R; Stillwell, L; Stauber, A; Sasser, L; Lingohr, M; Thrall, B. (2002). Contribution of dichloroacetate and trichloroacetate to liver tumor induction in mice by trichloroethylene. *Toxicol Appl Pharmacol* 182: 55-65. <http://dx.doi.org/10.1006/taap.2002.9427>.
- Bull, R. (2004a). Trichloroethylene and liver tumors in mice Proceedings of New Scientific Research Related to the Health Effects of Trichloroethylene. Washington, DC: U.S. Environmental Protection Agency.
- Bull, R. (2004b) Trichloroethylene and liver tumors in mice. Presentation at the U.S. EPA Symposium on New Scientific Research Related to Health Effects of TCE: U.S. Environmental Protection Agency.
- Bull, R; Sasser, L; Lei, X. (2004). Interactions in the tumor-promoting activity of carbon tetrachloride, trichloroacetate, and dichloroacetate in the liver of male B6C3F1 mice. *Toxicology* 199: 169-183.
<http://dx.doi.org/10.1016/j.tox.2004.02.018>.

- Bunce, N and Schneider, U. (1994). Chemical lifetimes of chlorinated aliphatic priority pollutants in the Canadian troposphere. *J Photochem Photobiol A* 81: 93 - 101. [http://dx.doi.org/10.1016/1010-6030\(94\)03781-7](http://dx.doi.org/10.1016/1010-6030(94)03781-7).
- Burg, J; Gist, G; Allred, S; Rudtke, T; Pallos, L; Cusack, C. (1995). The national exposure registry - morbidity analyses of noncancer outcomes from the trichloroethylene subregistry baseline data. *Int J Occup Med Toxicol* 4: 237 - 257.
- Burg, J and Gist, G. (1997). The potential impact on women from environmental exposures. *J Wom Health Gend Base Med* 6: 159-161.
- Burg, J and Gist, G. (1999). Health effects of environmental contaminant exposure: An intrafile comparison of the Trichloroethylene Subregistry. *Arch Environ Health* 54: 231-241. <http://dx.doi.org/10.1080/0003989909602480>.
- Burmaster, D. (1982). The new pollution: Groundwater contamination. *Environment* 24: 6-13;33-36.
- Bushnell, P. (1997). Concentration-time relationships for the effects of inhaled trichloroethylene on signal detection behavior in rats. *Fundam Appl Toxicol* 36: 30-38. <http://dx.doi.org/10.1093/toxsci/36.1.30>.
- Bushnell, P and Oshiro, W. (2000). Behavioral components of tolerance to repeated inhalation of trichloroethylene (TCE) in rats. *Neurotoxicol Teratol* 22: 221-229. [http://dx.doi.org/10.1016/S0892-0362\(99\)00078-1](http://dx.doi.org/10.1016/S0892-0362(99)00078-1).
- Butler, T. (1949). Metabolic transformations of trichloroethylene. *J Pharmacol Exp Ther* 97: 84-92.
- Buxton, P and Hayward, M. (1967). Polyneuritis cranialis associated with industrial trichloroethylene poisoning. *J Neurol Neurosurg Psychiatry* 30: 511-518.
- Byers, V; Levin, A; Ozonoff, D; Baldwin, R. (1988). Association between clinical symptoms and lymphocyte abnormalities in a population with chronic domestic exposure to industrial solvent-contaminated domestic water supply and a high incidence of leukaemia. *Cancer Immunol Immunother* 27: 77-81.
- Cai, H and Guengerich, F. (1999). Mechanisms of aqueous decomposition of trichloroethylene oxide. *J Am Chem Soc* 121: 11656-11663. <http://dx.doi.org/10.1021/ja9914240>.
- Cai, H and Guengerich, F. (2001a). Reaction of trichloroethylene oxide with proteins and dna: instability of adducts and modulation of functions. *Chem Res Toxicol* 14: 54-61.
- Cai, H and Guengerich, F. (2001b). Reaction of trichloroethylene and trichloroethylene oxide with cytochrome P450 enzymes: Inactivation and sites of modification. *Chem Res Toxicol* 14: 451-458. <http://dx.doi.org/10.1021/tx0002280>.
- Cai, P; König, R; Khan, M; Qiu, S; Kaphalia, B; Ansari, G. (2006). Autoimmune response in MRL+/+ mice following treatment with dichloroacetyl chloride or dichloroacetic anhydride. *Toxicol Appl Pharmacol* 216: 248-255. <http://dx.doi.org/10.1016/j.taap.2006.05.010>.
- Cai, P; König, R; Khan, M; Kaphalia, B; Ansari, G. (2007). Differential immune responses to albumin adducts of reactive intermediates of trichloroethene in MRL+/+ mice. *Toxicol Appl Pharmacol* 220: 278-283. <http://dx.doi.org/10.1016/j.taap.2007.01.020>.
- Cai, P; König, R; Boor, P; Kondraganti, S; Kaphalia, B; Khan, M; Ansari, G. (2008). Chronic exposure to trichloroethene causes early onset of SLE-like disease in female MRL +/+ mice. *Toxicol Appl Pharmacol* 228: 68-75. <http://dx.doi.org/10.1016/j.taap.2007.11.031>.
- Cakmak, A; Ekici, A; Ekici, M; Arslan, M; Iteginli, A; Kurtipek, E; Kara, T. (2004). Respiratory findings in gun factory workers exposed to solvents. *Respir Med* 98: 52-56.
- Caldwell, J and Keshava, N. (2006). Key issues in the modes of action and effects of trichloroethylene metabolites for liver and kidney tumorigenesis. *Environ Health Perspect* 114: 1457-1463. <http://dx.doi.org/10.1289/ehp.8692>.
- Caldwell, J; Jinot, J; Devoney, D; Gift, J. (2008a). Evaluation of evidence for infection as a mode of action for induction of rat lymphoma. *Environ Mol Mutagen* 49: 155-164.
- Caldwell, J; Keshava, N; Evans, M. (2008b). Difficulty of mode of action determination for trichloroethylene: An example of complex interactions of metabolites and other chemical exposures. *Environ Mol Mutagen* 49: 142-154. <http://dx.doi.org/10.1002/em.20350>.

- Caldwell, P; Thorne, P; Johnson, P; Boitano, S; Runyan, R; Selmin, O. (2008c). Trichloroethylene disrupts cardiac gene expression and calcium homeostasis in rat myocytes. *Toxicol Sci* 104: 135-143. <http://dx.doi.org/10.1093/toxsci/kfn078>.
- Caldwell, P; Manziello, A; Howard, J; Palbykin, B; Runyan, R; Selmin, O. (2010). Gene expression profiling in the fetal cardiac tissue after folate and low-dose trichloroethylene exposure. *Birth Defects Res A Clin Mol Teratol* 88: 111-127. <http://dx.doi.org/10.1002/bdra.20631>.
- Callen, D; Wolf, C; Philpot, R. (1980). Cytochrome P-450 mediated genetic activity and cytotoxicity of seven halogenated aliphatic hydrocarbons in *Saccharomyces cerevisiae*. *Mutat Res Genet Toxicol Environ Mutagen* 77: 55-63. [http://dx.doi.org/10.1016/0165-1218\(80\)90120-2](http://dx.doi.org/10.1016/0165-1218(80)90120-2).
- Calvisi, D; Factor, V; Ladu, S; Conner, E; Thorgeirsson, S. (2004a). Disruption of beta-catenin pathway or genomic instability define two distinct categories of liver cancer in transgenic mice. *Gastroenterology* 126: 1374-1386. <http://dx.doi.org/10.1053/j.gastro.2004.02.014>.
- Calvisi, D; Ladu, S; Conner, E; Factor, V; Thorgeirsson, S. (2004b). Disregulation of E-cadherin in transgenic mouse models of liver cancer. *Lab Invest* 84: 1137-1147. <http://dx.doi.org/10.1038/labinvest.3700147>.
- Calvisi, D; Ladu, S; Gorden, A; Farina, M; Conner, E; Lee, J-S; Factor, V; Thorgeirsson, S. (2006). Ubiquitous activation of Ras and Jak/Stat pathways in human HCC. *Gastroenterology* 130: 1117-1128. <http://dx.doi.org/10.1053/j.gastro.2006.01.006>.
- Calvisi, D; Ladu, S; Gorden, A; Farina, M; Lee, J-S; Conner, E; Schroeder, I; Factor, V; Thorgeirsson, S. (2007). Mechanistic and prognostic significance of aberrant methylation in the molecular pathogenesis of human hepatocellular carcinoma. *J Clin Invest* 117: 2713-2722. <http://dx.doi.org/10.1172/JCI31457>.
- Campo, P; Loquet, G; Blachère, V; Roure, M. (1999). Toluene and styrene intoxication route in the rat cochlea. *Neurotoxicol Teratol* 21: 427-434. [http://dx.doi.org/10.1016/S0892-0362\(99\)00010-0](http://dx.doi.org/10.1016/S0892-0362(99)00010-0).
- Campo, P; Blachère, V; Payan, J; Cossec, B; Ducos, P. (2006). Toluene toxicokinetics and metabolism parameters in the rat and guinea pig. *Environ Toxicol Pharmacol* 21: 276-282. <http://dx.doi.org/10.1016/j.etap.2005.09.004>.
- Campo, P; Maguin, K; Lataye, R. (2007). Effects of aromatic solvents on acoustic reflexes mediated by central auditory pathways. *Toxicol Sci* 99: 582-590. <http://dx.doi.org/10.1093/toxsci/kfm180>.
- Cao, K; Hollenbach, J; Shi, X; Shi, W; Chopek, M; Fernández-Viña, M. (2001). Analysis of the frequencies of HLA-A, B, and C alleles and haplotypes in the five major ethnic groups of the United States reveals high levels of diversity in these loci and contrasting distribution patterns in these populations. *Hum Immunol* 62: 1009-1030.
- Cappaert, N; Klis, S; Muijser, H; de Groot, J; Kulig, B; Smoorenburg, G. (1999). The ototoxic effects of ethyl benzene in rats. *Hear Res* 137: 91-102. [http://dx.doi.org/10.1016/S0378-5955\(99\)00141-0](http://dx.doi.org/10.1016/S0378-5955(99)00141-0).
- Cappaert, N; Klis, S; Baretta, A; Muijser, H; Smoorenburg, G. (2000). Ethyl benzene-induced ototoxicity in rats: A dose-dependent mid-frequency hearing loss. *J Assoc Res Otolaryngol* 1: 292-299. <http://dx.doi.org/10.1007/s101620010050>.
- Carney, E; Thorsrud, B; Dugard, P; Zablotny, C. (2006). Developmental toxicity studies in Crl:CD (SD) rats following inhalation exposure to trichloroethylene and perchloroethylene. *Birth Defects Res B Dev Reprod Toxicol* 77: 405-412. <http://dx.doi.org/10.1002/bdrb.20091>.
- Carpenter, S; Lasker, J; Raucy, J. (1996). Expression, induction, and catalytic activity of the ethanol-inducible cytochrome P450 (CYP2E1) in human fetal liver and hepatocytes. *Mol Pharmacol* 49: 260-268.
- Carrieri, M; Magosso, D; Piccoli, P; Zanetti, E; Trevisan, A; Bartolucci, G. (2007). Acute, nonfatal intoxication with trichloroethylene. *Arch Toxicol* 81: 529-532. <http://dx.doi.org/10.1007/s00204-007-0180-y>.
- Carter, J; Carter, H; DeAngelo, A. (1995). Biochemical, pathologic and morphometric alterations induced in male B6C3F1 mouse liver by short-term exposure to dichloroacetic acid. *Toxicol Lett* 81: 55-71. [http://dx.doi.org/10.1016/0378-4274\(95\)03409-9](http://dx.doi.org/10.1016/0378-4274(95)03409-9).

- Carter, J; Carter, H; Deddens, J; Hurst, B; George, M; DeAngelo, A. (2003). A 2-year dose-response study of lesion sequences during hepatocellular carcinogenesis in the male B6C3F(1) mouse given the drinking water chemical dichloroacetic acid. Environ Health Perspect 111: 53-64. <http://dx.doi.org/10.1289/ehp.5442>.
- CDC. (Centers for Disease Control and Prevention). (2007). National Vital Statistics Reports: Deaths: Final data for 2004. pp. 120. Hyattsville, MD. <http://www.cdc.gov/nchs/products/nvsr.htm>.
- CDHS. (California Department of Health Services). (1986). Organic chemical contamination of large public water systems in California. (86870001977). Sacramento, CA.
- Chakrabarti, S and Tuchweber, B. (1988). Studies of acute nephrotoxic potential of trichloroethylene in Fischer 344 rats. J Toxicol Environ Health 23: 147-158.
- Chalupa, B; Synková, J; Ševčík, M. (1960). The assessment of electroencephalographic changes and memory disturbances in acute intoxications with industrial poisons. Br J Ind Med 17: 238-241.
- Chang, L; Daniel, F; DeAngelo, A. (1992). Analysis of DNA strand breaks induced in rodent liver in vivo, hepatocytes in primary culture, and a human cell line by chlorinated acetic acids and chlorinated acetaldehydes. Environ Mol Mutagen 20: 277-288. <http://dx.doi.org/10.1002/em.2850200406>.
- Chang, Y-M; Tai, C-F; Yang, S-C; Chen, C; Shih, T; Lin, R; Liou, S-H. (2003). A cohort mortality study of workers exposed to chlorinated organic solvents in Taiwan. Ann Epidemiol 13: 652-660. [http://dx.doi.org/10.1016/S1047-2797\(03\)00038-3](http://dx.doi.org/10.1016/S1047-2797(03)00038-3).
- Chang, Y; Tai, C; Yang, S; Lin, R; Sung, F; Shih, T; Liou, S. (2005). Cancer incidence among workers potentially exposed to chlorinated solvents in an electronics factory. J Occup Health 47: 171-180.
- Channel, S; Latendresse, J; Kidney, J; Grabau, J; Lane, J; Steel-Goodwin, L; Gothaus, M. (1998). A subchronic exposure to trichloroethylene causes lipid peroxidation and hepatocellular proliferation in male B6C3F1 mouse liver. Toxicol Sci 43: 145-154. <http://dx.doi.org/10.1093/toxsci/43.2.145>.
- Chapin, R; Norton, R; Popp, J; Bus, J. (1982). The effects of 2,5-hexanedione on reproductive hormones and testicular enzyme activities in the F-344 rat. Toxicol Appl Pharmacol 62: 262-272.
- Chapin, R and Sloane, R. (1997). Reproductive assessment by continuous breeding: Evolving study design and summaries of ninety studies. Environ Health Perspect 105: 199-205.
- Charbonneau, M; Couture, J; Plaa, G. (1991). Inhalation versus oral administration of acetone: effect of the vehicle on the potentiation of CCl₄-induced liver injury. Toxicol Lett 57: 47-54.
- Charbotel, B; Fevotte, J; Hours, M; Martin, J; Bergeret, A. (2005). Case-control study on renal cell cancer and occupational trichloroethylene exposure, in the Arve Valley (France). Lyon, France: Institut Universitaire de Médecine du Travail, UMRESTTE, Université Claude Bernard. http://hal.archives-ouvertes.fr/docs/00/54/59/80/PDF/charbotel_octobre_05.pdf.
- Charbotel, B; Fevotte, J; Hours, M; Martin, J-L; Bergeret, A. (2006). Case-control study on renal cell cancer and occupational exposure to trichloroethylene. Part II: Epidemiological aspects. Ann Occup Hyg 50: 777-787. <http://dx.doi.org/10.1093/annhyg/mel039>.
- Charbotel, B; Gad, S; Caïola, D; Béroud, C; Fevotte, J; Bergeret, A; Ferlicot, S; Richard, S. (2007). Trichloroethylene exposure and somatic mutations of the VHL gene in patients with Renal Cell Carcinoma. J Occup Med Toxicol 2: 13. <http://dx.doi.org/10.1186/1745-6673-2-13>.
- Charbotel, B (2008) [E mail from Barbara Charbotel, University of Lyon, to Cheryl Scott, U.S. EPA].
- Charbotel, B; Fevotte, J; Martin, J-L; Bergeret, A. (2009). [Renal cell carcinoma and exposure to trichloroethylene: Are French occupational exposure limits relevant?]. Rev Epidemiol Sante Publique 57: 41-47. <http://dx.doi.org/10.1016/j.respe.2008.09.008>.
- Charles JM, Bond DM, Jeffries TK, Yano BL, Stott WT, Johnson KA, Cunny HC, Wilson RD, Bust JS. 1996. Chronic dietary toxicity/oncogenicity studies on 2,4-dichlorophenoxyacetic acid in rodents. Fundamental and applied toxicology 11:166-172. <http://www.ncbi.nlm.nih.gov/pubmed/8921335>.

- Charles, J and Leeming, N. (1998). Chronic dietary toxicity/oncogenicity studies on 2,4-dichlorophenoxybutyric acid in rodents. *Toxicol Sci* 46: 21-30. <http://dx.doi.org/10.1006/toxs.1998.2502>.
- Checkoway, H; Pearce, N; Crawford-Brown, D. (1989). Research methods in occupational epidemiology (Vol. 13). New York, NY: Oxford University Press.
- Chemical Marketing Reporter. (1997). Chemical Profile: Trichloroethylene. New York, NY: Schnell Pub Co.
- Chen, C. (2000). Biologically based dose-response model for liver tumors induced by trichloroethylene. *Environ Health Perspect* 108 Suppl 2: 335-342.
- Chen, C; Hennig, G; Whiteley, H; Manautou, J. (2002a). Protection against acetaminophen hepatotoxicity by clofibrate pretreatment: role of catalase induction. *J Biochem Mol Toxicol* 16: 227-234. <http://dx.doi.org/10.1002/jbt.10043>.
- Chen, X; Cheung, S; So, S; Fan, S; Barry, C; Higgins, J; Lai, K; Ji, J; Dudoit, S; Ng, I; Van De Rijn, M; Botstein, D; Brown, P. (2002b). Gene expression patterns in human liver cancers. *Mol Biol Cell* 13: 1929-1939. <http://dx.doi.org/10.1091/mbc.02-02-0023>.
- Chen, Y; Cai, J; Anders, M; Stevens, J; Jones, D. (2001). Role of mitochondrial dysfunction in S-(1,2-dichlorovinyl)-l-cysteine-induced apoptosis. *Toxicol Appl Pharmacol* 170: 172-180. <http://dx.doi.org/10.1006/taap.2000.9107>.
- Chen, Z; White, C; He, C; Liu, Y; Eaton, D. (1995). Zonal differences in DNA synthesis activity and cytochrome P450 gene expression in livers of male F344 rats treated with five nongenotoxic carcinogens. *J Environ Pathol Toxicol Oncol* 14: 83-99.
- Cherrie, J; Kromhout, H; Semple, S. (2001). The importance of reliable exposure estimates in deciding whether trichloroethylene can cause kidney cancer. *J Cancer Res Clin Oncol* 127: 400-404.
- Cheung, C; Akiyama, T; Ward, J; Nicol, C; Feigenbaum, L; Vinson, C; Gonzalez, F. (2004). Diminished hepatocellular proliferation in mice humanized for the nuclear receptor peroxisome proliferator-activated receptor alpha. *Cancer Res* 64: 3849-3854. <http://dx.doi.org/10.1158/0008-5472.CAN-04-0322>.
- Chia, S-E; Ong, C-N; Tsakok, M; Ho, A. (1996). Semen parameters in workers exposed to trichloroethylene. *Reprod Toxicol* 10: 295-299. [http://dx.doi.org/10.1016/0890-6238\(96\)00058-5](http://dx.doi.org/10.1016/0890-6238(96)00058-5).
- Chia, S; Goh, V; Ong, C. (1997). Endocrine profiles of male workers with exposure to trichloroethylene. *Am J Ind Med* 32: 217-222. [http://dx.doi.org/10.1002/\(SICI\)1097-0274\(199709\)32:3<217::AID-AJIM6>3.0.CO;2-U](http://dx.doi.org/10.1002/(SICI)1097-0274(199709)32:3<217::AID-AJIM6>3.0.CO;2-U).
- Chieco, P; Moslen, M; Reynolds, E. (1981). Effect of administrative vehicle on oral 1,1-dichloroethylene toxicity. *Toxicol Appl Pharmacol* 57: 146-155. [http://dx.doi.org/10.1016/0041-008X\(81\)90274-X](http://dx.doi.org/10.1016/0041-008X(81)90274-X).
- Chiesa, R; Donato, F; Tagger, A; Favret, M; Ribero, M; Nardi, G; Gelatti, U; Bucella, E; Tomasi, E; Portolani, N; Bonetti, M; Bettini, L; Pelizzari, G; Salmi, A; Savio, A; Garatti, M; Callea, F. (2000). Etiology of hepatocellular carcinoma in Italian patients with and without cirrhosis. *Cancer Epidemiol Biomarkers Prev* 9: 213-216.
- Chiu, B and Weisenburger, D. (2003). An update of the epidemiology of non-Hodgkin's lymphoma. 4: 161-168.
- Chiu, W and White, P. (2004). PBPK model-based route extrapolation: Exact and approximate solutions for use in risk assessment.
- Chiu, W. (2006). Statistical issues in physiologically based pharmacokinetic modeling. In JC Lipscomb & EV Ohanian (Eds.), *Toxicokinetics And Risk Assessment* (pp. 269). New York: Informa Healthcare.
- Chiu, W; Caldwell, J; Keshava, N; Scott, C. (2006a). Key scientific issues in the health risk assessment of trichloroethylene. *Environ Health Perspect* 114: 1445-1449. <http://dx.doi.org/10.1289/ehp.8690>.
- Chiu, W; Okino, M; Lipscomb, J; Evans, M. (2006b). Issues in the pharmacokinetics of trichloroethylene and its metabolites. *Environ Health Perspect* 114: 1450-1456. <http://dx.doi.org/10.1289/ehp.8691>.

- Chiu, W and White, P. (2006). Steady-state solutions to PBPK models and their applications to risk assessment I: Route-to-route extrapolation of volatile chemicals. *Risk Anal* 26: 769-780.
<http://dx.doi.org/10.1111/j.1539-6924.2006.00762.x>.
- Chiu, W and Bois, F. (2007). An approximate method for population toxicokinetic analysis with aggregated data. *Journal of Agricultural, Biological, and Environmental Statistics* 12: 346-363.
<http://dx.doi.org/10.1198/108571107X229340>.
- Chiu, W; Micallef, S; Monster, A; Bois, F. (2007). Toxicokinetics of inhaled trichloroethylene and tetrachloroethylene in humans at 1 ppm: Empirical results and comparisons with previous studies. *Toxicol Sci* 95: 23-36. <http://dx.doi.org/10.1093/toxsci/kfl129>.
- Chiu, W; Okino, M; Evans, M. (2009). Characterizing uncertainty and population variability in the toxicokinetics of trichloroethylene and metabolites in mice, rats, and humans using an updated database, physiologically based pharmacokinetic (PBPK) model, and Bayesian approach. *Toxicol Appl Pharmacol* 241: 36-60. <http://dx.doi.org/10.1016/j.taap.2009.07.032>.
- Chiu, W. (2011). Trichloroacetic acid: Updated estimates of its bioavailability and its contribution to trichloroethylene-induced mouse hepatomegaly. *Toxicology* 285: 114-125.
<http://dx.doi.org/10.1016/j.tox.2011.04.009>.
- Cho, W. (2007). OncomiRs: The discovery and progress of microRNAs in cancers. *Mol Cancer* 6: 60.
<http://dx.doi.org/10.1186/1476-4598-6-60>.
- Chow, W; Gridley, G; McLaughlin, J; Mandel, J; Wacholder, S; Blot, W; Niwa, S; Fraumeni, J. (1994). Protein intake and risk of renal cell cancer. *J Natl Cancer Inst* 86: 1131-1139.
- Chow, W and Devesa, S. (2008). Contemporary epidemiology of renal cell cancer. *Cancer J* 14: 288-301.
<http://dx.doi.org/10.1097/PPO.0b013e3181867628>.
- Cicmanec, J; Condie, L; Olson, G; Wang, S. (1991). 90-Day toxicity study of dichloroacetate in dogs. *Fundam Appl Toxicol* 17: 376-389.
- Clapp, R and Hoffman, K. (2008). Cancer mortality in IBM Endicott plant workers, 1969-2001: An update on a NY production plant. *Environ Health* 7: 13. <http://dx.doi.org/10.1186/1476-069X-7-13>.
- Clay, P. (2008). Assessment of the genotoxicity of trichloroethylene and its metabolite, S-(1,2-dichlorovinyl)-L-cysteine (DCVC), in the comet assay in rat kidney. *Mutagenesis* 23: 27-33.
<http://dx.doi.org/10.1093/mutage/gem034>.
- Clegg, E; Cook, J; Chapin, R; Foster, P; Daston, G. (1997). Leydig cell hyperplasia and adenoma formation: Mechanisms and relevance to humans. *Reprod Toxicol* 11: 107-121.
[http://dx.doi.org/10.1016/S0890-6238\(96\)00203-1](http://dx.doi.org/10.1016/S0890-6238(96)00203-1).
- Clewell, H and Andersen, M. (2004). Applying mode-of-action and pharmacokinetic considerations in contemporary cancer risk assessments: An example with trichloroethylene. *Crit Rev Toxicol* 34: 385-445. <http://dx.doi.org/10.1080/10408440490500795>.
- Clewell, H, III; Gentry, P; Gearhart, J; Allen, B; Covington, T; Andersen, M. (1995). Applying mechanistic information, pharmacokinetic modeling, and margin of exposure evaluations in a novel risk assessment for trichloroethylene and its active carcinogenic metabolites: chloral hydrate, trichloroacetic acid, dichloroacetic acid, and dichlorovinylcysteine.
- Clewell, H, III; Gentry, P; Covington, T; Gearhart, J. (2000). Development of a physiologically based pharmacokinetic model of trichloroethylene and its metabolites for use in risk assessment. *Environ Health Perspect* 108: 283-305.
- Coberly, S; Oudiz, D; Overstreet, J; Wiley, L. (1992). Effects of maternal exposure to trichloroethylene (TCE) on cell proliferation in the mouse preimplantation embryo. *Reprod Toxicol* 6: 241-245.
- Cocco, P (2010) [Sezione di Medicina del Lavoro, Università di Cagliari to Cheryl Siegel Scott, U. S. EPA, December 12, 2010].
- Cocco, P. (2011). personal communication from Pierluigi Cocco, University of Cagliari, Italy, to Cheryl Scott, U.S. EPA, 19 March 2011
- Cocco, P; t'Mannetje, A; Fadda, D; Melis, M; Becker, N; de Sanjosé, S; Foretova, L; Mareckova, J; Staines, A; Kleefeld, S; Maynadié, M; Nieters, A; Brennan, P; Boffetta, P. (2010). Occupational

- exposure to solvents and risk of lymphoma subtypes: results from the Epilymph case-control study. Occup Environ Med 67: 341-347. <http://dx.doi.org/10.1136/oem.2009.046839>.
- Cogger, V; Warren, A; Fraser, R; Ngu, M; McLean, A; Le Couteur, D. (2003). Hepatic sinusoidal pseudocapillarization with aging in the non-human primate. Exp Gerontol 38: 1101-1107.
- Cohen, P; Nishimura, K; Zhu, L; Pollard, J. (1999). Macrophages: Important accessory cells for reproductive function. J Leukoc Biol 66: 765-772.
- Cohn, P; Fagliano, J; Klotz, J. (1994a). Assessing human health effects from chemical contaminants in drinking water. J Med Soc New J 91: 719-722.
- Cohn, P; Klotz, J; Bove, F; Berkowitz, M; Fagliano, J. (1994b). Drinking water contamination and the incidence of leukemia and non-Hodgkin's lymphoma. Environ Health Perspect 102: 556-561.
- Cojocel, C; Beuter, W; Müller, W; Mayer, D. (1989). Lipid peroxidation: A possible mechanism of trichloroethylene-induced nephrotoxicity. Toxicology 55: 131-141.
[http://dx.doi.org/10.1016/0300-483X\(89\)90180-7](http://dx.doi.org/10.1016/0300-483X(89)90180-7).
- Coleridge, H; Coleridge, J; Luck, J; Norman, J. (1968). The effect of four volatile anaesthetic agents on the impulse activity of two types of pulmonary receptor. Br J Anaesth 40: 484-492.
<http://dx.doi.org/10.1093/bja/40.7.484>.
- Collier, J; Selmin, O; Johnson, P; Runyan, R. (2003). Trichloroethylene effects on gene expression during cardiac development. Birth Defects Res A Clin Mol Teratol 67: 488-495.
<http://dx.doi.org/10.1002/bdra.10073>.
- Colt, J and Blair, A. (1998). Parental occupational exposures and risk of childhood cancer. Environ Health Perspect 106 Suppl 3: 909-925.
- Columbano, A; Ledda-Columbano, G; Coni, P; Faa, G; Liguori, C; Santa Cruz, G; Pani, P. (1985). Occurrence of cell death (apoptosis) during the involution of liver hyperplasia. Lab Invest 52: 670-675.
- Columbano, A; Ledda-Columbano, G; Lee, G; Rajalakshmi, S; Sarma, D. (1987). Inability of mitogen-induced liver hyperplasia to support the induction of enzyme-altered islands induced by liver carcinogens. Cancer Res 47: 5557-5559.
- Columbano, A; Ledda-Columbano, G; Ennas, M; Curto, M; Chelo, A; Pani, P. (1990). Cell proliferation and promotion of rat liver carcinogenesis: Different effect of hepatic regeneration and mitogen induced hyperplasia on the development of enzyme-altered foci. Carcinogenesis 11: 771-776.
<http://dx.doi.org/10.1093/carcin/11.5.771>.
- Columbano, A and Ledda-Columbano, G. (2003). Mitogenesis by ligands of nuclear receptors: an attractive model for the study of the molecular mechanisms implicated in liver growth. Cell Death Differ 10: S19-S21. <http://dx.doi.org/10.1038/sj.cdd.4401113>.
- Colvin, M; Myers, J; Nell, V; Rees, D; Cronje, R. (1993). A cross-sectional survey of neurobehavioral effects of chronic solvent exposure on workers in a paint manufacturing plant. Environ Res 63: 122-132. <http://dx.doi.org/10.1006/enrs.1993.1134>.
- Commandeur, J and Vermeulen, N. (1990). Identification of N-acetyl(2,2-dichlorovinyl)- and N-acetyl(1,2-dichlorovinyl)-L-cysteine as two regioisomeric mercapturic acids of trichloroethylene in the rat. Chem Res Toxicol 3: 212-218. <http://dx.doi.org/10.1021/tx00015a005>.
- Coni, P; Simbula, G; De Prati, A; Menegazzi, M; Suzuki, H; Sarma, D; Ledda-Columbano, G; Columbano, A. (1993). Differences in the steady-state levels of c-fos, c-jun and c-myc messenger RNA during mitogen-induced liver growth and compensatory regeneration. Hepatology 17: 1109-1116.
- Conner, E; Lemmer, E; Omori, M; Wirth, P; Factor, V; Thorgeirsson, S. (2000). Dual functions of E2F-1 in a transgenic mouse model of liver carcinogenesis. Oncogene 19: 5054-5062.
<http://dx.doi.org/10.1038/sj.onc.1203885>.
- Conner, E; Lemmer, E; Sánchez, A; Factor, V; Thorgeirsson, S. (2003). E2F1 blocks and c-Myc accelerates hepatic ploidy in transgenic mouse models. Biochem Biophys Res Commun 302: 114-120.

- Contrera, J; Jacobs, A; DeGeorge, J. (1997). Carcinogenicity testing and the evaluation of regulatory requirements for pharmaceuticals. *Regul Toxicol Pharmacol* 25: 130-145.
<http://dx.doi.org/10.1006/rtpb.1997.1085>.
- Cook, J; Klinefelter, G; Hardisty, J; Sharpe, R; Foster, P. (1999). Rodent leydig cell tumorigenesis: A review of the physiology, pathology, mechanisms and relevance to humans. *Crit Rev Toxicol* 29: 169-261.
- Cooper, A. (2004). The role of glutamine transaminase K (GTK) in sulfur and alpha-keto acid metabolism in the brain, and in the possible bioactivation of neurotoxicants. *Neurochem Int* 44: 557-577.
<http://dx.doi.org/10.1016/j.neuint.2003.12.002>.
- Cooper, G; Miller, F; Pandey, J. (1999). The role of genetic factors in autoimmune disease: implications for environmental research. *Environ Health Perspect* 107 Suppl 5: 693-700.
- Cooper, G and Stroehla, B. (2003). The epidemiology of autoimmune diseases. *Autoimmun Rev* 2: 119-125.
- Cooper, G; Parks, C; Treadwell, E; St Clair, E; Gilkeson, G; Dooley, M. (2004). Occupational risk factors for the development of systemic lupus erythematosus. *J Rheumatol* 31: 1928-1933.
- Cooper, G; Makris, S; Nietert, P; Jinot, J. (2009). Evidence of autoimmune-related effects of trichloroethylene exposure from studies in mice and humans. *Environ Health Perspect* 117: 696-702. <http://dx.doi.org/10.1289/ehp.11782>.
- Coopman, V; Cordonnier, J; De Letter, E; Piette, M. (2003). Tissue distribution of trichloroethylene in a case of accidental acute intoxication by inhalation. *Forensic Sci Int* 134: 115-119.
[http://dx.doi.org/10.1016/S0379-0738\(03\)00131-2](http://dx.doi.org/10.1016/S0379-0738(03)00131-2).
- Copeland, K; Checkoway, H; McMichael, A; Holbrook, R. (1977). Bias due to misclassification in the estimation of relative risk. *Am J Epidemiol* 105: 488-495.
- Cornett, R; James, M; Henderson, G; Cheung, J; Shroads, A; Stacpoole, P. (1999). Inhibition of glutathione S-transferase zeta and tyrosine metabolism by dichloroacetate: A potential unifying mechanism for its altered biotransformation and toxicity. *Biochem Biophys Res Commun* 262: 752-756. <http://dx.doi.org/10.1006/bbrc.1999.1287>.
- Corton, J and Lapinskas, P. (2005). Peroxisome proliferator-activated receptors: Mediators of phthalate ester-induced effects in the male reproductive tract? *Toxicol Sci* 83: 4-17.
<http://dx.doi.org/10.1093/toxsci/kfi011>.
- Corton, J. (2008). Evaluation of the role of peroxisome proliferator-activated receptor alpha (PPARalpha) in mouse liver tumor induction by trichloroethylene and metabolites. *Crit Rev Toxicol* 38: 857-875. <http://dx.doi.org/10.1080/10408440802209796>.
- Cosby, N and Dukelow, W. (1992). Toxicology of maternally ingested trichloroethylene (TCE) on embryonal and fetal development in mice and of TCE metabolites on in vitro fertilization. *Toxicol Sci* 19: 268-274.
- Costa, A and Ivanetich, K. (1984). Chlorinated ethylenes: their metabolism and effect on DNA repair in rat hepatocytes. *Carcinogenesis* 5: 1629-1636.
- Costa, G; Merletti, F; Segnan, N. (1989). A mortality cohort study in a north Italian aircraft factory. *Br J Ind Med* 46: 738-743.
- Costantini, A; Benvenuti, A; Vineis, P; Kriebel, D; Tumino, R; Ramazzotti, V; Rodella, S; Stagnaro, E; Crosignani, P; Amadori, D; Mirabelli, D; Sommani, L; Belletti, I; Troschel, L; Romeo, L; Miceli, G; Tozzi, G; Mendico, I; Maltoni, S; Miligi, L. (2008). Risk of leukemia and multiple myeloma associated with exposure to benzene and other organic solvents: Evidence from the Italian Multicenter Case-control study. *Am J Ind Med* 51: 803-811.
<http://dx.doi.org/10.1002/ajim.20592>.
- Costas, K; Knorr, R; Condon, S. (2002). A case-control study of childhood leukemia in Woburn, Massachusetts: The relationship between leukemia incidence and exposure to public drinking water. *Sci Total Environ* 300: 23-35. [http://dx.doi.org/10.1016/S0048-9697\(02\)00169-9](http://dx.doi.org/10.1016/S0048-9697(02)00169-9).

- Costet, P; Legendre, C; Moré, J; Edgar, A; Galtier, P; Pineau, T. (1998). Peroxisome proliferator-activated receptor alpha-isoform deficiency leads to progressive dyslipidemia with sexually dimorphic obesity and steatosis. *J Biol Chem* 273: 29577-29585.
- Counts, J; Sarmiento, J; Harbison, M; Downing, J; McClain, R; Goodman, J. (1996). Cell proliferation and global methylation status changes in mouse liver after phenobarbital and/or choline-devoid, methionine-deficient diet administration. *Carcinogenesis* 17: 1251-1257.
- Coyle, Y; Hynan, L; Euhus, D; Minhajuddin, A. (2005). An ecological study of the association of environmental chemicals on breast cancer incidence in Texas. *Breast Cancer Res Treat* 92: 107-114. <http://dx.doi.org/10.1007/s10549-004-8268-z>.
- Crabb, D; Pinaire, J; Chou, W; Sissom, S; Peters, J; Harris, R; Stewart, M. (2001). Peroxisome proliferator-activated receptors (PPAR) and the mitochondrial aldehyde dehydrogenase (ALDH2) promoter in vitro and in vivo. *Alcohol Clin Exp Res* 25: 945-952.
- Crebelli, R; Bignami, M; Conti, L; Carere, A. (1982). Mutagenicity of trichloroethylene in *Salmonella typhimurium* TA100. *Ann Ist Super Sanita* 18: 117-121.
- Crebelli, R; Conti, G; Conti, L; Carere, A. (1985). Mutagenicity of trichloroethylene, trichloroethanol and chloral hydrate in *Aspergillus nidulans*. *Mutat Res Genet Toxicol Environ Mutagen* 155: 105-111. [http://dx.doi.org/10.1016/0165-1218\(85\)90126-0](http://dx.doi.org/10.1016/0165-1218(85)90126-0).
- Crebelli, R and Carere, A. (1989). Genetic toxicology of 1,1,2-trichloroethylene. *Mutat Res* 221: 11-37. [http://dx.doi.org/10.1016/0165-1110\(89\)90043-2](http://dx.doi.org/10.1016/0165-1110(89)90043-2).
- Crebelli, R; Conti, G; Conti, L; Carere, A. (1991). In vitro studies with nine known or suspected spindle poisons: results in tests for chromosome malsegregation in *Aspergillus nidulans*. *Mutagenesis* 6: 131-136. <http://dx.doi.org/10.1093/mutage/6.2.131>.
- Crofton, K and Zhao, X. (1993). Mid-frequency hearing loss in rats following inhalation exposure to trichloroethylene: evidence from reflex modification audiometry. *Neurotoxicol Teratol* 15: 413-423.
- Crofton, K; Lassiter, T; Rebert, C. (1994). Solvent-induced ototoxicity in rats: An atypical selective mid-frequency hearing deficit. *Hear Res* 80: 25-30. [http://dx.doi.org/10.1016/0378-5955\(94\)90005-1](http://dx.doi.org/10.1016/0378-5955(94)90005-1).
- Crofton, K and Zhao, X. (1997). The ototoxicity of trichloroethylene: extrapolation and relevance of high-concentration, short-duration animal exposure data. *Fundam Appl Toxicol* 38: 101-106.
- Crump, K; Allen, B; Shipp, A. (1989). Choice of dose measure for extrapolating carcinogenic risk from animals to humans: An empirical investigation of 23 chemicals. *Health Phys* 57: 387-393.
- Cruzan, G; Bus, J; Banton, M; Gingell, R; Carlson, G. (2009). Mouse specific lung tumors from CYP2F2-mediated cytotoxic metabolism: an endpoint/toxic response where data from multiple chemicals converge to support a mode of action. *Regul Toxicol Pharmacol* 55: 205-218. <http://dx.doi.org/10.1016/j.yrph.2009.07.002>.
- Cummings, B; Zangar, R; Novak, R; Lash, L. (1999). Cellular distribution of cytochromes P-450 in the rat kidney. *Drug Metab Dispos* 27: 542-548.
- Cummings, B and Lash, L. (2000). Metabolism and toxicity of trichloroethylene and S-(1,2-dichlorovinyl)-L-cysteine in freshly isolated human proximal tubular cells. *Toxicol Sci* 53: 458-466.
- Cummings, B; Lasker, J; Lash, L. (2000a). Expression of glutathione-dependent enzymes and cytochrome P450s in freshly isolated and primary cultures of proximal tubular cells from human kidney. *J Pharmacol Exp Ther* 293: 677-685.
- Cummings, B; Parker, J; Lash, L. (2000b). Role of cytochrome P450 and glutathione S-transferase alpha in the metabolism and cytotoxicity of trichloroethylene in rat kidney. *Biochem Pharmacol* 59: 531-543. [http://dx.doi.org/10.1016/S0006-2952\(99\)00374-3](http://dx.doi.org/10.1016/S0006-2952(99)00374-3).
- Cummings, B; Zangar, R; Novak, R; Lash, L. (2000c). Cytotoxicity of trichloroethylene and S-(1,2-dichlorovinyl)-L-cysteine in primary cultures of rat renal proximal tubular and distal tubular cells. *Toxicology* 150: 83-98. [http://dx.doi.org/10.1016/S0300-483X\(00\)00252-3](http://dx.doi.org/10.1016/S0300-483X(00)00252-3).
- Cummings, B; Parker, J; Lash, L. (2001). Cytochrome p450-dependent metabolism of trichloroethylene in rat kidney. *Toxicol Sci* 60: 11-19.

- Cutler, J; Parker, G; Rosen, S; Prenney, B; Healey, R; Caldwell, G. (1986). Childhood leukemia in Woburn, Massachusetts. Public Health Rep 101: 201-205.
- D'Souza, R; Bruckner, J; Feldman, S. (1985). Oral and intravenous trichloroethylene pharmacokinetics in the rat. J Toxicol Environ Health 15: 587-601.
- Dahlgren, J; Takhar, H; Anderson-Mahoney, P; Kotlerman, J; Tarr, J; Warshaw, R. (2007). Cluster of systemic lupus erythematosus (SLE) associated with an oil field waste site: a cross sectional study. Environ Health 6: 8. <http://dx.doi.org/10.1186/1476-069X-6-8>.
- Dai, X; De Souza, A; Dai, H; Lewis, D; Lee, C; Spencer, A; Herweijer, H; Hagstrom, J; Linsley, P; Bassett, D; Ulrich, R; He, Y. (2007). PPARalpha siRNA-treated expression profiles uncover the causal sufficiency network for compound-induced liver hypertrophy. PLoS Comput Biol 3: e30. <http://dx.doi.org/10.1371/journal.pcbi.0030030>.
- Dai, Y; Leng, S; Li, L; Niu, Y; Huang, H; Cheng, J; Zheng, Y. (2004). Genetic polymorphisms of cytokine genes and risk for trichloroethylene-induced severe generalized dermatitis: a case-control study. Biomarkers 9: 470-478. <http://dx.doi.org/10.1080/13547500400026920>.
- Dajee, M; Lazarov, M; Zhang, J; Cai, T; Green, C; Russell, A; Marinkovich, M; Tao, S; Lin, Q; Kubo, Y; Khavari, P. (2003). NF-kappaB blockade and oncogenic Ras trigger invasive human epidermal neoplasia. Nature 421: 639-643. <http://dx.doi.org/10.1038/nature01283>.
- Dalbey, W and Bingham, E. (1978). Metabolism of trichloroethylene by the isolated perfused lung. Toxicol Appl Pharmacol 43: 267-277.
- Dallas, C. (1991). Validation and application of pharmacokinetic models for interspecies extrapolations in toxicity risk assessments of volatile organics. Washington, DC: Air Force Office of Scientific Research. <http://www.biomedsearch.com/sci/Validation-Application-Pharmacokinetic-Models-Interspecies/0033755846.html>.
- Dallas, C; Gallo, J; Ramanathan, R; Muralidhara, S; Bruckner, J. (1991). Physiological pharmacokinetic modeling of inhaled trichloroethylene in rats. Toxicol Appl Pharmacol 110: 303-314.
- Danhof, M; de Jongh, J; De Lange, E; Della Pasqua, O; Ploeger, B; Voskuyl, R. (2007). Mechanism-based pharmacokinetic-pharmacodynamic modeling: biophase distribution, receptor theory, and dynamical systems analysis. Annu Rev Pharmacol Toxicol 47: 357-400. <http://dx.doi.org/10.1146/annurev.pharmtox.47.120505.105154>.
- Daniel, F; DeAngelo, A; Stober, J; Olson, G; Page, N. (1992). Hepatocarcinogenicity of chloral hydrate, 2-chloroacetaldehyde, and dichloroacetic acid in the male B6C3F1 mouse. Fundam Appl Toxicol 19: 159-168. [http://dx.doi.org/10.1016/0272-0590\(92\)90147-A](http://dx.doi.org/10.1016/0272-0590(92)90147-A).
- Daniell, W; Claypoole, K; Checkoway, H; Smith-Weller, T; Dager, S; Townes, B; Rosenstock, L. (1999). Neuropsychological function in retired workers with previous long-term occupational exposure to solvents. Occup Environ Med 56: 93-105.
- Darnerud, P; Brandt, I; Feil, V; Bakke, J. (1989). Dichlorovinyl cysteine (DCVC) in the mouse kidney: tissue-binding and toxicity after glutathione depletion and probenecid treatment. Arch Toxicol 63: 345-350.
- Das, R and Scott, J. (1994). Trichloroethylene-induced pneumotoxicity in fetal and neonatal mice. Toxicol Lett 73: 227-239.
- David, R; Moore, M; Cifone, M; Finney, D; Guest, D. (1999). Chronic peroxisome proliferation and hepatomegaly associated with the hepatocellular tumorigenesis of di(2-ethylhexyl)phthalate and the effects of recovery. Toxicol Sci 50: 195-205.
- David, R; Clewell, H; Gentry, P; Covington, T; Morgott, D; Marino, D. (2006). Revised assessment of cancer risk to dichloromethane II. Application of probabilistic methods to cancer risk determinations. Regul Toxicol Pharmacol 45: 55-65. <http://dx.doi.org/10.1016/j.yrtph.2005.12.003>.
- Davidson, I and Beliles, R. (1991). Consideration of the target organ toxicity of trichloroethylene in terms of metabolite toxicity and pharmacokinetics. Drug Metab Rev 23: 493-599. <http://dx.doi.org/10.3109/03602539109029772>.

- Davis, R; Murphy, W; Snawder, J; Striley, C; Henderson, D; Khan, A; Krieg, E. (2002). Susceptibility to the ototoxic properties of toluene is species specific. *Hear Res* 166: 24-32.
[http://dx.doi.org/10.1016/S0378-5955\(02\)00280-0](http://dx.doi.org/10.1016/S0378-5955(02)00280-0).
- Davis, S; Laszlo Pallos, L; Wu, J; Sapp, J; Cusack, C. (2005). ATSDR's trichloroethylene subregistry methods and results: 1989-2000. *Arch Environ Occup Health* 60: 130-139.
<http://dx.doi.org/10.3200/AEOH.60.3.130-139>.
- Dawson, B; Johnson, P; Goldberg, S; Ulreich, J. (1990). Cardiac teratogenesis of trichloroethylene and dichloroethylene in a mammalian model. *J Am Coll Cardiol* 16: 1304-1309.
- Dawson, B; Johnson, P; Goldberg, S; Ulreich, J. (1993). Cardiac teratogenesis of halogenated hydrocarbon-contaminated drinking water. *J Am Coll Cardiol* 21: 1466-1472.
[http://dx.doi.org/10.1016/0735-1097\(93\)90325-U](http://dx.doi.org/10.1016/0735-1097(93)90325-U).
- De Baere, S; Meyer, E; Dirinck, I; Lambert, W; Piette, M; Van Peteghem, C; De Leenheer, A. (1997). Tissue distribution of trichloroethylene and its metabolites in a forensic case. *J Anal Toxicol* 21: 223-227.
- de Jong, P and Brenner, B. (2004). From secondary to primary prevention of progressive renal disease: the case for screening for albuminuria. *Kidney Int* 66: 2109-2118.
<http://dx.doi.org/10.1111/j.1523-1755.2004.66001.x>.
- De Roos, A; Olshan, A; Teschke, K; Poole, C; Savitz, D; Blatt, J; Bondy, M; Pollock, B. (2001). Parental occupational exposures to chemicals and incidence of neuroblastoma in offspring. *Am J Epidemiol* 154: 106-114. <http://dx.doi.org/10.1093/aje/154.2.106>.
- Deane, M; Swan, S; Harris, J; Epstein, D; Neutra, R. (1989). Adverse pregnancy outcomes in relation to water contamination, Santa Clara County, California, 1980-1981. *Am J Epidemiol* 129: 894-904.
- DeAngelo, A; Daniel, F; McMillan, L; Wernsing, P; Savage, R, Jr. (1989). Species and strain sensitivity to the induction of peroxisome proliferation by chloroacetic acids. *Toxicol Appl Pharmacol* 101: 285-298. [http://dx.doi.org/10.1016/0041-008X\(89\)90277-9](http://dx.doi.org/10.1016/0041-008X(89)90277-9).
- DeAngelo, A; Daniel, F; Most, B; Olson, G. (1996). The carcinogenicity of dichloroacetic acid in the male fischer 344 rat. *Toxicology* 114: 207-221. [http://dx.doi.org/10.1016/S0300-483X\(96\)03510-X](http://dx.doi.org/10.1016/S0300-483X(96)03510-X).
- DeAngelo, A; Daniel, F; Most, B; Olson, G. (1997). Failure of monochloroacetic acid and trichloroacetic acid administered in the drinking water to produce liver cancer in male F344/N rats. *J Toxicol Environ Health* 52: 425-445.
- DeAngelo, A; George, M; House, D. (1999). Hepatocarcinogenicity in the male B6C3F1 mouse following a lifetime exposure to dichloroacetic acid in the drinking water: Dose-response determination and modes of action. *J Toxicol Environ Health A* 58: 485-507.
- DeAngelo, A; Daniel, F; Wong, D; George, M. (2008). The induction of hepatocellular neoplasia by trichloroacetic acid administered in the drinking water of the male B6C3F1 mouse. *J Toxicol Environ Health A* 71: 1056-1068. <http://dx.doi.org/10.1080/15287390802111952>.
- Deerberg, F and Müller-Peddinghaus, R. (1970). [Comparative studies on the occurrence of spontaneous diseases and spontaneous deaths in conventional and specific germ-free female NMRI-HAN mice]. *Z Versuchstierkd* 12: 341-347.
- Deerberg, F; Pittermann, W; Rapp, K. (1974). [Lung tumor in the mouse: a progressive tumor]. *Vet Pathol* 11: 430-441.
- Dees, C and Travis, C. (1993). The mitogenic potential of trichloroethylene in B6C3F1 mice. *Toxicol Lett* 69: 129-137.
- Dees, C and Travis, C. (1994). Trichloroacetate stimulation of liver DNA synthesis in male and female mice. *Toxicol Lett* 70: 343-355. [http://dx.doi.org/10.1016/0378-4274\(94\)90129-5](http://dx.doi.org/10.1016/0378-4274(94)90129-5).
- Degrassi, F and Tanzarella, C. (1988). Immunofluorescent staining of kinetochores in micronuclei: A new assay for the detection of aneuploidy. *Mutat Res Genet Toxicol Environ Mutagen* 203: 339-345.
[http://dx.doi.org/10.1016/0165-1161\(88\)90030-1](http://dx.doi.org/10.1016/0165-1161(88)90030-1).

- Dehon, B; Humbert, L; Devisme, L; Stievenart, M; Mathieu, D; Houdret, N; Lhermitte, M. (2000). Tetrachloroethylene and trichloroethylene fatality: Case report and simple headspace SPME-capillary gas chromatographic determination in tissues. *J Anal Toxicol* 24: 22-26.
- Dekant, W; Metzler, M; Henschler, D. (1984). Novel metabolites of trichloroethylene through dechlorination reactions in rats, mice and humans. *Biochem Pharmacol* 33: 2021-2027.
- Dekant, W; Metzler, M; Henschler, D. (1986a). Identification of S-1,2-dichlorovinyl-N-acetyl-cysteine as a urinary metabolite of trichloroethylene: A possible explanation for its nephrocarcinogenicity in male rats. *Biochem Pharmacol* 35: 2455-2458. [http://dx.doi.org/10.1016/0006-2952\(86\)90039-0](http://dx.doi.org/10.1016/0006-2952(86)90039-0).
- Dekant, W; Schulz, A; Metzler, M; Henschler, D. (1986b). Absorption, elimination, and metabolism of trichlorethylene: A quantitative comparison between rats and mice. *Xenobiotica* 16: 143-152.
- Dekant, W; Vamvakas, S; Berthold, K; Schmidt, S; Wild, D; Henschler, D. (1986c). Bacterial beta-lyase mediated cleavage and mutagenicity of cysteine conjugates derived from the nephrocarcinogenic alkenes trichloroethylene, tetrachloroethylene and hexachlorobutadiene. *Chem Biol Interact* 60: 31-45. [http://dx.doi.org/10.1016/0009-2797\(86\)90015-3](http://dx.doi.org/10.1016/0009-2797(86)90015-3).
- Dekant, W; Berthold, K; Vamvakas, S; Henschler, D; Anders, M. (1988). Thioacylating intermediates as metabolites of S-(1,2-dichlorovinyl)-L-cysteine and S-(1,2,2-trichlorovinyl)-L-cysteine formed by cysteine conjugate beta-lyase. *Chem Res Toxicol* 1: 175-178. <http://dx.doi.org/10.1021/tx00003a008>.
- Dekant, W. (1990). Bioactivation of halogenated alkenes by glutathione conjugation a mechanistic explanation for their renal carcinogenicity. *J Cancer Res Clin Oncol* 116: 1063.
- Delahunt, B; Bethwaite, P; Nacey, J. (1995). Occupational risk for renal cell carcinoma. A case-control study based on the New Zealand Cancer Registry. *Br J Urol* 75: 578-582. <http://dx.doi.org/10.1111/j.1464-410X.1995.tb07410.x>.
- Delinsky, A; Delinsky, D; Muralidhara, S; Fisher, J; Bruckner, J; Bartlett, M. (2005). Analysis of dichloroacetic acid in rat blood and tissues by hydrophilic interaction liquid chromatography with tandem mass spectrometry. *Rapid Commun Mass Spectrom* 19: 1075-1083. <http://dx.doi.org/10.1002/rcm.189>.
- DeMarini, D; Perry, E; Shelton, M. (1994). Dichloroacetic acid and related compounds: Induction of prophage in *E. coli* and mutagenicity and mutation spectra in *Salmonella TA100*. *Mutagenesis* 9: 429-437. <http://dx.doi.org/10.1093/mutage/9.5.429>.
- DerSimonian, R and Laird, N. (1986). Meta-analysis in clinical trials. *Contr Clin Trials* 7: 177-188.
- Dewar, R; Siemiatyckia, J; Geacuterin, M. (1994). Loss of statistical power associated with the use of a job-exposure matrix in occupational case-control studies. *Appl Occup Environ Hyg* 6: 508-515.
- Ding, R; Shen, T; Zhu, Q. (2006). [Effects of TCE and PCE on cultured human keratinocyte lipid peroxidation and protective effect of vitamin E on it]. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi* 24: 662-665.
- Diot, E; Lesire, V; Guilmot, J; Metzger, M; Pilore, R; Rogier, S; Stadler, M; Diot, P; Lemarie, E; Lasfargues, G. (2002). Systemic sclerosis and occupational risk factors: a case-control study. *Occup Environ Med* 59: 545-549.
- DiRenzo, A; Gandolfi, A; Sipes, I. (1982). Microsomal bioactivation and covalent binding of aliphatic halides to DNA. *Toxicol Lett* 11: 243-252.
- Dix, K; Kedderis, G; Borghoff, S. (1997). Vehicle-dependent oral absorption and target tissue dosimetry of chloroform in male rats and female mice. *Toxicol Lett* 91: 197-209.
- Djouadi, F; Weinheimer, C; Saffitz, J; Pitchford, C; Bastin, J; Gonzalez, F; Kelly, D. (1998). A gender-related defect in lipid metabolism and glucose homeostasis in peroxisome proliferator- activated receptor alpha- deficient mice. *J Clin Invest* 102: 1083-1091. <http://dx.doi.org/10.1172/JCI3949>.
- Dobrev, I; Andersen, M; Yang, R. (2001). Assessing interaction thresholds for trichloroethylene in combination with tetrachloroethylene and 1,1,1-trichloroethane using gas uptake studies and PBPK modeling. *Arch Toxicol* 75: 134-144. <http://dx.doi.org/10.1007/s002040100216>.

- Dobrev, I; Andersen, M; Yang, R. (2002). In silico toxicology: Simulating interaction thresholds for human exposure to mixtures of trichloroethylene, tetrachloroethylene, and 1,1,1-trichloroethane. Environ Health Perspect 110: 1031-1039.
- Dohn, D and Anders, M. (1982). Assay of cysteine conjugate beta-lyase activity with S-(2-benzothiazolyl)cysteine as the substrate. Anal Biochem 120: 379-386.
[http://dx.doi.org/10.1016/0003-2697\(82\)90361-X](http://dx.doi.org/10.1016/0003-2697(82)90361-X).
- Dolinoy, D; Weidman, J; Jirtle, R. (2007). Epigenetic gene regulation: Linking early developmental environment to adult disease. Reprod Toxicol 23: 297-307.
<http://dx.doi.org/10.1016/j.reprotox.2006.08.012>.
- Donoghue, A; Dryson, E; Wynn-Williams, G. (1995). Contrast sensitivity in organic-solvent-induced chronic toxic encephalopathy. J Occup Environ Med 37: 1357-1363.
- Dorfmueller, M; Henne, S; York, R; Bornschein, R; Manson, J. (1979). Evaluation of teratogenicity and behavioral toxicity with inhalation exposure of maternal rats to trichloroethylene. Toxicology 14: 153-166.
- Dorne, J; Walton, K; Renwick, A. (2005). Human variability in xenobiotic metabolism and pathway-related uncertainty factors for chemical risk assessment: a review. Food Chem Toxicol 43: 203-216. <http://dx.doi.org/10.1016/j.fct.2004.05.011>.
- Dosemeci, M; Wacholder, S; Lubin, J. (1990). Does nondifferential misclassification of exposure always bias a true effect toward the null value? Am J Epidemiol 132: 746-748.
- Dosemeci, M; Cocco, P; Gómez, M; Stewart, P; Heineman, E. (1994). Effects of three features of a job-exposure matrix on risk estimates. Epidemiology 5: 124-127.
- Dosemeci, M; Cocco, P; Chow, W-H. (1999). Gender differences in risk of renal cell carcinoma and occupational exposures to chlorinated aliphatic hydrocarbons. Am J Ind Med 36: 54-59.
[http://dx.doi.org/10.1002/\(SICI\)1097-0274\(199907\)36:1<54::AID-AJIM8>3.0.CO;2-0](http://dx.doi.org/10.1002/(SICI)1097-0274(199907)36:1<54::AID-AJIM8>3.0.CO;2-0).
- Døssing, M; Petersen, K; Vyberg, M; Olsen, J. (1997). Liver cancer among employees in Denmark. Am J Ind Med 32: 248-254.
- Douglas, G; Gingerich, J; Soper, L; Potvin, M; Bjarnason, S. (1999). Evidence for the lack of base-change and small-deletion mutation induction by trichloroethylene in lacZ transgenic mice. Environ Mol Mutagen 34: 190-194.
- Dow, J and Green, T. (2000). Trichloroethylene induced vitamin B(12) and folate deficiency leads to increased formic acid excretion in the rat. Toxicology 146: 123-136.
- Dowsley, T; Ulreich, J; Bolton, J; Park, S; Forkert, P. (1996). CYP2E1-dependent bioactivation of 1,1-dichloroethylene in murine lung: Formation of reactive intermediates and glutathione conjugates. Toxicol Appl Pharmacol 139: 42-48. <http://dx.doi.org/10.1006/taap.1996.0141>.
- Drake, V; Koprowski, S; Hu, N; Smith, S; Lough, J. (2006a). Cardiogenic effects of trichloroethylene and trichloroacetic acid following exposure during heart specification of avian development. Toxicol Sci 94: 153-162. <http://dx.doi.org/10.1093/toxsci/kfl083>.
- Drake, V; Koprowski, S; Lough, J; Hu, N; Smith, S. (2006b). Trichloroethylene exposure during cardiac valvuloseptal morphogenesis alters cushion formation and cardiac hemodynamics in the avian embryo. Environ Health Perspect 114: 842-847.
- Driscoll, T; Hamdan, H; Wang, G; Wright, P; Stacey, N. (1992). Concentrations of individual serum or plasma bile acids in workers exposed to chlorinated aliphatic hydrocarbons. Br J Ind Med 49: 700-705.
- Dryver, E; Brandt, L; Kauppinen, T; Olsson, H. (2004). Occupational exposures and non-Hodgkin's lymphoma in Southern Sweden. Int J Occup Environ Health 10: 13-21.
- Du, C and Wang, J. (1998). Increased morbidity odds ratio of primary liver cancer and cirrhosis of the liver among vinyl chloride monomer workers. Occup Environ Med 55: 528-532.
- Duffel, M and Jakoby, W. (1982). Cysteine S-conjugate N-acetyltransferase from rat kidney microsomes. Mol Pharmacol 21: 444-448.
- Dugard, P. (2000). Effects of trichloroethylene (TCE) on an in vitro chick atrioventricular canal culture. Toxicol Sci 56: 437-438.

- Dumas, S; Parent, M; Siemiatycki, J; Brisson, J. (2000). Rectal cancer and occupational risk factors: A hypothesis-generating, exposure-based case-control study. *Int J Cancer* 87: 874-879. [http://dx.doi.org/10.1002/1097-0215\(20000915\)87:6<874::AID-IJC18>3.0.CO;2-L](http://dx.doi.org/10.1002/1097-0215(20000915)87:6<874::AID-IJC18>3.0.CO;2-L).
- DuTeaux, S; Hengel, M; DeGroot, D; Jelks, K; Miller, M. (2003). Evidence for trichloroethylene bioactivation and adduct formation in the rat epididymis and efferent ducts. *Biol Reprod* 69: 771-779. <http://dx.doi.org/10.1095/biolreprod.102.014845>.
- DuTeaux, S; Berger, T; Hess, R; Sartini, B; Miller, M. (2004a). Male reproductive toxicity of trichloroethylene: Sperm protein oxidation and decreased fertilizing ability. *Biol Reprod* 70: 1518-1526. <http://dx.doi.org/10.1095/biolreprod.103.022210>.
- DuTeaux, S; Newman, J; Morisseau, C; Fairbairn, E; Jelks, K; Hammock, B; Miller, M. (2004b). Epoxide hydrolases in the rat epididymis: possible roles in xenobiotic and endogenous fatty acid metabolism. *Toxicol Sci* 78: 187-195. <http://dx.doi.org/10.1093/toxsci/kfh066>.
- Duval, S and Tweedie, R. (2000). A nonparametric “trim and fill” method of accounting for publication bias in meta-analysis. *J Am Stat Assoc* 95: 89-98.
- Eacho, P; Lanier, T; Brodhecker, C. (1991). Hepatocellular DNA synthesis in rats given peroxisome proliferating agents: Comparison of WY-14,643 to clofibrate acid, nafenopin and LY171883. *Carcinogenesis* 12: 1557-1561.
- Eastin, W; Mennear, J; Tenant, R; Stoll, R; Branstetter, D; Bucher, J; McCullough, B; Binder, R; Spalding, J; Mahler, J. (2001). Tg.AC genetically altered mouse: assay working group overview of available data. *Toxicol Pathol* 29 Suppl: 60-80. <http://dx.doi.org/10.1080/019262301753178483>.
- Eastmond, D and Tucker, J. (1989). Identification of aneuploidy-inducing agents using cytokinesis-blocked human lymphocytes and an antikinetochore antibody. *Environ Mol Mutagen* 13: 34-43.
- Eastmond, D; Hartwig, A; Anderson, D; Anwar, W; Cimino, M; Dobrev, I; Douglas, G; Nohmi, T; Phillips, D; Vickers, C. (2009). Mutagenicity testing for chemical risk assessment: Update of the WHO/IPCS Harmonized Scheme. *Mutagenesis* 24: 341-349. <http://dx.doi.org/10.1093/mutage/ gep014>.
- EFSA (European Food Safety Authority). (2006). Opinion of the scientific panel on food additives, flavourings, processing aids and materials in contact with food (AFC) on a request from the commission related to a new long-term carcinogenicity study on aspartame. *EFSA J* 4: 1-44. <http://dx.doi.org/10.2903/j.efsa.2006.356>.
- Egger, M; Smith, G; Schneider, M; Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *Br Med J* 315: 629-634.
- EHS. (Environmental Health Strategies). (1997). Final Report. Correspondence from Paul A. Cammer, Ph.D., Trichloroethylene Issues Group, to Cheryl Siegel Scott, US Environmental Protection Agency. December 22, 1997. Menlo Park, CA.
- Eichenlaub-Ritter, U; Baart, E; Yin, H; Betzendahl, I. (1996). Mechanisms of spontaneous and chemically-induced aneuploidy in mammalian oogenesis: basis of sex-specific differences in response to aneugens and the necessity for further tests. *Mutat Res* 372: 279-294.
- El-Serag, H. (2004). Hepatocellular carcinoma: Recent trends in the United States. *Gastroenterology* 127: S27-S34. <http://dx.doi.org/10.1053/j.gastro.2004.09.013>.
- El-Serag, H. (2007). Epidemiology of hepatocellular carcinoma in USA. *Hepatology Research* 37: S88-S94. <http://dx.doi.org/10.1111/j.1872-034X.2007.00168.x>.
- El-Serag, H and Rudolph, K. (2007). Hepatocellular carcinoma: epidemiology and molecular carcinogenesis. *Gastroenterology* 132: 2557-2576. <http://dx.doi.org/10.1053/j.gastro.2007.04.061>.
- El Ghawabi, S; Mansoor, M; El Gamel, M; El Saharti, A; El Enany, F. (1973). Chronic trichloroethylene exposure. *J Egypt Med Assoc* 56: 715-724.
- Elcombe, C. (1985). Species differences in carcinogenicity and peroxisome proliferation due to trichloroethylene: A biochemical human hazard assessment. *Arch Toxicol Suppl* 8: 6-17.
- Elcombe, C; Rose, M; Pratt, I. (1985). Biochemical, histological, and ultrastructural changes in rat and mouse liver following the administration of trichloroethylene: Possible relevance to species

- differences in hepatocarcinogenicity. *Toxicol Appl Pharmacol* 79: 365-376.
[http://dx.doi.org/10.1016/0041-008X\(85\)90135-8](http://dx.doi.org/10.1016/0041-008X(85)90135-8).
- Elfarra, A and Anders, M. (1984). Renal processing of glutathione conjugates role in nephrotoxicity. *Biochem Pharmacol* 33: 3729-3732.
- Elfarra, A; Jakobson, I; Anders, M. (1986). Mechanism of S-(1,2-dichlorovinyl)glutathione-induced nephrotoxicity. *Biochem Pharmacol* 35: 283-288.
- Elfarra, A; Lash, L; Anders, M. (1987). Alpha-ketoacids stimulate rat renal cysteine conjugate beta-lyase activity and potentiate the cytotoxicity of S-(1,2-dichlorovinyl)-L-cysteine. *Mol Pharmacol* 31: 208-212.
- Elfarra, A and Hwang, I. (1990). In vivo metabolites of S-(2-benzothiazolyl)-L-cysteine as markers of in vivo cysteine conjugate beta-lyase and thiol glucuronosyl transferase activities. *Drug Metab Dispos* 18: 917-922.
- Elfarra, A; Krause, R; Last, A; Lash, L; Parker, J. (1998). Species- and sex-related differences in metabolism of trichloroethylene to yield chloral and trichloroethanol in mouse, rat, and human liver microsomes. *Drug Metab Dispos* 26: 779-785.
- El-Hage J. (2007). Center for Drug Evaluation and Research. U.S. Food and Drug Administration. Peroxisome proliferator-activated receptor (PPAR) antagonists: preclinical and clinical cardiac safety considerations.
<http://www.fda.gov/downloads/AboutFDA/CentersOffices/CDER/ucm119071.pdf>
- Elizur, A; Adair-Kirk, T; Kelley, D; Griffin, G; Demello, D; Senior, R. (2008). Tumor necrosis factor-alpha from macrophages enhances LPS-induced clara cell expression of keratinocyte-derived chemokine. *Am J Respir Cell Mol Biol* 38: 8-15. <http://dx.doi.org/10.1165/rcmb.2007-0203OC>.
- Elofsson, S; Gamberale, F; Hindmarsh, T; Iregren, A; Isaksson, A; Johnsson, I; Knave, B; Lydahl, E; Mindus, P; Persson, H; Philipson, B; Steby, M; Struve, G; Soderman, E; Wennberg, A; Widén, L. (1980). Exposure to organic solvents: A cross-sectional epidemiologic investigation on occupationally exposed car and industrial spray painters with special reference to the nervous system. *Scand J Work Environ Health* 6: 239-273.
- Elovaara, E; Hemminki, K; Vainio, H. (1979). Effects of methylene chloride, trichloroethane, trichloroethylene, tetrachloroethylene and toluene on the development of chick embryos. *Toxicology* 12: 111-119. [http://dx.doi.org/10.1016/0300-483X\(79\)90037-4](http://dx.doi.org/10.1016/0300-483X(79)90037-4).
- Emmert, B; Bünger, J; Keuch, K; Müller, M; Emmert, S; Hallier, E; Westphal, G. (2006). Mutagenicity of cytochrome P450 2E1 substrates in the Ames test with the metabolic competent *S. typhimurium* strain YG7108pin3ERb5. *Toxicology* 228: 66-76.
<http://dx.doi.org/10.1016/j.tox.2006.08.013>.
- Engel, L; Vaughan, T; Gammon, M; Chow, W; Risch, H; Dubrow, R; Mayne, S; Rotterdam, H; Schoenberg, J; Stanford, J; West, A; Blot, W; Fraumeni, J. (2002). Occupation and risk of esophageal and gastric cardia adenocarcinoma. *Am J Ind Med* 42: 11-22.
<http://dx.doi.org/10.1002/ajim.10077>.
- Epstein, D; Nolen, G; Randall, J; Christ, S; Read, E; Stober, J; Smith, M. (1992). Cardiopathic effects of dichloroacetate in the fetal Long-Evans rat. *Teratology* 46: 225-235.
<http://dx.doi.org/10.1002/tera.1420460306>.
- Ertle, T; Henschler, D; Mueller, G; Spassowski, M. (1972). Metabolism of trichloroethylene in man. I. The significance of trichloroethanol in long-term exposure conditions. *Arch Toxicol* 29: 171-188.
<http://dx.doi.org/10.1007/BF00315598>.
- Evans, M; Chiu, W; Okino, M; Caldwell, J. (2009). Development of an updated PBPK model for trichloroethylene and metabolites in mice, and its application to discern the role of oxidative metabolism in TCE-induced hepatomegaly. *Toxicol Appl Pharmacol* 236: 329-340.
<http://dx.doi.org/10.1016/j.taap.2009.02.013>.
- Evert, M; Schneider-Stock, R; Dombrowski, F. (2003). Apolipoprotein A-IV mRNA overexpression in early preneoplastic hepatic foci induced by low-number pancreatic islet transplants in

- streptozotocin-diabetic rats. Pathol Res Pract 199: 373-379. <http://dx.doi.org/10.1078/0344-0338-00433>.
- Eyre, R; Stevens, D; Parker, J; Bull, R. (1995a). Renal activation of trichloroethene and S-(1,2-dichlorovinyl)-L-cysteine and cell proliferative responses in the kidneys of F344 rats and B6C3F1 mice. J Toxicol Environ Health 46: 465-481.
- Eyre, R; Stevens, D; Parker, J; Bull, R. (1995b). Acid-labile adducts to protein can be used as indicators of the cysteine S-conjugate pathway of trichloroethene metabolism. J Toxicol Environ Health 46: 443-464.
- Fagliano, J; Berry, M; Bove, F; Burke, T. (1990). Drinking water contamination and the incidence of leukemia: An ecologic study. Am J Public Health 80: 1209-1212.
- Fahrig, R; Madle, S; Baumann, H. (1995). Genetic toxicology of trichloroethylene (TCE). Mutat Res 340: 1-36.
- Fairman, M; Scott, B; Toothill, C; Losowsky, M. (1977). Jejunal mucosal gamma glutamyl transferase activity in coeliac disease. Gut 18: 484-487.
- Fan, A. (1988). Trichloroethylene: water contamination and health risk assessment. Rev Environ Contam Toxicol 101: 55-92.
- Fang, H; Strom, S; Cai, H; Falany, C; Kocarek, T; Runge-Morris, M. (2005). Regulation of human hepatic hydroxysteroid sulfotransferase gene expression by the peroxisome proliferator-activated receptor alpha transcription factor. Mol Pharmacol 67: 1257-1267.
<http://dx.doi.org/10.1124/mol.104.005389>.
- Farazi, P and Depinho, R. (2006). Hepatocellular carcinoma pathogenesis: From genes to environment. Nat Rev Cancer 6: 674-687. <http://dx.doi.org/10.1038/nrc1934>.
- Fattovich, G; Stroffolini, T; Zagni, I; Donato, F. (2004). Hepatocellular carcinoma in cirrhosis: incidence and risk factors. Gastroenterology 127: S35-S50. <http://dx.doi.org/10.1053/j.gastro.2004.09.014>.
- Fechter, L; Liu, Y; Herr, D; Crofton, K. (1998). Trichloroethylene ototoxicity: Evidence for a cochlear origin. Toxicol Sci 42: 28-35. <http://dx.doi.org/10.1093/toxsci/42.1.28>.
- Fechter, L; Gearhart, C; Fulton, S; Campbell, J; Fisher, J; Na, K; Cocker, D; Nelson-Miller, A; Moon, P; Pouyatos, B. (2007). Promotion of noise-induced cochlear injury by toluene and ethylbenzene in the rat. Toxicol Sci 98: 542-551.
- Feinberg, A; Ohlsson, R; Henikoff, S. (2006). The epigenetic progenitor origin of human cancer. Nat Rev Genet 7: 21-33. <http://dx.doi.org/10.1038/nrg1748>.
- Feitelson, M; Sun, B; Satiroglu Tufan, N; Liu, J; Pan, J; Lian, Z. (2002). Genetic mechanisms of hepatocarcinogenesis. Oncogene 21: 2593-2604. <http://dx.doi.org/10.1038/sj.onc.1205434>.
- Feldman, R; Chirico-Post, J; Proctor, S. (1988). Blink reflex latency after exposure to trichloroethylene in well water. Arch Environ Health 43: 143-148. <http://dx.doi.org/10.1080/00039896.1988.9935843>.
- Feldman, R; Niles, C; Proctor, S; Jabre, J. (1992). Blink reflex measurement of effects of trichloroethylene exposure on the trigeminal nerve. Muscle Nerve 15: 490-495.
<http://dx.doi.org/10.1002/mus.880150413>.
- Ferlay, J; Bray, F; Pisani, P; Parkin, D. (2004). GLOBOCAN 2002: Cancer incidence, mortality and prevalence worldwide. Lyon, France: International Agency for Research on Cancer.
- Ferlay, J; Shin, H; Bray, F; Mathers, C; Parkin, D. (2008). GLOBOCAN 2008, cancer incidence and mortality worldwide: IARC CancerBase no. 10. Lyon, France: International Agency for Research on Cancer. <http://globocan.iarc.fr>.
- Fernandez, J; Droz, P; Humbert, B; Caperos, J. (1977). Trichloroethylene exposure: Simulation of uptake, excretion, and metabolism using a mathematical model. Br J Ind Med 34: 43-55.
- Ferreira-Gonzalez, A; DeAngelo, A; Nasim, S; Garrett, C. (1995). Ras oncogene activation during hepatocarcinogenesis in B6C3F1 male mice by dichloroacetic and trichloroacetic acids. Carcinogenesis 16: 495-500. <http://dx.doi.org/10.1093/carcin/16.3.495>.
- Fevotte, J; Charbotel, B; Muller-Beauté, P; Martin, J; Hours, M; Bergeret, A. (2006). Case-control study on renal cell cancer and occupational exposure to trichloroethylene. Part I: Exposure assessment. Ann Occup Hyg 50: 765-775. <http://dx.doi.org/10.1093/annhyg/mel040>.

- FIFRA SAP. (Federal Insecticide, Fungicide, and Rodenticide Act Science Advisory Panel). (2004). A Set of Scientific Issues Being Considered by the Environmental Protection Agency Regarding Proposed Science Policy: PPAR-a Agonist-Mediated Hepatocarcinogenesis in Rodents and Relevance to Human Health Risk Assessment. Minutes No. 2003-05 (5 March 2004),. <http://www.epa.gov/scipoly/sap/meetings/2003/december9/meetingminutes.pdf>.
- Figgs, L; Dosemeci, M; Blair, A. (1995). United States non-Hodgkin's lymphoma surveillance by occupation 1984-1989: a twenty-four state death certificate study. Am J Ind Med 27: 817-835.
- Filipsson, A and Victorin, K. (2003). Comparison of available benchmark dose softwares and models using trichloroethylene as a model substance. Regul Toxicol Pharmacol 37: 343-355.
- Filser, J and Bolt, H. (1979). Pharmacokinetics of halogenated ethylenes in rats. Arch Toxicol 42: 123-136. <http://dx.doi.org/10.1007/BF00316492>.
- Finckh, A; Cooper, G; Chibnik, L; Costenbader, K; Watts, J; Pankey, H; Fraser, P; Karlson, E. (2006). Occupational silica and solvent exposures and risk of systemic lupus erythematosus in urban women. Arthritis Rheum 54: 3648-3654. <http://dx.doi.org/10.1002/art.22210>.
- Fiserova-Bergerova, V; Tichy, M; Di Carlo, F. (1984). Effects of biosolubility on pulmonary uptake and disposition of gases and vapors of lipophilic chemicals. Drug Metab Rev 15: 1033-1070.
- Fisher, J; Whittaker, T; Taylor, D; Clewell, H, III; Andersen, M. (1989). Physiologically based pharmacokinetic modeling of the pregnant rat: A multiroute exposure model for trichloroethylene and its metabolite, trichloroacetic acid. Toxicol Appl Pharmacol 99: 395-414. [http://dx.doi.org/10.1016/0041-008X\(89\)90149-X](http://dx.doi.org/10.1016/0041-008X(89)90149-X).
- Fisher, J; Whittaker, T; Taylor, D; Clewell, H, III; Andersen, M. (1990). Physiologically based pharmacokinetic modeling of the lactating rat and nursing pup: A multiroute exposure model for trichloroethylene and its metabolite, trichloroacetic acid. Toxicol Appl Pharmacol 102: 497-513.
- Fisher, J; Gargas, M; Allen, B; Andersen, M. (1991). Physiologically based pharmacokinetic modeling with trichloroethylene and its metabolite, trichloroacetic acid, in the rat and mouse. Toxicol Appl Pharmacol 109: 183-195.
- Fisher, J and Allen, B. (1993). Evaluating the risk of liver cancer in humans exposed to trichloroethylene using physiological models. Risk Anal 13: 87-95.
- Fisher, J; Mahle, D; Bankston, L; Greene, R; Gearhart, J. (1997). Lactational transfer of volatile chemicals in breast milk. Am Ind Hyg Assoc J 58: 425-431. <http://dx.doi.org/10.1080/15428119791012667>.
- Fisher, J; Mahle, D; Abbas, R. (1998). A human physiologically based pharmacokinetic model for trichloroethylene and its metabolites, trichloroacetic acid and free trichloroethanol. Toxicol Appl Pharmacol 152: 339-359. <http://dx.doi.org/10.1006/taap.1998.8486>.
- Fisher, J. (2000). Physiologically based pharmacokinetic models for trichloroethylene and its oxidative metabolites. Environ Health Perspect 108: 265-273.
- Fisher, J; Channel, S; Eggers, J; Johnson, P; MacMahon, K; Goodyear, C; Sudberry, G; Warren, D; Latendresse, J; Graeter, L. (2001). Trichloroethylene, trichloroacetic acid, and dichloroacetic acid: Do they affect fetal rat heart development. Int J Toxicol 20: 257-267.
- Fitzgerald, J; Sanyer, J; Schardein, J; Lake, R; McGuire, E; de la Iglesia, F. (1981). Carcinogen bioassay and mutagenicity studies with the hypolipidemic agent gemfibrozil. J Natl Cancer Inst 67: 1105-1116.
- Fleming-Jones, M and Smith, R. (2003). Volatile organic compounds in foods: A five year study. J Agric Food Chem 51: 8120-8127. <http://dx.doi.org/10.1021/jf0303159>.
- Flodin, U; Fredriksson, M; Persson, B. (1987). Multiple myeloma and engine exhausts, fresh wood, and creosote: A case-referent study. Am J Ind Med 12: 519-529.
- Flood, T. (1997a). Case-referent study of childhood leukemia in Maricopa County, Arizona: 1965-1990. Phoenix, AZ: Arizona Dept. of Health Services, Bureau of Epidemiology and Disease Control Services, Office of Chronic Disease Epidemiology, Office of Environmental Health.

- Flood, TC, CA. (1988). Report on mortality in Maricopa County 1966-1986. Phoenix, AZ: Arizona Department of Health Services, Division of Disease Prevention, Office of Chronic Disease Epidemiology, Office of Risk Assessment and Investigation.
- Flood, TJM, F. J. Vertz, P. D. Laubham, K. A. Porter, R. S. Aickinm, M. (1997b). Case-referent study of childhood leukemia in Maricopa County, Arizona: 1965-1990. Phoenix, AZ: Arizona Dept. of Health Services, Bureau of Epidemiology and Disease Control Services, Office of Chronic Disease Epidemiology, Office of Environmental Health.
- Forand, S. (2004). Leukaemia incidence among workers in the shoe and boot manufacturing industry: A case-control study. Environ Health 3: 7. <http://dx.doi.org/10.1186/1476-069X-3-7>.
- Ford, E; Rhodes, S; McDiarmid, M; Schwartz, S; Brown, J. (1995). Deaths from acute exposure to trichloroethylene. J Occup Environ Med 37: 749-754.
- Forkert, P-G; Baldwin, R; Millen, B; Lash, L; Putt, D; Shultz, M; Collins, K. (2005). Pulmonary bioactivation of trichloroethylene to chloral hydrate: Relative contributions of CYP2E1, CYP2F, and CYP2B1. Drug Metab Dispos 33: 1429-1437. <http://dx.doi.org/10.1124/dmd.105.005074>.
- Forkert, P-G; Millen, B; Lash, L; Putt, D; Ghanayem, B. (2006). Pulmonary bronchiolar cytotoxicity and formation of dichloroacetyl lysine protein adducts in mice treated with trichloroethylene. J Pharmacol Exp Ther 316: 520-529. <http://dx.doi.org/10.1124/jpet.105.093062>.
- Forkert, P; Sylvestre, P; Poland, J. (1985). Lung injury induced by trichloroethylene. Toxicology 35: 143-160.
- Forkert, P and Birch, D. (1989). Pulmonary toxicity of trichloroethylene in mice: covalent binding and morphological manifestations. Drug Metab Dispos 17: 106-113.
- Forkert, P and Forkert, L. (1994). Trichloroethylene induces pulmonary fibrosis in mice. Can J Physiol Pharmacol 72: 205-210.
- Forkert, P. (1999a). In vivo formation and localization of 1,1-dichloroethylene epoxide in murine liver: Identification of its glutathione conjugate 2-S-glutathionyl acetate. J Pharmacol Exp Ther 290: 1299-1306.
- Forkert, P. (1999b). 1,1-Dichloroethylene-induced Clara cell damage is associated with in situ formation of the reactive epoxide. Immunohistochemical detection of its glutathione conjugate. Am J Respir Cell Mol Biol 20: 1310-1318.
- Forkert, P; Malkinson, A; Rice, P; Moussa, M. (1999). Diminished CYP2E1 expression and formation of 2-S-glutathionyl acetate, a glutathione conjugate derived from 1,1-dichloroethylene epoxide, in murine lung tumors. Drug Metab Dispos 27: 68-73.
- Forkert, P; Lash, L; Nadeau, V; Tardif, R; Simmonds, A. (2002). Metabolism and toxicity of trichloroethylene in epididymis and testis. Toxicol Appl Pharmacol 182: 244-254.
- Forkert, P; Lash, L; Tardif, R; Tanphaichitr, N; Vandevoot, C; Moussa, M. (2003). Identification of trichloroethylene and its metabolites in human seminal fluid of workers exposed to trichloroethylene. Drug Metab Dispos 31: 306-311.
- Fort, D; Rayburn, J; DeYoung, D; Bantle, J. (1991). Assessing the efficacy of an Aroclor 1254-induced exogenous metabolic activation system for FETAX. Drug Chem Toxicol 14: 143-160. <http://dx.doi.org/10.3109/01480549109017873>.
- Fort, D; Stover, E; Rayburn, J; Hull, M; Bantle, J. (1993). Evaluation of the developmental toxicity of trichloroethylene and detoxification metabolites using Xenopus. Birth Defects Res B Dev Reprod Toxicol 13: 35-45.
- Fox, A; Sullivan, B; Buffini, J; Neichin, M; Nicora, R; Hoehler, F; O'Rourke, R; Stoltz, R. (1996a). Reduction of serum lactate by sodium dichloroacetate, and human pharmacokinetic-pharmacodynamic relationships. J Pharmacol Exp Ther 279: 686-693.
- Fox, A; Yang, X; Murli, H; Lawlor, T; Cifone, M; Reno, F. (1996b). Absence of mutagenic effects of sodium dichloroacetate. Fundam Appl Toxicol 32: 87-95. <http://dx.doi.org/10.1006/faat.1996.0110>.

- Fox, T; Schumann, A; Watanabe, P; Yano, B; Maher, V; McCormick, J. (1990). Mutational analysis of the H-ras oncogene in spontaneous C57BL/6 x C3H/He mouse liver tumors and tumors induced with genotoxic and nongenotoxic hepatocarcinogens. *Cancer Res* 50: 4014-4019.
- Fraga, M; Herranz, M; Espada, J; Ballestar, E; Paz, M; Ropero, S; Erkek, E; Bozdogan, O; Peinado, H; Niveleau, A; Mao, J; Balmain, A; Cano, A; Esteller, M. (2004). A mouse skin multistage carcinogenesis model reflects the aberrant DNA methylation patterns of human tumors. *Cancer Res* 64: 5527-5534. <http://dx.doi.org/10.1158/0008-5472.CAN-03-4061>.
- Fraga, M; Ballestar, E; Villar-Garea, A; Boix-Chornet, M; Espada, J; Schotta, G; Bonaldi, T; Haydon, C; Ropero, S; Petrie, K; Iyer, N; Pérez-Rosado, A; Calvo, E; Lopez, J; Cano, A; Calasanz, M; Colomer, D; Piris, M; Ahn, N; Imhof, A; Caldas, C; Jenuwein, T; Esteller, M. (2005). Loss of acetylation at Lys16 and trimethylation at Lys20 of histone H4 is a common hallmark of human cancer. *Nat Genet* 37: 391-400. <http://dx.doi.org/10.1038/ng1531>.
- Franco, A; Costoya, M; Roca, E. (2007). Estimating risk during showering exposure to VOCs of workers in a metal-degreasing facility. *J Toxicol Environ Health A* 70: 627-637. <http://dx.doi.org/10.1080/15287390600974551>.
- Franco, J; Teixeira, A; Meotti, F; Ribas, C; Stringari, J; Garcia Pomblum, S; Moro, A; Bohrer, D; Bairros, A; Dafre, A; Santos, A; Farina, M. (2006). Cerebellar thiol status and motor deficit after lactational exposure to methylmercury. *Environ Res* 102: 22-28. <http://dx.doi.org/10.1016/j.envres.2006.02.003>.
- Fredriksson, A; Danielsson, B; Eriksson, P. (1993). Altered behaviour in adult mice orally exposed to tri- and tetrachloroethylene as neonates. *Toxicol Lett* 66: 13-19. [http://dx.doi.org/10.1016/0378-4274\(93\)90074-8](http://dx.doi.org/10.1016/0378-4274(93)90074-8).
- Fredriksson, M; Bengtsson, N; Hardell, L; Axelson, O. (1989). Colon cancer, physical activity, and occupational exposures. A case-control study. *Cancer* 63: 1838-1842.
- Fritschi, L and Siemiatycki, J. (1996a). Lymphoma, myeloma and occupation: Results of a case-control study. *Int J Cancer* 67: 498-503. [http://dx.doi.org/10.1002/\(SICI\)1097-0215\(19960807\)67:4<498::AID-IJC6>3.0.CO;2-N](http://dx.doi.org/10.1002/(SICI)1097-0215(19960807)67:4<498::AID-IJC6>3.0.CO;2-N).
- Fritschi, L and Siemiatycki, J. (1996b). Melanoma and occupation: Results of a case-control study. *Occup Environ Med* 53: 168-173.
- Fritschi, L; Siemiatycki, J; Richardson, L. (1996). Self-assessed versus expert-assessed occupational exposures. *Am J Epidemiol* 144: 521-527.
- Fritschi, L; Nadon, L; Benke, G; Lakhani, R; Latreille, B; Parent, M; Siemiatycki, J. (2003). Validation of expert assessment of occupational exposures. *Am J Ind Med* 43: 519-522. <http://dx.doi.org/10.1002/ajim.10208>.
- Froyland, L; Madsen, L; Vaagenes, H; Totland, G; Auwerx, J; Kryvi, H; Staels, B; Berge, R. (1997). Mitochondrion is the principal target for nutritional and pharmacological control of triglyceride metabolism. *J Lipid Res* 38: 1851-1858.
- Fujita, J; Marino, M; Wada, H; Jungbluth, A; Mackrell, P; Rivadeneira, D; Stapleton, P; Daly, J. (2001). Effect of TNF gene depletion on liver regeneration after partial hepatectomy in mice. *Surgery* 129: 48-54.
- Fukuda, K; Takemoto, K; Tsuruta, H. (1983). Inhalation carcinogenicity of trichloroethylene in mice and rats. *Ind Health* 21: 243-254.
- Furge, K; Dykema, K; Petillo, D; Westphal, M; Zhang, Z; Kort, E; Teh, B. (2007). Combining differential expression, chromosomal and pathway analyses for the molecular characterization of renal cell carcinoma. *Canadian Urological Association Journal* 1: S21-S27.
- Furnus, C; Ulrich, M; Terreros, M; Dulout, F. (1990). The induction of aneuploidy in cultured Chinese hamster cells by propionaldehyde and chloral hydrate. *Mutagenesis* 5: 323-326.
- Fürstenberger, G and Senn, H. (2002). Insulin-like growth factors and cancer. *Lancet Oncol* 3: 298-302.
- Fuscoe, J; Afshari, A; George, M; DeAngelo, A; Tice, R; Salman, T; Allen, J. (1996). In vivo genotoxicity of dichloroacetic acid: Evaluation with the mouse peripheral blood micronucleus

- assay and the single cell gel assay. Environ Mol Mutagen 27: 1-9.
[http://dx.doi.org/10.1002/\(SICI\)1098-2280\(1996\)27:1<1::AID-EM1>3.0.CO;2-L](http://dx.doi.org/10.1002/(SICI)1098-2280(1996)27:1<1::AID-EM1>3.0.CO;2-L).
- Gaglio, P; Liu, H; Dash, S; Cheng, S; Dunne, B; Ratterree, M; Baskin, G; Blanchard, J; Bohm, R; Theise, N; LaBrecque, D. (2002). Liver regeneration investigated in a non-human primate model (*Macaca mulatta*). J Hepatol 37: 625-632. [http://dx.doi.org/10.1016/S0168-8278\(02\)00262-3](http://dx.doi.org/10.1016/S0168-8278(02)00262-3).
- Gagnaire, F; Marignac, B; Langlais, C; Bonnet, P. (2001). Ototoxicity in rats exposed to ortho-, meta- and para-xylene vapours for 13 weeks. Pharmacol Toxicol 89: 6-14. <http://dx.doi.org/10.1111/j.1600-0773.2001.890102.x>.
- Gallou, C; Longuemaux, S; Deloménie, C; Méjean, A; Martin, N; Martinet, S; Palais, G; Bouvier, R; Droz, D; Krishnamoorthy, R; Junien, C; Béroud, C; Dupret, J. (2001). Association of GSTT1 non-null and NAT1 slow/rapid genotypes with von Hippel-Lindau tumour suppressor gene transversions in sporadic renal cell carcinoma. Pharmacogenetics 11: 521-535.
- Galloway, S; Armstrong, M; Reuben, C; Colman, S; Brown, B; Cannon, C; Bloom, A; Nakamura, F; Ahmed, M; Duk, S; Rimpo, J; Margolin, B; Resnick, M; Anderson, B; Zeiger, E. (1987). Chromosome aberrations and sister chromatid exchanges in Chinese hamster ovary cells: Evaluations of 108 chemicals. Environ Mol Mutagen 10: 1-175.
<http://dx.doi.org/10.1002/em.2850100502>.
- Gama, C and Meira, J. (1978). Occupational acro-osteolysis. J Bone Joint Surg Am 60: 86-90.
- Gamberale, G; Annwall, G; Olson, B. (1976). Exposure to trichloroethylene III. Psychological functions. Scand J Work Environ Health 4: 220-224.
- Gandhi, M; Aweeka, F; Greenblatt, R; Blaschke, T. (2004). Sex differences in pharmacokinetics and pharmacodynamics. Annu Rev Pharmacol Toxicol 44: 499-523.
<http://dx.doi.org/10.1146/annurev.pharmtox.44.101802.121453>.
- Gandillet, A; Alexandre, E; Royer, C; Cinqualbre, J; Jaeck, D; Richert, L. (2003). Hepatocyte ploidy in regenerating livers after partial hepatectomy, drug-induced necrosis, and cirrhosis. Eur Surg Res 35: 148-160. <http://dx.doi.org/10.1159/000070044>.
- Garabrant, D; Held, J; Langholz, B; Bernstein, L. (1988). Mortality of aircraft manufacturing workers in southern California. Am J Ind Med 13: 683-693. <http://dx.doi.org/10.1002/ajim.4700130607>.
- Garabrant, D; Lacey JV, J; Laing, T; Gillespie, B; Mayes, M; Cooper, B; Schottenfeld, D. (2003). Scleroderma and solvent exposure among women. Am J Epidemiol 157: 493-500.
<http://dx.doi.org/10.1093/aje/kwf223>.
- Garfinkel, L and Stellman, S. (1988). Smoking and lung cancer in women: findings in a prospective study. Cancer Res 48: 6951-6955.
- Gargas, M; Seybold, P; Andersen, M. (1988). Modeling the tissue solubilities and metabolic rate constant (Vmax) of halogenated methanes, ethanes, and ethylenes. Toxicol Lett 43: 235-256.
[http://dx.doi.org/10.1016/0378-4274\(88\)90031-8](http://dx.doi.org/10.1016/0378-4274(88)90031-8).
- Gargas, M; Burgess, R; Voisard, D; Cason, G; Andersen, M. (1989). Partition coefficients of low-molecular-weight volatile chemicals in various liquids and tissues. Toxicol Appl Pharmacol 98: 87-99.
- Gart, J; Chu, K; Tarone, R. (1979). Statistical issues in interpretation of chronic bioassay tests for carcinogenicity. J Natl Cancer Inst 62: 957-974.
- Garte, S; Gaspari, L; Alexandrie, A; Ambrosone, C; Autrup, H; Autrup, J; Baranova, H; Bathum, L; Benhamou, S; Boffetta, P; Bouchardy, C; Breskvar, K; Brockmoller, J; Cascorbi, I; Clapper, M; Coutelle, C; Daly, A; Dell'Omoo, M; Dolzan, V; Dresler, C; Fryer, A; Haugen, A; Hein, D; Hildesheim, A; Hirvonen, A; Hsieh, L; Ingelman-Sundberg, M; Kalina, I; Kang, D; Kihara, M; Kiyohara, C; Kremers, P; Lazarus, P; Le Marchand, L; Lechner, M; van Lieshout, E; London, S; Manni, J; Maugard, C; Morita, S; Nazar-Stewart, V; Noda, K; Oda, Y; Parl, F; Pastorelli, R; Persson, I; Peters, W; Rannug, A; Rebbeck, T; Risch, A; Roelandt, L; Romkes, M; Ryberg, D; Salagovic, J; Schoket, B; Seidegard, J; Shields, P; Sim, E; Sinnet, D; Strange, R; Stüber, I; Sugimura, H; To-Figueras, J; Vineis, P; Yu, M; Taioli, E. (2001). Metabolic gene polymorphism frequencies in control populations. Cancer Epidemiol Biomarkers Prev 10: 1239-1248.

- Gash, D; Rutland, K; Hudson, N; Sullivan, P; Bing, G; Cass, W; Pandya, J; Liu, M; Choi, D; Hunter, R; Gerhardt, G; Smith, C; Slevin, J; Prince, T. (2008). Trichloroethylene: Parkinsonism and complex 1 mitochondrial neurotoxicity. Ann Neurol 63: 184-192. <http://dx.doi.org/10.1002/ana.21288>.
- Ge, R; Wang, W; Kramer, P; Yang, S; Tao, L; Pereira, M. (2001a). Wy-14,643-induced hypomethylation of the c-myc gene in mouse liver. Toxicol Sci 62: 28-35.
- Ge, R; Yang, S; Kramer, P; Tao, L; Pereira, M. (2001b). The effect of dichloroacetic acid and trichloroacetic acid on DNA methylation and cell proliferation in B6C3F1 mice. J Biochem Mol Toxicol 15: 100-106. <http://dx.doi.org/10.1002/jbt.5>.
- Ge, R; Tao, L; Kramer, P; Cunningham, M; Pereira, M. (2002). Effect of peroxisome proliferators on the methylation and protein level of the c-myc protooncogene in B6C3F1 mice liver. J Biochem Mol Toxicol 16: 41-47. <http://dx.doi.org/10.1002/jbt.10019>.
- Gelderblom, W; Galendo, D; Abel, S; Swanevelder, S; Marasas, W; Wild, C. (2001). Cancer initiation by fumonisin B(1) in rat liver: Role of cell proliferation. Cancer Lett 169: 127-137.
- Gelfand, A and Smith, A. (1990). Sampling based approaches to calculating marginal densities. J Am Stat Assoc 85: 398 - 409.
- Gelman, A; Bois, F; Jiang, J. (1996). Physiological pharmacokinetic analysis using population modeling and informative prior distributions. J Am Stat Assoc 91: 1400-1412.
- Gelman, A; Carlin, J; Stern, H; Rubin, D. (2003). Bayesian data analysis: Second edition. Boca Raton, FL: Chapman and Hall/CRC Press.
- George, J; Reel, J; Myers, C; Lawton, A. (1985). Trichloroethylene: Reproduction and fertility assessment in CD-1 mice when administered in the feed. (NTP-86-068). pp. 242p. Research Triangle Park, NC: NIEHS.
- George, J; Reel, J; Myers, C; Lawton, A; Lamb, J, IV. (1986). Trichloroethylene: Reproduction and fertility assessment in F344 rats when administered in the feed. (NTP-86-085). Research Triangle Park, NC: National Institute of Environmental Health Sciences, National Toxicology Program.
- George, J; Byth, K; Farrell, G. (1995a). Age but not gender selectively affects expression of individual cytochrome P450 proteins in human liver. Biochem Pharmacol 50: 727-730.
- George, J; Price, C; Navarro, H; Marr, M; Myers, C; Hunter ES Schwetz, B; Shelby, M. (1995b). Developmental toxicity of thiophenol (THIO) in rats and rabbits. Toxicologist 15: 160.
- George, M; Moore, T; Kilburn, S; Olson, G; DeAngelo, A. (2000). Carcinogenicity of chloral hydrate administered in drinking water to the male F344/N rat and male B6C3F1 mouse. Toxicol Pathol 28: 610-618.
- Ghantous, H; Danielsson, B; Dencker, L; Gorczak, J; Vesterberg, O. (1986). Trichloroacetic acid accumulates in murine amniotic fluid after tri- and tetrachloroethylene inhalation. Acta Pharmacol Toxicol 58: 105-114. <http://dx.doi.org/10.1111/j.1600-0773.1986.tb00078.x>.
- Ghoshal, A and Farber, E. (1984). The induction of liver cancer by dietary deficiency of choline and methionine without added carcinogens. Carcinogenesis 5: 1367-1370.
- Giardino, N and Andelman, J. (1996). Characterization of the emissions of trichloroethylene, chloroform, and 1,2-dibromo-3-chloropropane in a full-size, experimental shower. J Expo Sci Environ Epidemiol 6: 413-423.
- Gibson, D; Aardema, M; Kerckaert, G; Carr, G; Brauninger, R; LeBoeuf, R. (1995). Detection of aneuploidy-inducing carcinogens in the Syrian hamster embryo (SHE) cell transformation assay. Mutat Res 343: 7-24. [http://dx.doi.org/10.1016/0165-1218\(95\)90058-6](http://dx.doi.org/10.1016/0165-1218(95)90058-6).
- Gilbert, K; Griffin, J; Pumford, N. (1999). Trichloroethylene activates CD4+ T cells: potential role in an autoimmune response. Drug Metab Rev 31: 901-916. <http://dx.doi.org/10.1081/DMR-100101945>.
- Gilbert, K; Whitlow, A; Pumford, N. (2004). Environmental contaminant and disinfection by-product trichloroacetaldehyde stimulates T cells in vitro. Int Immunopharmacol 4: 25-36. <http://dx.doi.org/10.1016/j.intimp.2003.10.001>.
- Gilbert, K; Pumford, N; Blossom, S. (2006). Environmental contaminant trichloroethylene promotes autoimmune disease and inhibits T-cell apoptosis in MRL(+/+) mice. J Immunotoxicol 3: 263-267. <http://dx.doi.org/10.1080/15476910601023578>.

- Gilkeson, G; Keil, D; Peden-Adams, M. (2004). Immune effects of trichloroethylene on autoimmune disease in mice. 87 - 98 PUBL- The Medical University of South Carolina Press.
- Gilks, W; Richardson, S; Spiegelhalter, D. (1995). Markov chain Monte Carlo in practice. Boca Raton, FL: Chapman & Hall/CRC Press.
- Giller, S; Le Curieux, F; Gauthier, L; Erb, F; Marzin, D. (1995). Genotoxicity assay of chloral hydrate and chloropicrin. Mutat Res 348: 147-152. [http://dx.doi.org/10.1016/0165-7992\(95\)90002-0](http://dx.doi.org/10.1016/0165-7992(95)90002-0).
- Giller, S; Le Curieux, F; Erb, F; Marzin, D. (1997). Comparative genotoxicity of halogenated acetic acids found in drinking water. Mutagenesis 12: 321-328.
- Ginsberg, G; Hattis, D; Sonawane, B; Russ, A; Banati, P; Kozlak, M; Smolenski, S; Goble, R. (2002). Evaluation of child/adult pharmacokinetic differences from a database derived from the therapeutic drug literature. Toxicol Sci 66: 185-200.
- Ginsberg, G; Hattis, D; Miller, R; Sonawane, B. (2004a). Pediatric pharmacokinetic data: Implications for environmental risk assessment for children. Pediatrics 113: 973-983.
- Ginsberg, G; Hattis, D; Sonawane, B. (2004b). Incorporating pharmacokinetic differences between children and adults in assessing children's risks to environmental toxicants. Toxicol Appl Pharmacol 198: 164-183. <http://dx.doi.org/10.1016/j.taap.2003.10.010>.
- Ginsberg, G; Hattis, D; Russ, A; Sonawane, B. (2005). Pharmacokinetic and pharmacodynamic factors that can affect sensitivity to neurotoxic sequelae in elderly individuals. Environ Health Perspect 113: 1243-1249. <http://dx.doi.org/10.1289/ehp.7568>.
- Gochfeld, M. (2007). Framework for gender differences in human and animal toxicology. Environ Res 104: 4-21. <http://dx.doi.org/10.1016/j.envres.2005.12.005>.
- Goel, S; Rao, G; Pandya, K; Shanker, R. (1992). Trichloroethylene toxicity in mice: A biochemical, hematological and pathological assessment. Indian J Exp Biol 30: 402-406.
- Goeptar, A; Commandeur, J; van Ommen, B; van Bladeren, P; Vermeulen, N. (1995). Metabolism and kinetics of trichloroethylene in relation to toxicity and carcinogenicity: Relevance of the mercapturic acid pathway. Chem Res Toxicol 8: 3-21.
- Goh, V; Chia, S; Ong, C. (1998). Effects of chronic exposure to low doses of trichloroethylene on steroid hormone and insulin levels in normal men. Environ Health Perspect 106: 41-44.
- Gold, L; Manley, N; Slone, T; Rohrbach, L; Garfinkel, G. (2005). Supplement to the Carcinogenic Potency Database (CPDB): results of animal bioassays published in the general literature through 1997 and by the National Toxicology Program in 1997-1998. Toxicol Sci 85: 747-808. <http://dx.doi.org/10.1093/toxsci/kfi161>.
- Gold, L; Stewart, P; Milliken, K; Purdue, M; Severson, R; Seixas, N; Blair, A; Hartge, P; Davis, S; De Roos, A. (2011). The relationship between multiple myeloma and occupational exposure to six chlorinated solvents. Occup Environ Med 68: 391-399. <http://dx.doi.org/10.1136/oem.2009.054809>.
- Gold, LS and Zeiger, E (Eds.). (1997). Handbook of carcinogenic potency and genotoxicity databases. Boca Raton, FL: CRC Press.
- Goldberg, M; Parent, M; Siemiatycki, J; Désy, M; Nadon, L; Richardson, L; Lakhani, R; Latreille, B; Valois, M. (2001). A case-control study of the relationship between the risk of colon cancer in men and exposures to occupational agents. Am J Ind Med 39: 531-546.
- Goldberg, S; Lebowitz, M; Graver, E; Hicks, S. (1990). An association of human congenital cardiac malformations and drinking water contaminants. J Am Coll Cardiol 16: 155-164.
- Goldsworthy, T and Popp, J. (1987). Chlorinated hydrocarbon-induced peroxisomal enzyme activity in relation to species and organ carcinogenicity. Toxicol Appl Pharmacol 88: 225-233. [http://dx.doi.org/10.1016/0041-008X\(87\)90008-1](http://dx.doi.org/10.1016/0041-008X(87)90008-1).
- Goldsworthy, T; Lyght, O; Burnett, V; Popp, J. (1988). Potential role of [alpha]-2[mu]-globulin, protein droplet accumulation, and cell replication in the renal carcinogenicity of rats exposed to trichloroethylene, perchloroethylene, and pentachloroethane. Toxicol Appl Pharmacol 96: 367-379. [http://dx.doi.org/10.1016/0041-008X\(88\)90095-6](http://dx.doi.org/10.1016/0041-008X(88)90095-6).

- Gollin, S. (2005). Mechanisms leading to chromosomal instability. Semin Cancer Biol 15: 33-42. <http://dx.doi.org/10.1016/j.semcancer.2004.09.004>.
- Gómez, M; Cocco, P; Dosemeci, M; Stewart, P. (1994). Occupational exposure to chlorinated aliphatic hydrocarbons: Job exposure matrix. Am J Ind Med 26: 171-183. <http://dx.doi.org/10.1002/ajim.4700260204>.
- Gonzalez-Leon, A; Merdink, J; Bull, R; Schultz, I. (1999). Effect of pre-treatment with dichloroacetic or trichloroacetic acid in drinking water on the pharmacokinetics of a subsequent challenge dose in B6C3F1 mice. Chem Biol Interact 123: 239-253.
- Gorla, G; Malhi, H; Gupta, S. (2001). Polyploidy associated with oxidative injury attenuates proliferative potential of cells. J Cell Sci 114: 2943-2951.
- Graham, D; Staffa, J; Shatin, D; Andrade, S; Schech, S; La Grenade, L; Gurwitz, J; Chan, K; Goodman, M; Platt, R. (2004). Incidence of hospitalized rhabdomyolysis in patients treated with lipid-lowering drugs. JAMA 292: 2585-2590. <http://dx.doi.org/10.1001/jama.292.21.2585>.
- Gramantieri, L; Melchiorri, C; Chieco, P; Gaiani, S; Stecca, B; Casali, A; Bolondi, L. (1996). Alteration of DNA ploidy and cell nuclearity in human hepatocellular carcinoma associated with HBV infection. J Hepatol 25: 848-853. [http://dx.doi.org/10.1016/S0168-8278\(96\)80288-1](http://dx.doi.org/10.1016/S0168-8278(96)80288-1).
- Grandjean, E; Munchinger, R; Turrian, V; Haas, P; Knoepfel, H-K; Rosenmund, H. (1955). Investigations into the effects of exposure to trichlorethylene in mechanical engineering. Br J Ind Med 12: 131-142.
- Grasl-Kraupp, B; Waldhör, T; Huber, W; Schulte-Hermann, R. (1993). Glutathione S-transferase isoenzyme patterns in different subtypes of enzyme-altered rat liver foci treated with the peroxisome proliferator nafenopin or with phenobarbital. Carcinogenesis 14: 2407-2412. <http://dx.doi.org/10.1093/carcin/14.11.2407>.
- Grawé, J; Nüsse, M; Adler, I-D. (1997). Quantitative and qualitative studies of micronucleus induction in mouse erythrocytes using flow cytometry: I: Measurement of micronucleus induction in peripheral blood polychromatic erythrocytes by chemicals with known and suspected genotoxicity. Mutagenesis 12: 1-8. <http://dx.doi.org/10.1093/mutage/12.1.1>.
- Green, L and Lash, T. (1999). Re: "Renal cell cancer correlated with occupational exposure to trichloroethylene". J Cancer Res Clin Oncol 125: 430-432.
- Green, T and Prout, M. (1985). Species differences in response to trichloroethylene: II. Biotransformation in rats and mice. Toxicol Appl Pharmacol 79: 401-411. [http://dx.doi.org/10.1016/0041-008X\(85\)90138-3](http://dx.doi.org/10.1016/0041-008X(85)90138-3).
- Green, T; Odum, J; Nash, J; Foster, J. (1990). Perchloroethylene-induced rat kidney tumors: An investigation of the mechanisms involved and their relevance to humans. Toxicol Appl Pharmacol 103: 77-89. [http://dx.doi.org/10.1016/0041-008X\(90\)90264-U](http://dx.doi.org/10.1016/0041-008X(90)90264-U).
- Green, T; Dow, J; Ellis, M; Foster, J; Odum, J. (1997a). The role of glutathione conjugation in the development of kidney tumours in rats exposed to trichloroethylene. Chem Biol Interact 105: 99-117.
- Green, T; Mainwaring, G; Foster, J. (1997b). Trichloroethylene-induced mouse lung tumors: Studies of the mode of action and comparisons between species. Fundam Appl Toxicol 37: 125-130. <http://dx.doi.org/10.1006/faat.1997.2312>.
- Green, T; Dow, J; Foster, J; Hext, P. (1998). Formic acid excretion in rats exposed to trichloroethylene: a possible explanation for renal toxicity in long-term studies. Toxicology 127: 39-47.
- Green, T. (2000). Pulmonary toxicity and carcinogenicity of trichloroethylene: Species differences and modes of action. Environ Health Perspect 108: 261-264.
- Green, T. (2003a). Trichloroacetic acid: Drinking water study in male and female B6C3F1, male PPARalpha knockout and male 129/sv wild type mice. (CTL/024741). Cheshire, UK: Central Toxicology Laboratory.
- Green, T. (2003b). The concentrations of trichloroacetic acid in blood following administration of trichloroacetic acid in drinking water. Cheshire, UK: Central Toxicology Laboratory.

- Green, T; Dow, J; Foster, J. (2003). Increased formic acid excretion and the development of kidney toxicity in rats following chronic dosing with trichloroethanol, a major metabolite of trichloroethylene. *Toxicology* 191: 109-119. [http://dx.doi.org/10.1016/S0300-483X\(03\)00206-3](http://dx.doi.org/10.1016/S0300-483X(03)00206-3).
- Green, T; Dow, J; Ong, C; Ng, V; Ong, H; Zhuang, Z; Yang, X; Bloemen, L. (2004). Biological monitoring of kidney function among workers occupationally exposed to trichloroethylene. *Occup Environ Med* 61: 312-317.
- Greenberg, M; Burton, G; Fisher, J. (1999). Physiologically based pharmacokinetic modeling of inhaled trichloroethylene and its oxidative metabolites in B6C3F1 mice. *Toxicol Appl Pharmacol* 154: 264-278. <http://dx.doi.org/10.1006/taap.1998.8594>.
- Greene, A; Wiener, S; Puder, M; Yoshida, A; Shi, B; Perez-Atayde, A; Efstathiou, J; Holmgren, L; Adamis, A; Rupnick, M; Folkman, J; O'Reilly, M. (2003). Endothelial-directed hepatic regeneration after partial hepatectomy. *Ann Surg* 237: 530-535. <http://dx.doi.org/10.1097/01.SLA.0000059986.96051.EA>.
- Greenland, S. (1992). A semi-Bayes approach to the analysis of correlated multiple associations, with an application to an occupational cancer-mortality study. *Stat Med* 11: 219-230. <http://dx.doi.org/10.1002/sim.4780110208>.
- Greenland, S; Salvan, A; Wegman, D; Hallock, M; Smith, T. (1994). A case-control study of cancer mortality at a transformer-assembly facility. *Int Arch Occup Environ Health* 66: 49-54. <http://dx.doi.org/10.1007/BF00386579>.
- Greenland, S. (2004). Model-based estimation of relative risks and other epidemiologic measures in studies of common outcomes and in case-control studies. *Am J Epidemiol* 160: 301-305. <http://dx.doi.org/10.1093/aje/kwh221>.
- Gregersen, P. (1988). Neurotoxic effects of organic solvents in exposed workers: Two controlled follow-up studies after 5.5 and 10.6 years. *Am J Ind Med* 14: 681-701.
- Greim, H; Bonse, G; Radwan, Z; Reichert, D; Henschler, D. (1975). Mutagenicity in vitro and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. *Biochem Pharmacol* 24: 2013-2017. [http://dx.doi.org/10.1016/0006-2952\(75\)90396-2](http://dx.doi.org/10.1016/0006-2952(75)90396-2).
- Greim, H; Wolff, T; Höfler, M; Lahaniatis, E. (1984). Formation of dichloroacetylene from trichloroethylene in the presence of alkaline material--possible cause of intoxication after abundant use of chloroethylene-containing solvents. *Arch Toxicol* 56: 74-77.
- Greten, F; Eckmann, L; Greten, T; Park, J; Li, Z-W; Egan, L; Kagnoff, M; Karin, M. (2004). IKKbeta links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* 118: 285-296. <http://dx.doi.org/10.1016/j.cell.2004.07.013>.
- Griffin, J; Blossom, S; Jackson, S; Gilbert, K; Pumford, N. (2000a). Trichloroethylene accelerates an autoimmune response by Th1 T cell activation in MRL +/+ mice. *Immunopharmacology* 46: 123-137. [http://dx.doi.org/10.1016/S0162-3109\(99\)00164-2](http://dx.doi.org/10.1016/S0162-3109(99)00164-2).
- Griffin, J; Gilbert, K; Lamps, L; Pumford, N. (2000b). CD4+ T-cell activation and induction of autoimmune hepatitis following trichloroethylene treatment in MRL+/+ mice. *Toxicol Sci* 57: 345-352.
- Griffin, J; Gilbert, K; Pumford, N. (2000c). Inhibition of CYP2E1 reverses CD4+ T-cell alterations in trichloroethylene-treated MRL+/+ mice. *Toxicol Sci* 54: 384-389.
- Grosse, Y; Baan, R; Secretan-lauby, B; El Ghissassi, F; Bouvard, V; Benbrahim-Tallaa, L; Guha, N; Islami, F; Galichet, L; Straif, K. (2011). Carcinogenicity of chemicals in industrial and consumer products, food contaminants and flavourings, and water chlorination byproducts. *Lancet Oncol* 12: 328-329. [http://dx.doi.org/10.1016/S1470-2045\(11\)70088-2](http://dx.doi.org/10.1016/S1470-2045(11)70088-2).
- Groupwater Vapor Project, Endicott, New York: Summary of findings, working draft. Cited in ATSDR. (2006). Endicott Area, Town of Union, Broome County, New York: Health Statistics Reviews, Cancer and Birth Outcome Analysis.
- Groves, C; Hayden, P; Lock, E; Schnellmann, R. (1993). Differential cellular effects in the toxicity of haloalkene and haloalkane cysteine conjugates to rabbit renal proximal tubules. *J Biochem Toxicol* 8: 49-56.

- Gu, Z; Sele, B; Chmara, D; Jalbert, P; Vincent, M; Vincent, F; Marka, C; Faure, J. (1981a). [Effects of trichlorethylene and its metabolites on the rate of sister chromatid exchange. In vivo and in vitro study on the human lymphocytes]. Ann Genet 24: 105-106.
- Gu, Z; Sele, B; Jalbert, P; Vincent, M; Vincent, F; Marka, C; Chmara, D; Faure, J. (1981b). [Induction of sister chromatid exchange by trichloroethylene and its metabolites]. Toxicol Eur Res 3: 63-67.
- Gualandi, G. (1987). Use of alpha- and beta-tubulin mutants for the study of spontaneous and induced chromosomal mis-distribution in Aspergillus nidulans. Mutat Res-Fundam Mol Mech Mutagen 178: 33-41. [http://dx.doi.org/10.1016/0027-5107\(87\)90083-2](http://dx.doi.org/10.1016/0027-5107(87)90083-2).
- Guehl, D; Bezard, E; Dovero, S; Boraud, T; Bioulac, B; Gross, C. (1999). Trichloroethylene and parkinsonism: A human and experimental observation. Eur J Neurol 6: 609-611. <http://dx.doi.org/10.1046/j.1468-1331.1999.650609.x>.
- Guengerich, F; Crawford, WJ; Watanabe, P. (1979). Activation of vinyl chloride to covalently bound metabolites: roles of 2-chloroethylene oxide and 2-chloroacetaldehyde. Biochemistry 18: 5177-5182.
- Guengerich, F; Kim, D; Iwasaki, M. (1991). Role of human cytochrome P-450 IIE1 in the oxidation of many low molecular weight cancer suspects. Chem Res Toxicol 4: 168-179. <http://dx.doi.org/10.1021/tx00020a008>.
- Guengerich, F and Shimada, T. (1991). Oxidation of toxic and carcinogenic chemicals by human cytochrome P-450 enzymes. Chem Res Toxicol 4: 391-407. <http://dx.doi.org/10.1021/tx00022a001>.
- Guengerich, F. (2004). Metabolism of trichlorethylene and covalent binding of reaction products. Proceedings of symposium on new scientific research related to the health effects of trichlorethylene. Washington, DC.
- Guerrero-Bosagna, C and Skinner, M. (2009). Epigenetic transgenerational effects of endocrine disruptors on male reproduction. Seminars in Reproductive Medicine 27: 403-408. <http://dx.doi.org/10.1055/s-0029-1237428>.
- Gun, R; Grygorcewicz, C; Nettelbeck, T. (1978). Choice reaction time in workers using trichloroethylene. Med J Aust 1: 535-536.
- Guo, H; Lee, S; Chan, L; Li, W. (2004). Risk assessment of exposure to volatile organic compounds in different indoor environments. Environ Res 94: 57-66.
- Gupta, S. (2000). Hepatic polyploidy and liver growth control. Semin Cancer Biol 10: 161-171. <http://dx.doi.org/10.1006/scbi.2000.0317>.
- Gustavsson, P; Jakobsson, R; Johansson, H; Lewin, F; Norell, S; Rutqvist, L. (1998). Occupational exposures and squamous cell carcinoma of the oral cavity, pharynx, larynx, and oesophagus: A case-control study in Sweden. Occup Environ Med 55: 393-400.
- Guyton, K; Barone, S; Brown, R; Euling, S; Jinot, J; Makris, S. (2008). Mode of action frameworks: a critical analysis. J Toxicol Environ Health B Crit Rev 11: 16-31. <http://dx.doi.org/10.1080/10937400701600321>.
- Guyton, K; Chiu, W; Bateson, T; Jinot, J; Scott, C; Brown, R; Caldwell, J. (2009). A reexamination of the PPAR-alpha activation mode of action as a basis for assessing human cancer risks of environmental contaminants. Environ Health Perspect 117: 1664-1672. <http://dx.doi.org/10.1289/ehp.0900758>.
- Guzelian, PS; Henry, CJ; Olin, SS (Eds.). (1992). Similarities and differences between children and adults: Implications for risk assessment. New York: ILSI Press.
- Gyorke, S and Terentyev, D. (2008). Modulation of ryanodine receptor by luminal calcium and accessory proteins in health and cardiac disease. Cardiovasc Res 77: 245-255. <http://dx.doi.org/10.1093/cvr/cvm038>.
- Habib, G; Barrios, R; Shi, Z; Lieberman, M. (1996). Four distinct membrane-bound dipeptidase RNAs are differentially expressed and show discordant regulation with gamma-glutamyl transpeptidase. J Biol Chem 271: 16273-16280.

- Hack, C; Chiu, W; Jay Zhao, Q; Clewell, H. (2006). Bayesian population analysis of a harmonized physiologically based pharmacokinetic model of trichloroethylene and its metabolites. *Regul Toxicol Pharmacol* 46: 63-83. <http://dx.doi.org/10.1016/j.yrtph.2006.05.012>.
- Haddad, S; Charest-Tardif, G; Krishnan, K. (2000). Physiologically based modeling of the maximal effect of metabolic interactions on the kinetics of components of complex chemical mixtures. *J Toxicol Environ Health A* 61: 209-223. <http://dx.doi.org/10.1080/00984100050131350>.
- Haglid, K; Briving, C; Hansson, H-A; Rosengren, L; Kjellstrand, P; Stavron, D; Swedin, U; Wronski, A. (1981). Trichloroethylene: Long-lasting changes in the brain after rehabilitation. *Neurotoxicology* 2: 659-673.
- Hakkola, J; Pasanen, M; Hukkanen, J; Pelkonen, O; Maenpaa, J; Edwards, R; Boobis, A; Raunio, H. (1996a). Expression of xenobiotic-metabolizing cytochrome P450 forms in human full-term placenta. *Biochem Pharmacol* 51: 403-411. [http://dx.doi.org/10.1016/0006-2952\(95\)02184-1](http://dx.doi.org/10.1016/0006-2952(95)02184-1).
- Hakkola, J; Raunio, H; Purkunen, R; Pelkonen, O; Saarikoski, S; Cresteil, T; Pasanen, M. (1996b). Detection of cytochrome P450 gene expression in human placenta in first trimester of pregnancy. *Biochem Pharmacol* 52: 379-383. [http://dx.doi.org/10.1016/0006-2952\(96\)00216-X](http://dx.doi.org/10.1016/0006-2952(96)00216-X).
- Hakkola, J; Pelkonen, O; Pasanen, M; Raunio, H. (1998a). Xenobiotic-metabolizing cytochrome P450 enzymes in the human feto-placental unit: Role in intrauterine toxicity. *Crit Rev Toxicol* 28: 35-72. <http://dx.doi.org/10.1080/10408449891344173>.
- Hakkola, J; Tanaka, E; Pelkonen, O. (1998b). Developmental expression of cytochrome P450 enzymes in human liver. *Pharmacol Toxicol* 82: 209-217. <http://dx.doi.org/10.1111/j.1600-0773.1998.tb01427.x>.
- Halmes, N; Perkins, E; McMillan, D; Pumford, N. (1997). Detection of trichloroethylene-protein adducts in rat liver and plasma. *Toxicol Lett* 92: 187-194.
- Hamada, T and Tanaka, H. (1995). Transfer of methyl chloroform, trichloroethylene and tetrachloroethylene to milk, tissues and expired air following intraruminal or oral administration in lactating goats and milk-fed kids. *Environ Pollut* 87: 313-318. [http://dx.doi.org/10.1016/0269-7491\(94\)P4163-I](http://dx.doi.org/10.1016/0269-7491(94)P4163-I).
- Hamdan, H and Stacey, N. (1993). Mechanism of trichloroethylene-induced elevation of individual serum bile acids: I. Correlation of trichloroethylene concentrations to bile acids in rat serum. *Toxicol Appl Pharmacol* 121: 291-295. <http://dx.doi.org/10.1006/taap.1993.1156>.
- Hanahan, D and Weinberg, R. (2000). The hallmarks of cancer. *Cell* 100: 57-70. [http://dx.doi.org/10.1016/S0092-8674\(00\)81683-9](http://dx.doi.org/10.1016/S0092-8674(00)81683-9).
- Hansch, C; Leo, A; Hoekman, D. (1995). Exploring QSAR: Hydrophobic, electronic, and steric constants. Washington, DC: American Chemical Society.
- Hansen, J; Raaschou-Nielsen, O; Christensen, J; Johansen, I; McLaughlin, J; Lipworth, L; Blot, W; Olsen, J. (2001). Cancer incidence among Danish workers exposed to trichloroethylene. *J Occup Environ Med* 43: 133-139.
- Hardell, L; Eriksson, M; Lenner, P; Lundgren, E. (1981). Malignant lymphoma and exposure to chemicals, especially organic solvents, chlorophenols and phenoxy acids: A case-control study. *Br J Cancer* 43: 169-176.
- Hardell, L; Bengtsson, N; Jonsson, U; Eriksson, S; Larsson, L. (1984). Aetiological aspects on primary liver cancer with special regard to alcohol, organic solvents and acute intermittent porphyria - An epidemiological investigation. *Br J Cancer* 50: 389-397.
- Hardell, L; Eriksson, M; Degerman, A. (1994). Exposure to phenoxyacetic acids, chlorophenols, or organic solvents in relation to histopathology, stage, and anatomical localization of non-Hodgkin's lymphoma. *Cancer Res* 54: 2386-2389.
- Hardin, B; Bond, G; Sikov, M; Andrew, F; Beliles, R; Niemeier, R. (1981). Testing of selected workplace chemicals for teratogenic potential. *Scand J Work Environ Health* 7: 66-75.
- Hardin, B; Kelman, B; Brent, R. (2004). Trichloroethylene and cardiac malformations. *Environ Health Perspect* 112: A607-608; author reply A608-609.

- Hardin, B; Kelman, B; Brent, R. (2005). Trichloroethylene and dichloroethylene: a critical review of teratogenicity. Birth Defects Res A Clin Mol Teratol 73: 931-955.
<http://dx.doi.org/10.1002/bdra.20192>.
- Harrington-Brock, K; Doerr, C; Moore, M. (1998). Mutagenicity of three disinfection by-products: Di- and trichloroacetic acid and chloral hydrate in L5178Y(+-) --3.7.2C mouse lymphoma cells. Mutat Res Genet Environ Mutagen 413: 265-276.
- Harrington, J; Whitby, H; Gray, C; Reid, F; Aw, T; Waterhouse, J. (1989). Renal disease and occupational exposure to organic solvents: A case referent approach. Br J Ind Med 46: 643-650.
<http://dx.doi.org/10.1136/oem.46.9.643>.
- Harris, N; Jaffe, E; Diebold, J; Flandrin, G; Muller-Hermelink, H; Vardiman, J. (2000). Lymphoma classification - from controversy to consensus: The R.E.A.L. and WHO Classification of lymphoid neoplasms. Ann Oncol 11: 3-10. http://dx.doi.org/10.1093/annonc/11.suppl_1.S3.
- Haseman, J; Huff, J; Boorman, G. (1984). Use of historical control data in carcinogenicity studies in rodents. Toxicol Pathol 12: 126-135. <http://dx.doi.org/10.1177/019262338401200203>.
- Hasmall, S and Roberts, R. (2000). The nongenotoxic hepatocarcinogens diethylhexylphthalate and methylclofenapate induce DNA synthesis preferentially in octoploid rat hepatocytes. Toxicol Pathol 28: 503-509. <http://dx.doi.org/10.1177/019262330002800401>.
- Hassall, C; Brendel, K; Gandolfi, A. (1983). Regulation of a S(trans-1,2-dichlorovinyl)-L-cysteine-induced renal tubular toxicity by glutathione. J Appl Toxicol 3: 321-325.
- Hassan, M; Frome, A; Patt, Y; El-Serag, H. (2002). Rising prevalence of hepatitis C virus infection among patients recently diagnosed with hepatocellular carcinoma in the United States. J Clin Gastroenterol 35: 266-269.
- Hattis, D; Ginsberg, G; Sonawane, B; Smolenski, S; Russ, A; Kozlak, M; Goble, R. (2003). Differences in pharmacokinetics between children and adults: II. Childrens variability in drug elimination half-lives and in some parameters needed for physiologically-based pharmacokinetic modeling. Risk Anal 23: 117-142. <http://dx.doi.org/10.1111/1539-6924.00295>.
- Hawley, G and Lewis, R, Sr (Eds.). (2001). Hawley's Condensed Chemical Dictionary (14 ed.). New York, NY: John Wiley & Sons, Inc.
- Haworth, S; Lawlor, T; Mortelmans, K; Speck, W; Zeiger, E. (1983). Salmonella mutagenicity test results for 250 chemicals. Environ Mutagen 5: 3-142. <http://dx.doi.org/10.1002/em.2860050703>.
- Hayden, M and Ghosh, S. (2004). Signaling to NF-kappaB. Genes Dev 18: 2195-2224.
<http://dx.doi.org/10.1101/gad.1228704>.
- Hays, T; Rusyn, I; Burns, A; Kennett, M; Ward, J; Gonzalez, F; Peters, J. (2005). Role of peroxisome proliferator-activated receptor-alpha (PPARalpha) in bezafibrate-induced hepatocarcinogenesis and cholestasis. Carcinogenesis 26: 219-227. <http://dx.doi.org/10.1093/carcin/bgh285>.
- Healy, T; Poole, T; Hopper, A. (1982). Rat fetal development and maternal exposure to trichloroethylene 100 ppm. Br J Anaesth 54: 337-341.
- Heath, C; Lally, C; Calle, E; McLaughlin, J; Thun, M. (1997). Hypertension, diuretics, and antihypertensive medications as possible risk factors for renal cell cancer. Am J Epidemiol 145: 607-613.
- Hegi, M; Fox, T; Belinsky, S; Devereux, T; Anderson, M. (1993). Analysis of activated protooncogenes in B6C3F1 mouse liver tumors induced by ciprofibrate, a potent peroxisome proliferator. Carcinogenesis 14: 145-149. <http://dx.doi.org/10.1093/carcin/14.1.145>.
- Hein, M; Waters, M; Ruder, A; Stenzel, M; Blair, A; Stewart, P. (2010). Statistical modeling of occupational chlorinated solvent exposures for case-control studies using a literature-based database. Ann Occup Hyg 54: 459-472. <http://dx.doi.org/10.1093/annhyg/meq027>.
- Heineman, E; Cocco, P; Gomez, M; Dosemeci, M; Stewart, P; Hayes, R; Zahm, S; Thomas, T; Blair, A. (1994). Occupational exposure to chlorinated aliphatic hydrocarbons and risk of astrocytic brain cancer. Am J Ind Med 26: 155-169. <http://dx.doi.org/10.1002/ajim.4700260203>.

- Heinemann, K; Willich, S; Heinemann, L; DoMinh, T; Möhner, M; Heuchert, G. (2000). Occupational exposure and liver cancer in women: results of the Multicentre International Liver Tumour Study (MILTS). *Occup Med (Lond)* 50: 422-429.
- Hejtmancik, M; Trela, B; Kurtz, P; Persing, R; Ryan, M; Yarrington, J; Chhabra, R. (2002). Comparative gavage subchronic toxicity studies of o-chloroaniline and m-chloroaniline in F344 rats and B6C3F1 mice. *Toxicol Sci* 69: 234-243.
- Helliwell, P and Hutton, A. (1950). Trichlorethylene anesthesia. *Anaesthesia* 5: 4-13.
<http://dx.doi.org/10.1111/j.1365-2044.1950.tb13680.x>.
- Henderson, G; Yan, Z; James, M; Davydova, N; Stacpoole, P. (1997). Kinetics and metabolism of chloral hydrate in children: Identification of dichloroacetate as a metabolite. *Biochem Biophys Res Commun* 235: 695-698. <http://dx.doi.org/10.1006/bbrc.1997.6868>.
- Handler, A and Crow, W. (1992). Paper No. 92-75.05.
- Hendriksen, P; Freidig, A; Jonker, D; Thissen, U; Bogaards, J; Mumtaz, M; Groten, J; Stierum, R. (2007). Transcriptomics analysis of interactive effects of benzene, trichloroethylene and methyl mercury within binary and ternary mixtures on the liver and kidney following subchronic exposure in the rat. *Toxicol Appl Pharmacol* 225: 171-188. <http://dx.doi.org/10.1016/j.taap.2007.08.017>.
- Hennig, U; Rudd, N; Hoar, D. (1988). Kinetochore immunofluorescence in micronuclei: A rapid method for the in situ detection of aneuploidy and chromosome breakage in human fibroblasts. *Mutat Res* 203: 405-414.
- Henning, S and Swendseid, M. (1996). The role of folate, choline, and methionine in carcinogenesis induced by methyl-deficient diets. *Adv Exp Med Biol* 399: 143-155.
- Henschler, D; Eder, E; Neudecker, T; Metzler, M. (1977). Carcinogenicity of trichloroethylene: fact or artifact? *Arch Toxicol* 37: 233-236.
- Henschler, D; Romen, W; Elsaesser, H; Reichert, D; Eder, E; Radwan, Z. (1980). Carcinogenicity study of trichloroethylene by longterm inhalation in three animal species. *Arch Toxicol* 43: 237-248. <http://dx.doi.org/10.1007/BF00366179>.
- Henschler, D; Elsaesser, H; Romen, W; Eder, E. (1984). Carcinogenicity study of trichloroethylene, with and without epoxide stabilizers, in mice. *J Cancer Res Clin Oncol* 107: 149-156. <http://dx.doi.org/10.1007/BF01032599>.
- Henschler, D; Vamvakas, S; Lammert, M; Dekant, W; Kraus, B; Thomas, B; Ulm, K. (1995). Increased incidence of renal cell tumors in a cohort of cardboard workers exposed to trichloroethene. *Arch Toxicol* 69: 291-299. <http://dx.doi.org/10.1007/s002040050173>.
- Herbert, V; Gardner, A; Colman, N. (1980). Mutagenicity of dichloroacetate, an ingredient of some formulations of pangamic acid (trade-named "vitamin B15"). *Am J Clin Nutr* 33: 1179-1182.
- Herman, J; Umar, A; Polyak, K; Graff, J; Ahuja, N; Issa, J; Markowitz, S; Willson, J; Hamilton, S; Kinzler, K; Kane, M; Kolodner, R; Vogelstein, B; Kunkel, T; Baylin, S. (1998). Incidence and functional consequences of hMLH1 promoter hypermethylation in colorectal carcinoma. *PNAS* 95: 6870-6875.
- Hernberg, S; Korkala, M; Asikainen, U; Riala, R. (1984). Primary liver cancer and exposure to solvents. *Int Arch Occup Environ Health* 54: 147-153. <http://dx.doi.org/10.1007/BF00378517>.
- Hernberg, S; Kauppinen, T; Riala, R; Korkala, M; Asikainen, U. (1988). Increased risk for primary liver cancer among women exposed to solvents. *Scand J Work Environ Health* 14: 356-365.
- Herren-Freund, S; Pereira, M; Khouri, M; Olson, G. (1987). The carcinogenicity of trichloroethylene and its metabolites, trichloroacetic acid and dichloroacetic acid, in mouse liver. *Toxicol Appl Pharmacol* 90: 183-189. [http://dx.doi.org/10.1016/0041-008X\(87\)90325-5](http://dx.doi.org/10.1016/0041-008X(87)90325-5).
- Herrera, E; Samper, E; Martín-Caballero, J; Flores, J; Lee, H; Blasco, M. (1999). Disease states associated with telomerase deficiency appear earlier in mice with short telomeres. *EMBO J* 18: 2950-2960. <http://dx.doi.org/10.1093/emboj/18.11.2950>.
- Hers, I; Zapf-Gilje, R; Li, L; Atwater, J. (2001). The use of indoor air measurements to evaluate intrusion of subsurface VOC vapors into buildings. *J Air Waste Manag Assoc* 51: 1318-1331.

- Higgins, J; Thompson, S; Deeks, J; Altman, D. (2003). Measuring inconsistency in meta-analyses. *BMJ* 327: 557-560. <http://dx.doi.org/10.1136/bmj.327.7414.557>.
- Hill, A. (1965). The environment and disease: Association or causation? *Proc R Soc Med* 58: 295-300.
- Hinchman, C and Ballatori, N. (1990). Glutathione-degrading capacities of liver and kidney in different species. *Biochem Pharmacol* 40: 1131-1135. [http://dx.doi.org/10.1016/0006-2952\(90\)90503-D](http://dx.doi.org/10.1016/0006-2952(90)90503-D).
- Hines, R and McCarver, D. (2002). The ontogeny of human drug-metabolizing enzymes: Phase I oxidative enzymes. *J Pharmacol Exp Ther* 300: 355-360. <http://dx.doi.org/10.1124/jpet.300.2.355>.
- Hirsch, A; Rankin, K; Panelli, P. (1996). Trichloroethylene exposure and headache. *Headache Q* 7: 126-138.
- Hissink, E; Bogaards, J; Freidig, A; Commandeur, J; Vermeulen, N; van Bladeren, P. (2002). The use of in vitro metabolic parameters and physiologically based pharmacokinetic (PBPK) modeling to explore the risk assessment of trichloroethylene. *Environ Toxicol Pharmacol* 11: 259-271. [http://dx.doi.org/10.1016/S1382-6689\(02\)00019-4](http://dx.doi.org/10.1016/S1382-6689(02)00019-4).
- Hobara, T; Kobayashi, H; Higashihara, E; Kawamoto, T; Sakai, T. (1984). Acute effects of 1,1,1-trichloroethane, trichloroethylene, and toluene on the hematologic parameters in dogs. *Arch Environ Contam Toxicol* 13: 589-593.
- Hodgen, G and Sherins, R. (1973). Enzymes as markers of testicular growth and development in the rat. *Endocrinology* 93: 985-989.
- Hoel, D and Walburg, H, Jr. (1972). Statistical analysis of survival experiments. *J Natl Cancer Inst* 49: 361-372.
- Hoffman, S; Mishima, N; Krug, E. (2004). An improved model for evaluating trichloroethylene and its metabolites as cardiac specific teratogens. In D Jollow (Ed.), *Trichloroethylene: the scientific basis of risk assessment*: The Medical University of South Carolina Press.
- Hoivik, D; Qualls, C, Jr; Mirabile, R; Cariello, N; Kimbrough, C; Colton, H; Anderson, S; Santostefano, M; Morgan, R; Dahl, R; Brown, A; Zhao, Z; Mudd, P, Jr; Oliver, W, Jr; Brown, H; Miller, R. (2004). Fibrates induce hepatic peroxisome and mitochondrial proliferation without overt evidence of cellular proliferation and oxidative stress in cynomolgus monkeys. *Carcinogenesis* 25: 1757-1769. <http://dx.doi.org/10.1093/carcin/bgh182>.
- Hooper, N; Hesp, R; Tieku, S. (1994). Metabolism of aspartame by human and pig intestinal microvillar peptidases. *Biochem J* 298: 635-639.
- Hoover, R and Strasser, P. (1980). Artificial sweeteners and human bladder cancer: Preliminary results. *Lancet* 315: 837-840. [http://dx.doi.org/10.1016/S0140-6736\(80\)91350-1](http://dx.doi.org/10.1016/S0140-6736(80)91350-1).
- Horesovsky, G; Ginsler, J; Everitt, J; Walker, C. (1994). Increased susceptibility to in vitro transformation of cells carrying the Eker tumor susceptibility mutation. *Carcinogenesis* 15: 2183-2187.
- Hornung, R; Pinney, S; Lodwick, J; Killough, G; Brewer, D; Nasuta, J. (2008). Estimation of radon exposures to workers at the Fernald Feed Materials Production Center 1952-1988. *J Expo Sci Environ Epidemiol* 18: 512-523. <http://dx.doi.org/10.1038/sj.jes.7500645>.
- Hortelano, S; Dewez, B; Genaro, A; Díaz-Guerra, M; Boscá, L. (1995). Nitric oxide is released in regenerating liver after partial hepatectomy. *Hepatology* 21: 776-786. <http://dx.doi.org/10.1002/hep.1840210327>.
- Horvath, A; Getzen, F; Maczynska, Z. (1999). IUPAC-NIST Solubility data series 67: Halogenated ethanes and ethenes with water. *J Phys Chem Ref Data* 28: 395-627. <http://dx.doi.org/10.1063/1.556039>.
- Howard, P; Boethling, R; Jarvis, W; Meylan, W; Michalenko, E. (1991). *Handbook of environmental degradation rates*. Boca Raton, FL: Lewis Publishers/CRC Press.
- Howroyd, P; Swanson, C; Dunn, C; Cattley, R; Corton, J. (2004). Decreased longevity and enhancement of age-dependent lesions in mice lacking the nuclear receptor peroxisome proliferator-activated receptor alpha (PPARalpha). *Toxicol Pathol* 32: 591-599.

- Hrelia, P; Maffei, F; Vigagni, F; Fimognari, C; Flori, P; Stanzani, R; Forti, G. (1994). Interactive effects between trichloroethylene and pesticides at metabolic and genetic level in mice. Environ Health Perspect 102: 31-34.
- HSDB. (Hazardous Substances Data Bank). (1996). Trichloroethylene. Bethesda, MD: National Library of Medicine. Retrieved from <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>
- HSDB. (Hazardous Substances Data Bank). (2002). Trichloroethylene. Bethesda, MD: National Library of Medicine. Retrieved from <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>
- Hsia, C; Huo, T; Chiang, S; Lu, M-F; Sun, C; Wu, J; Lee, P; Chi, C; Lui, W; Lee, S. (2007). Evaluation of interleukin-6, interleukin-10 and human hepatocyte growth factor as tumor markers for hepatocellular carcinoma. Eur J Surg Oncol 33: 208-212.
<http://dx.doi.org/10.1016/j.ejso.2006.10.036>.
- Hu, C; Jiang, L; Geng, C; Zhang, X; Cao, J; Zhong, L. (2008). Possible involvement of oxidative stress in trichloroethylene-induced genotoxicity in human HepG2 cells. Mutat Res 652: 88-94.
<http://dx.doi.org/10.1016/j.mrgentox.2008.01.002>.
- Hu, Y; Hakkola, J; Oscarson, M; Ingelman-Sundberg, M. (1999). Structural and functional characterization of the 5'-flanking region of the rat and human cytochrome P450 2E1 genes: Identification of a polymorphic repeat in the human gene. Biochem Biophys Res Commun 263: 286-293. <http://dx.doi.org/10.1006/bbrc.1999.1362>.
- Huang, H; Li, L; Chen, B; Huang, J; Kuang, S. (2002). New problems caused by occupational trichloroethylene exposure. Int J Immunopathol Pharmacol 15: 30-32.
- Hunter, E; Rogers, E; Schmid, J; Richard, A. (1996). Comparative effects of haloacetic acids in whole embryo culture. Teratology 54: 57-64. [http://dx.doi.org/10.1002/\(SICI\)1096-9926\(199606\)54:2<57::AID-TERA1>3.0.CO;2-1](http://dx.doi.org/10.1002/(SICI)1096-9926(199606)54:2<57::AID-TERA1>3.0.CO;2-1).
- IARC. (International Agency for Research on Cancer). (1995a). Dry cleaning, some chlorinated solvents and other industrial chemicals: Summary of data reported and evaluation. In IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Lyon, France.
- IARC. (International Agency for Research on Cancer). (1995b). Dry cleaning, some chlorinated solvents and other industrial chemicals. In IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Lyon, France. <http://monographs.iarc.fr/ENG/Monographs/vol63/index.php>.
- IARC. (International Agency for Research on Cancer). (1995c). Tetrachloroethylene. In IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, pp. 159-221. Lyon, France. <http://monographs.iarc.fr/ENG/Monographs/vol63/index.php>.
- IARC. (International Agency for Research on Cancer). (2000). Di(2-ethylhexyl) phthalate. In: IARC monographs on the evaluation of carcinogenic risk of chemicals to humans Vol 77 Some industrial chemicals. Lyon, France: International Agency for Research on Cancer. <http://monographs.iarc.fr/ENG/Monographs/vol77/mono77.pdf>.
- IARC. (International Agency for Research on Cancer). (2001). Pathology and genetics of tumours of haematopoietic and lymphoid tissues. In IARC WHO classification of tumours, pp. 351. Lyon, France.
<http://apps.who.int/bookorders/anglais/detart1.jsp?sesslan=1&codlan=1&codcol=70&codcch=3>.
- IARC. (International Agency for Research on Cancer). (2004a). Tobacco smoke and involuntary smoking. pp. 1473. Lyon, France.
- IARC. (International Agency for Research on Cancer). (2004b). Some drinking-water disinfectants and contaminants, including arsenic. In IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Lyon, France. <http://monographs.iarc.fr/ENG/Monographs/vol84/index.php>.
- Iavicoli, I; Marinaccio, A; Carelli, G. (2005). Effects of occupational trichloroethylene exposure on cytokine levels in workers. J Occup Environ Med 47: 453-457.
- ICRP. (International Commission on Radiological Protection). (2003). Basic anatomical and physiological data for use in radiological protection: Reference values. (ICRP Publication 89). In Annals of the ICRP. New York, NY: Pergamon Press.
http://www.elsevier.com/wps/find/bookdescription.cws_home/672826/description#description.

- IEL (International Epidemiology Institute). (2005). Rocketdyne worker health study: Final report: Executive summary & appendices. pp. 189. Rockville, MD.
http://www.boeing.com/aboutus/environment/santa_susana/factsheets/Rocketdyne_Worker_Health_h_Study_Final_Report_July_2005.pdf
- Ikbal, M; Tastekin, A; Dogan, H; Pirim, I; Ors, R. (2004). The assessment of genotoxic effects in lymphocyte cultures of infants treated with chloral hydrate. Mutat Res 564: 159-164.
<http://dx.doi.org/10.1016/j.mrgentox.2004.08.007>.
- Ikeda, M; Otsuji, H; Kawai, H; Kuniyoshi, M. (1971). Excretion kinetics of urinary metabolites in a patient addicted to trichloroethylene. Br J Ind Med 28: 203-206.
- Ikeda, M; Ohtsuji, H; Imamura, T; Komoike, Y. (1972). Urinary excretion of total trichloro-compounds, trichloroethanol, and trichloroacetic acid as a measure of exposure to trichloroethylene and tetrachloroethylene. Br J Ind Med 29: 328-333. <http://dx.doi.org/10.1136/oem.29.3.328>.
- Ikeda, M and Imamura, T. (1973). Biological half-life of trichloroethylene and tetrachloroethylene in human subjects. Int Arch Occup Environ Health 31: 209-224.
<http://dx.doi.org/10.1007/BF00539241>.
- Ikeda, M. (1977). Metabolism of trichloroethylene and tetrachloroethylene in human subjects. Environ Health Perspect 21: 239-245. <http://dx.doi.org/10.1289/ehp.7721239>.
- Inoue, O; Seiji, K; Kawai, T; Jin, C; Liu, Y-T; Chen, Z; Cai, S-X; Yin, S-N; Li, G-L; Nakasutka, H; Watanabe, T; Ikeda, M. (1989). Relationship between vapor exposure and urinary metabolite excretion among workers exposed to trichloroethylene. Am J Ind Med 15: 103-110.
- IOM. (Institute of Medicine). (2003). Gulf war and health: Volume 2. Insecticides and solvents. pp. 616. Washington, DC: National Academies Press. http://www.nap.edu/catalog.php?record_id=10628.
- Isaacs, K; Evans, M; Harris, T. (2004). Visualization-based analysis for a mixed-inhibition binary PBPK model: Determination of inhibition mechanism. J Pharmacokinet Pharmacodyn 31: 215-242.
- Isaacson, L and Taylor, D. (1989). Maternal exposure to 1,1,2-trichloroethylene affects myelin in the hippocampal formation of the developing rat. Brain Res 488: 403-407.
[http://dx.doi.org/10.1016/0006-8993\(89\)90739-7](http://dx.doi.org/10.1016/0006-8993(89)90739-7).
- Isaacson, L; Spohler, S; Taylor, D. (1990). Trichloroethylene affects learning and decreases myelin in the rat hippocampus. Neurotoxicol Teratol 12: 375-381. [http://dx.doi.org/10.1016/0892-0362\(90\)90057-J](http://dx.doi.org/10.1016/0892-0362(90)90057-J).
- Isacson, P; Bean, J; Splinter, R; Olson, D; Kohler, J. (1985). Drinking water and cancer incidence in Iowa. III. Association of cancer with indices of contamination. Am J Epidemiol 121: 856-869.
- Ito, Y; Yokota, H; Wang, R; Yamanoshita, O; Ichihara, G; Wang, H; Kurata, Y; Takagi, K; Nakajima, T. (2005). Species differences in the metabolism of di(2-ethylhexyl) phthalate (DEHP) in several organs of mice, rats, and marmosets. Arch Toxicol 79: 147-154.
<http://dx.doi.org/10.1007/s00204-004-0615-7>.
- Ito, Y; Yamanoshita, O; Asaeda, N; Tagawa, Y; Lee, C; Aoyama, T; Ichihara, G; Furuhashi, K; Kamijima, M; Gonzalez, F; Nakajima, T. (2007). Di(2-ethylhexyl)phthalate induces hepatic tumorigenesis through a peroxisome proliferator-activated receptor alpha-independent pathway. J Occup Health 49: 172-182.
- Iyer, R; Frink, E, Jr; Ebert, T; Anders, M. (1998). Cysteine conjugate beta-lyase-dependent metabolism of compound A (2-[fluoromethoxy]-1,1,3,3,3-pentafluoro-1-propene) in human subjects anesthetized with sevoflurane and in rats given compound A. Anesthesiology 88: 611-618.
- Jacob, S; Hery, M; Protois, J; Rossert, J; Stengel, B. (2007). New insight into solvent-related end-stage renal disease: Occupations, products and types of solvents at risk. Occup Environ Med 64: 843-848. <http://dx.doi.org/10.1136/oem.2007.032482>.
- Jaffe, D; Gandolfi, A; Nagle, R. (1984). Chronic toxicity of S-(trans-1,2-dichlorovinyl)-L-cysteine in mice. J Appl Toxicol 4: 315-319.
- Jaffe, D; Hassall, C; Gandolfi, A; Brendel, K. (1985). Production of DNA single strand breaks in rabbit renal tissue after exposure to 1,2-dichlorovinylcysteine. Toxicology 35: 25-33.

- Jakobson, I; Wahlberg, J; Holmberg, B; Johansson, G. (1982). Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. *Toxicol Appl Pharmacol* 63: 181-187. [http://dx.doi.org/10.1016/0041-008X\(82\)90039-4](http://dx.doi.org/10.1016/0041-008X(82)90039-4).
- Jakobson, I; Holmberg, B; Ekner, A. (1986). Venous blood levels of inhaled trichloroethylene in female rats and changes induced by interacting agents. *Acta Pharmacol Toxicol* 59: 135-143. <http://dx.doi.org/10.1111/j.1600-0773.1986.tb00145.x>.
- Jakubowski, M and Wieczorek, H. (1988). The effects of physical effort on pulmonary uptake of selected organic compound vapours. *Pol J Occup Med* 1: 62-71.
- James, M; Cornett, R; Yan, Z; Henderson, G; Stacpoole, P. (1997). Glutathione-dependent conversion to glyoxylate, a major pathway of dichloroacetate biotransformation in hepatic cytosol from humans and rats, is reduced in dichloroacetate-treated rats. *Drug Metab Dispos* 25: 1223-1227.
- Jansson, C; Johansson, A; Bergdahl, I; Dickman, P; Plato, N; Adami, J; Boffetta, P; Lagergren, J. (2005). Occupational exposures and risk of esophageal and gastric cardia cancers among male Swedish construction workers. *Cancer Causes Control* 16: 755-764. <http://dx.doi.org/10.1007/s10552-005-1723-2>.
- Jansson, C; Plato, N; Johansson, A; Nyrén, O; Lagergren, J. (2006). Airborne occupational exposures and risk of oesophageal and cardia adenocarcinoma. *Occup Environ Med* 63: 107-112. <http://dx.doi.org/10.1136/oem.2005.022467>.
- Jasińska, M. (1965). [General anesthesia with trichloroethylene in children with congenital developmental defects of the jaw and face]. *Czasopismo Stomatologiczne* 18: 935-940.
- Jaspers, R; Muijsen, H; Lammers, J; Kulig, B. (1993). Mid-frequency hearing loss and reduction of acoustic startle responding in rats following trichloroethylene exposure. *Neurotoxicol Teratol* 15: 407-412. [http://dx.doi.org/10.1016/0892-0362\(93\)90058-V](http://dx.doi.org/10.1016/0892-0362(93)90058-V).
- Jemal, A; Siegel, R; Ward, E; Hao, Y; Xu, J; Murray, T; Thun, M. (2008). Cancer statistics, 2008. *CA Cancer J Clin* 58: 71-96. <http://dx.doi.org/10.3322/CA.2007.0010>.
- Jensen, O; Knudsen, J; McLaughlin, J; Sorensen, B. (1988). The Copenhagen case-control study of renal pelvis and ureter cancer: role of smoking and occupational exposures. *Int J Cancer* 41: 557-561.
- Ji, B; Blair, A; Shu, X; Chow, W; Hauptmann, M; Dosemeci, M; Yang, G; Lubin, J; Gao, Y; Rothman, N; Zheng, W. (2008). Occupation and breast cancer risk among Shanghai women in a population-based cohort study. *Am J Ind Med* 51: 100-110. <http://dx.doi.org/10.1002/ajim.20507>.
- Ji, J and Hemminki, K. (2005). Variation in the risk for liver and gallbladder cancers in socioeconomic and occupational groups in Sweden with etiological implications. *Int Arch Occup Environ Health* 78: 641-649. <http://dx.doi.org/10.1007/s00420-005-0015-1>.
- Jia, H-L; Ye, Q-H; Qin, L-X; Budhu, A; Forgue, M; Chen, Y; Liu, Y-K; Sun, H-C; Wang, L; Lu, H-Z; Shen, F; Tang, Z-Y; Wang, X. (2007). Gene expression profiling reveals potential biomarkers of human hepatocellular carcinoma. *Clin Cancer Res* 13: 1133-1139. <http://dx.doi.org/10.1158/1078-0432.CCR-06-1025>.
- Jia, M; Coats, B; Chadha, M; Frentzen, B; Perez-Rodriguez, J; Chadik, P; Yost, R; Henderson, G; Stacpoole, P. (2006). Human kinetics of orally and intravenously administered low-dose 1,2-(13)C-dichloroacetate. *J Clin Pharmacol* 46: 1449-1459. <http://dx.doi.org/10.1177/0091270006292627>.
- Jirtle, R; Meyer, S; Brockenbrough, J. (1991). Liver tumor promoter phenobarbital: A biphasic modulator of hepatocyte proliferation. *Prog Clin Biol Res* 369: 209-216.
- Johnson, A; VanDemark, W; Gomes, W. (1967). The effect of antispermatic and hormone treatments on testicular cholesterol. *Fed Proc* 26: 645.
- Johnson, P. (2009). personal communication from Paula Johnson, University of Arizona, to Susan Makris, U.S. EPA, 26 August 2009
- Johnson, P; Dawson, B; Goldberg, S. (1998a). Cardiac teratogenicity of trichloroethylene metabolites. *J Am Coll Cardiol* 32: 540-545. [http://dx.doi.org/10.1016/S0735-1097\(98\)00232-0](http://dx.doi.org/10.1016/S0735-1097(98)00232-0).
- Johnson, P; Dawson, B; Goldberg, S. (1998b). A review: trichloroethylene metabolites: potential cardiac teratogens. *Environ Health Perspect* 106 Suppl 4: 995-999.

- Johnson, P; Goldberg, S; Mays, M; Dawson, B. (2003). Threshold of trichloroethylene contamination in maternal drinking waters affecting fetal heart development in the rat. Environ Health Perspect 111: 289-292.
- Johnson, P; Dawson, B; Goldberg, S; Mays, M. (2004). Trichloroethylene: Johnson et al.'s Response. Environ Health Perspect 112: a608-a609. <http://dx.doi.org/10.1289/ehp.112-a608>.
- Johnson, P; Goldberg, S; Mays, M; Dawson, B. (2005). Correction: Threshold of trichloroethylene contamination in maternal drinking waters affecting fetal heart development in the rat. Environ Health Perspect 113: A18.
- Johnsrud, E; Koukouritaki, S; Divakaran, K; Brunengraber, L; Hines, R; McCarver, D. (2003). Human hepatic CYP2E1 expression during development. J Pharmacol Exp Ther 307: 402-407. <http://dx.doi.org/10.1124/jpet.102.053124>.
- Jones, T; Qin, C; Schaeffer, V; Stevens, J. (1988). Immunohistochemical localization of glutamine transaminase K, a rat kidney cysteine conjugate beta-lyase, and the relationship to the segment specificity of cysteine conjugate nephrotoxicity. Mol Pharmacol 34: 621-627.
- Jorquera, R and Tanguay, R. (2001). Fumarylacetoacetate, the metabolite accumulating in hereditary tyrosinemia, activates the ERK pathway and induces mitotic abnormalities and genomic instability. Hum Mol Genet 10: 1741-1752.
- Juntunen, J; Hupli, V; Hernberg, S; Luisto, M. (1980). Neurological picture of organic solvent poisoning in industry: A retrospective clinical study of 37 patients. Int Arch Occup Environ Health 46: 219-231.
- Juntunen, J; Antti-Poika, M; Tola, S; Partanen, T. (1982). Clinical prognosis of patients with diagnosed chronic solvent intoxication. Acta Neurol Scand 65: 488-503. <http://dx.doi.org/10.1111/j.1600-0404.1982.tb03105.x>.
- Juran, B and Lazaridis, K. (2006). Genomics and complex liver disease: Challenges and opportunities. Hepatology 44: 1380-1390. <http://dx.doi.org/10.1002/hep.21453>.
- Kadry, A; Turkall, R; Skowronski, G; Abdel-Rahman, M. (1991). Soil adsorption alters kinetics and bioavailability of trichloroethylene in orally exposed female rats. Toxicol Lett 58: 337-346. [http://dx.doi.org/10.1016/0378-4274\(91\)90046-9](http://dx.doi.org/10.1016/0378-4274(91)90046-9).
- Käfer, E. (1986). Tests which distinguish induced crossing-over and aneuploidy from secondary segregation in Aspergillus treated with chloral hydrate or gamma-rays. Mutat Res 164: 145-166. [http://dx.doi.org/10.1016/0165-1161\(86\)90006-3](http://dx.doi.org/10.1016/0165-1161(86)90006-3).
- Kamijima, M; Hisanaga, N; Wang, H; Nakajima, T. (2007). Occupational trichloroethylene exposure as a cause of idiosyncratic generalized skin disorders and accompanying hepatitis similar to drug hypersensitivities. Int Arch Occup Environ Health 80: 357-370. <http://dx.doi.org/10.1007/s00420-006-0147-y>.
- Kamijima, M; Wang, H; Huang, H; Li, L; Shibata, E; Lin, B; Sakai, K; Liu, H; Tsuchiyama, F; Chen, J; Okamura, A; Huang, X; Hisanaga, N; Huang, Z; Ito, Y; Takeuchi, Y; Nakajima, T. (2008). Trichloroethylene causes generalized hypersensitivity skin disorders complicated by hepatitis. J Occup Health 50: 328-338.
- Kamijo, Y; Hora, K; Nakajima, T; Kono, K; Takahashi, K; Ito, Y; Higuchi, M; Kiyosawa, K; Shigematsu, H; Gonzalez, F; Aoyama, T. (2007). Peroxisome proliferator-activated receptor alpha protects against glomerulonephritis induced by long-term exposure to the plasticizer di-(2-ethylhexyl)phthalate. J Am Soc Nephrol 18: 176-188. <http://dx.doi.org/10.1681/asn.2006060597>.
- Kan, F; Forkert, P; Wade, M. (2007). Trichloroethylene exposure elicits damage in epididymal epithelium and spermatozoa in mice. Histol Histopathol 22: 977-988.
- Kaneko, T; Wang, P-Y; Sato, A. (1994). Enzymes induced by ethanol differently affect the pharmacokinetics of trichloroethylene and 1,1,1-trichloroethane. Occup Environ Med 51: 113-119.
- Kaneko, T; Saegusa, M; Tasaka, K; Sato, A. (2000). Immunotoxicity of trichloroethylene: A study with MRL-lpr/lpr mice. J Appl Toxicol 20: 471-475. [http://dx.doi.org/10.1002/1099-1263\(200011/12\)20:6<471::AID-JAT716>3.0.CO;2-E](http://dx.doi.org/10.1002/1099-1263(200011/12)20:6<471::AID-JAT716>3.0.CO;2-E).

- Kanisawa, M and Suzuki, S. (1978). Induction of renal and hepatic tumors in mice by ochratoxin A, a mycotoxin. *Gann* 69: 599-600.
- Kappas, A. (1989). On the mechanisms of induced aneuploidy in *Aspergillus nidulans* and validation of tests for genomic mutations. In MA Resnick & BK Vig (Eds.), *Mechanisms of Chromosome Distribution and Aneuploidy* (Vol. 318, pp. 377-384). New York, NY: Wiley.
- Karami, S; Boffetta, P; Brennan, P; Stewart, P; Zaridze, D; Matveev, V; Janout, V; Kollarova, H; Bencko, V; Navratilova, M; Szczesnia-Dabrowska, N; Mates, D; Gromiec, J; Sobotka, R; Chow, W; Rothman, N; Moore, L. (2011). Renal cancer risk and occupational exposure to polycyclic aromatic hydrocarbons and plastics. *J Occup Environ Med* 53: 218-223. <http://dx.doi.org/10.1097/JOM.0b013e31820a40a3>.
- Kargalioglu, Y; McMillan, B; Minear, R; Plewa, M. (2002). Analysis of the cytotoxicity and mutagenicity of drinking water disinfection by-products in *Salmonella typhimurium*. *Teratog Carcinog Mutagen* 22: 113-128. <http://dx.doi.org/10.1002/tcm.10010>.
- Karpinets, T and Foy, B. (2005). Tumorigenesis: The adaptation of mammalian cells to sustained stress environment by epigenetic alterations and succeeding matched mutations. *Carcinogenesis* 26: 1323-1334. <http://dx.doi.org/10.1093/carcin/bgi079>.
- Karunananayake, C; McDuffie, H; Dosman, J; Spinelli, J; Pahwa, P. (2008). Occupational exposures and non-Hodgkin's lymphoma: Canadian case-control study. *Environ Health* 7: 44. <http://dx.doi.org/10.1186/1476-069X-7-44>.
- Kato-Weinstein, J; Lingohr, M; Orner, G; Thrall, B; Bull, R. (1998). Effects of dichloroacetate on glycogen metabolism in B6C3F1 mice. *Toxicology* 130: 141-154. [http://dx.doi.org/10.1016/S0300-483X\(98\)00106-1](http://dx.doi.org/10.1016/S0300-483X(98)00106-1).
- Kato-Weinstein, J; Stauber, A; Orner, G; Thrall, B; Bull, R. (2001). Differential effects of dihalogenated and trihalogenated acetates in the liver of B6C3F1 mice. *J Appl Toxicol* 21: 81-89. <http://dx.doi.org/10.1002/jat.717>.
- Kauffmann, B; White, K; Sanders, V; Douglas, K; Sain, L; Borzelleca, J; Munson, A. (1982). Humoral and cell-mediated immune status in mice exposed to chloral hydrate. *Environ Health Perspect* 44: 147-151.
- Kautiainen, A; Vogel, J; Turteltaub, K. (1997). Dose-dependent binding of trichloroethylene to hepatic DNA and protein at low doses in mice. *Chem Biol Interact* 106: 109-121.
- Kawamoto, T; Hobara, T; Aoki, T; Kobayashi, H; Iwamoto, S; Sakai, T; Takano, T; Miyazaki, Y; Imamura, A; Ogino, K. (1988a). Metabolism of chloral hydrate in the anoxic perfused liver. *Toxicol Lett* 40: 225-231. [http://dx.doi.org/10.1016/0378-4274\(88\)90045-8](http://dx.doi.org/10.1016/0378-4274(88)90045-8).
- Kawamoto, T; Hobara, T; Nakamura, K; Imamura, A; Ogino, K; Kobayashi, H; Iwamoto, S; Sakai, T. (1988b). Induction of cytochrome P-450, cytochrome b-5, NADPH-cytochrome c reductase and change of cytochrome P-450 isozymes with long-term trichloroethylene treatment. *Toxicology* 53: 239-249. [http://dx.doi.org/10.1016/0300-483X\(88\)90217-X](http://dx.doi.org/10.1016/0300-483X(88)90217-X).
- Keech, A; Simes, R; Barter, P; Best, J; Scott, R; Taskinen, M; Forder, P; Pillai, A; Davis, T; Glasziou, P; Drury, P; Kesäniemi, Y; Sullivan, D; Hunt, D; Colman, P; d'Emden, M; Whiting, M; Ehnholm, C; Laakso, M. (2005). Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): Randomised controlled trial. *Lancet* 366: 1849-1861. [http://dx.doi.org/10.1016/S0140-6736\(05\)67667-2](http://dx.doi.org/10.1016/S0140-6736(05)67667-2).
- Keech, A; Simes, J; Barter, P; Best, J; Scott, R; Taskinen, M-R. (2006). Correction to the FIELD study report. *Lancet* 368: 1415. [http://dx.doi.org/10.1016/S0140-6736\(06\)69594-9](http://dx.doi.org/10.1016/S0140-6736(06)69594-9).
- Keil, D; Peden-Adams, M; Wallace, S; Ruiz, P; Gilkeson, G. (2009). Assessment of trichloroethylene (TCE) exposure in murine strains genetically-prone and non-prone to develop autoimmune disease. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 44: 443-453. <http://dx.doi.org/10.1080/10934520902719738>.
- Keller, D and Hd'A, H. (1988). Mechanistic studies on chloral toxicity: Relationship to trichloroethylene carcinogenesis. *Toxicol Lett* 42: 183-191. [http://dx.doi.org/10.1016/0378-4274\(88\)90076-8](http://dx.doi.org/10.1016/0378-4274(88)90076-8).

- Kelsh, M; Weingart, M; Mandel, J; Mink, P; Alexander, D; Basu, R; Goodman, M. (2005). A meta-analysis of occupational trichloroethylene exposure and selected cancers. Ann Epidemiol 15: 632-632. <http://dx.doi.org/10.1016/j.annepidem.2005.06.008>.
- Kelsh, M; Alexander, D; Mink, P; Mandel, J. (2010). Occupational trichloroethylene exposure and kidney cancer: a meta-analysis. Epidemiology 21: 95-102. <http://dx.doi.org/10.1097/EDE.0b013e3181c30e92>.
- Kenck, C; Wilhelm, M; Bugert, P; Staehler, G; Kovacs, G. (1996). Mutation of the VHL gene is associated exclusively with the development of non-papillary renal cell carcinomas. J Pathol 179: 157-161. [http://dx.doi.org/10.1002/\(SICI\)1096-9896\(199606\)179:2<157::AID-PATH557>3.0.CO;2-S](http://dx.doi.org/10.1002/(SICI)1096-9896(199606)179:2<157::AID-PATH557>3.0.CO;2-S).
- Kennedy, R and Luhmann, J. (1999). The 'Ouchless Emergency Department' getting closer: Advances in decreasing distress during painful procedures in the emergency department. 46/6 (1215-1247).
- Kernan, G; Ji, B-T; Dosemeci, M; Silverman, D; Balbus, J; Zahm, S. (1999). Occupational risk factors for pancreatic cancer: A case-control study based on death certificates from 24 U.S. States. Am J Ind Med 36: 260-270. [http://dx.doi.org/10.1002/\(SICI\)1097-0274\(199908\)36:2<260::AID-AJIM5>3.0.CO;2-P](http://dx.doi.org/10.1002/(SICI)1097-0274(199908)36:2<260::AID-AJIM5>3.0.CO;2-P).
- Keshava, N and Caldwell, J. (2006). Key issues in the role of peroxisome proliferator-activated receptor agonism and cell signaling in trichloroethylene toxicity. Environ Health Perspect 114: 1464-1470.
- Keshava, N; Chiu, W; Caldwell, J. (2007). PPARalpha and TCE: Keshava et al. respond. Environ Health Perspect 115: A15-A16. <http://dx.doi.org/10.1289/ehp.115-a15>.
- Kessler, W; Numtip, W; Grote, K; Csanady, G; Chahoud, I; Filser, J. (2004). Blood burden of di(2-ethylhexyl) phthalate and its primary metabolite mono(2-ethylhexyl) phthalate in pregnant and nonpregnant rats and marmosets. Toxicol Appl Pharmacol 195: 142-153. <http://dx.doi.org/10.1016/j.taap.2003.11.014>.
- Ketcha, M; Stevens, D; Warren, D; Bishop, C; Brashear, W. (1996). Conversion of trichloroacetic acid to dichloroacetic acid in biological samples. J Anal Toxicol 20: 236-241.
- Keys, D; Bruckner, J; Muralidhara, S; Fisher, J. (2003). Tissue dosimetry expansion and cross-validation of rat and mouse physiologically based pharmacokinetic models for trichloroethylene. Toxicol Sci 76: 35-50. <http://dx.doi.org/10.1093/toxsci/kfg212>.
- Keys, D; Schultz, I; Mahle, D; Fisher, J. (2004). A quantitative description of suicide inhibition of dichloroacetic acid in rats and mice. Toxicol Sci 82: 381-393. <http://dx.doi.org/10.1093/toxsci/kfh276>.
- Kezic, S; Monster, A; Kruse, J; Verberk, M. (2000). Skin absorption of some vaporous solvents in volunteers. Int Arch Occup Environ Health 73: 415-422. <http://dx.doi.org/10.1007/s004200000161>.
- Kezic, S; Monster, A; van de Gevel, I; Kruse, J; Opdam, J; Verberk, M. (2001). Dermal absorption of neat liquid solvents on brief exposures in volunteers. 62: 12-18. <http://dx.doi.org/10.1080/15298660108984604>.
- Khan, M; Kaphalia, B; Prabhakar, B; Kanz, M; Ansari, G. (1995). Trichloroethene-induced autoimmune response in female MRL +/+ mice. Toxicol Appl Pharmacol 134: 155-160. <http://dx.doi.org/10.1006/taap.1995.1179>.
- Khan, M; Wu, X; Ansari, G. (2001). Anti-malondialdehyde antibodies in MRL+/- mice treated with trichloroethene and dichloroacetyl chloride: possible role of lipid peroxidation in autoimmunity. Toxicol Appl Pharmacol 170: 88-92. <http://dx.doi.org/10.1006/taap.2000.9086>.
- Khandwala, H; McCutcheon, I; Flyvbjerg, A; Friend, K. (2000). The effects of insulin-like growth factors on tumorigenesis and neoplastic growth. Endocr Rev 21: 215-244. <http://dx.doi.org/10.1210/er.21.3.215>.
- Kilburn, K; Warshaw, R; Thornton, J. (1987). Formaldehyde impairs memory, equilibrium, and dexterity in histology technicians: Effects which persist for days after exposure. Arch Environ Occup Health 42: 117-120.

- Kilburn, K and Warshaw, R. (1992a). Prevalence of symptoms of systemic lupus erythematosus (SLE) and of fluorescent antinuclear antibodies associated with chronic exposure to trichloroethylene and other chemicals in well water. Environ Res 57: 1-9.
- Kilburn, K and Warshaw, R. (1992b). Neurobehavioral effects of formaldehyde and solvents on histology technicians: Repeated testing across time. Environ Res 58: 134-146.
[http://dx.doi.org/10.1016/S0013-9351\(05\)80210-5](http://dx.doi.org/10.1016/S0013-9351(05)80210-5).
- Kilburn, K and Warshaw, R. (1993a). Effects on neurobehavioral performance of chronic exposure to chemically contaminated well water. Toxicol Ind Health 9: 391-404.
- Kilburn, K and Warshaw, R. (1993b). Neurobehavioral testing of subjects exposed residentially to groundwater contaminated from an aluminum die-casting plant and local referents. J Toxicol Environ Health 39: 483-496.
- Kilburn, K and Thornton, J. (1996). Prediction equations for simple and visual two-choice reactions times in environmental neurotoxicology. Arch Environ Health 51: 439-444.
<http://dx.doi.org/10.1080/00039896.1996.9936043>.
- Kilburn, K. (2002a). Is neurotoxicity associated with environmental trichloroethylene (TCE)? Arch Environ Health 57: 113-120. <http://dx.doi.org/10.1080/00039890209602925>.
- Kilburn, K. (2002b). Do duration of exposure, proximity to electronic manufacturing plants, and involvement in a lawsuit affect chlorinated solvent toxicity. Arch Environ Health 57: 121-126.
<http://dx.doi.org/10.1080/00039890209602926>.
- Kim, C; Jackson, E; Woolfenden, A; Lawrence, S; Babar, I; Vogel, S; Crowley, D; Bronson, R; Jacks, T. (2005). Identification of bronchioalveolar stem cells in normal lung and lung cancer. Cell 121: 823-835. <http://dx.doi.org/10.1016/j.cell.2005.03.032>.
- Kim, D and Ghanayem, B. (2006). Comparative metabolism and disposition of trichloroethylene in Cyp2e1-/- and wild-type mice. Drug Metab Dispos 34: 2020-2027.
<http://dx.doi.org/10.1124/dmd.106.010538>.
- Kim, H; Bruckner, J; Dallas, C; Gallo, J. (1990a). Effect of dosing vehicles on the pharmacokinetics of orally administered carbon tetrachloride in rats. Toxicol Appl Pharmacol 102: 50-60.
[http://dx.doi.org/10.1016/0041-008X\(90\)90082-6](http://dx.doi.org/10.1016/0041-008X(90)90082-6).
- Kim, H; Odend'hal, S; Bruckner, J. (1990b). Effect of oral dosing vehicles on the acute hepatotoxicity of carbon tetrachloride in rats. Toxicol Appl Pharmacol 102: 34-49. [http://dx.doi.org/10.1016/0041-008X\(90\)90081-5](http://dx.doi.org/10.1016/0041-008X(90)90081-5).
- Kim, S; Kupke, K; Ierardi-Curto, L; Holme, E; Greter, J; Tanguay, R; Poudrier, J; D'Astous, M; Lettre, F; Hahn, S; Levy, H. (2000). Hepatocellular carcinoma despite long-term survival in chronic tyrosinaemia I. J Inherit Metab Dis 23: 791-804. <http://dx.doi.org/10.1023/A:1026756501669>.
- Kim, S; Kim, D; Pollack, G; Collins, L; Rusyn, I. (2009). Pharmacokinetic analysis of trichloroethylene metabolism in male B6C3F1 mice: Formation and disposition of trichloroacetic acid, dichloroacetic acid, S-(1,2-dichlorovinyl)glutathione and S-(1,2-dichlorovinyl)-L-cysteine. Toxicol Appl Pharmacol 238: 90-99. <http://dx.doi.org/10.1016/j.taap.2009.04.019>.
- Kimmerle, G and Eben, A. (1973a). Metabolism, excretion and toxicology of trichloroethylene after inhalation: 2. Experimental human exposure. Arch Toxicol 30: 127-138.
<http://dx.doi.org/10.1007/BF02425930>.
- Kimmerle, G and Eben, A. (1973b). Metabolism, excretion and toxicology of trichloroethylene after inhalation: 1. Experimental exposure on rats. Arch Toxicol 30: 115-126.
<http://dx.doi.org/10.1007/BF02425929>.
- King, G; Smialek, J; Troutman, W. (1985). Sudden death in adolescents resulting from the inhalation of typewriter correction fluid. JAMA 253: 1604-1606.
- Kioski, C; Heubach, S; Petersen, N. (1990a). The incidence of childhood leukemia and testicular cancer in Pima County: 1970-1986. Phoenix, AZ: Arizona Department of Health Services.
- Kioski, C; Petersen, N; Heubach, H; Chapin, C. (1990b). Cancer incidence and mortality in an East Phoenix area overlaying groundwater contaminated with volatile organic compounds. Phoenix, AZ: Arizona Department of Health Services.

- Kiran, S; Cocco, P; Mannetje, A; Satta, G; D'Andrea, I; Becker, N; de Sanjosé, S; Foretova, L; Staines, A; Kleefeld, S; Maynadié, M; Nieters, A; Brennan, P; Boffetta, P. (2010). Occupational exposure to ethylene oxide and risk of lymphoma. *Epidemiology* 21: 905-910. <http://dx.doi.org/10.1097/EDE.0b013e3181f4cc0f>.
- Kishi, R; Harabuchi, I; Ikeda, T; Katakura, Y; Miyake, H. (1993). Acute effects of trichloroethylene on blood concentrations and performance decrements in rats and their relevance to humans. *Occup Environ Med* 50: 470-480.
- Kitada, T; Seki, S; Kawakita, N; Kuroki, T; Monna, T. (1995). Telomere shortening in chronic liver diseases. *Biochem Biophys Res Commun* 211: 33-39. <http://dx.doi.org/10.1006/bbrc.1995.1774>.
- Kjellstrand, P; Lanke, J; Bjerkemo, M; Zetterqvist, L; Måansson, L. (1980). Irreversible effects of trichloroethylene exposure on the central nervous system. *Scand J Work Environ Health* 6: 40-47.
- Kjellstrand, P; Bjerkemo, M; Mortensen, I; Måansson, L. (1981a). Effects of long-term exposure to trichloroethylene on the behavior of Mongolian gerbils (*Meriones unguiculatus*). *J Toxicol Environ Health A* 8: 787-793. <http://dx.doi.org/10.1080/15287398109530114>.
- Kjellstrand, P; Kanje, M; Måansson, L; Bjerkemo, M; Mortensen, I; Lanke, J; Holmquist, B. (1981b). Trichloroethylene: Effects on body and organ weights in mice, rats and gerbils. *Toxicology* 21: 105-115. [http://dx.doi.org/10.1016/0300-483X\(81\)90121-9](http://dx.doi.org/10.1016/0300-483X(81)90121-9).
- Kjellstrand, P; Holmquist, B; Alm, P; Kanje, M; Romare, S; Jonsson, I; Måansson, L; Bjerkemo, M. (1983a). Trichloroethylene: Further studies of the effects on body and organ weights and plasma butyrylcholinesterase activity in mice. *Acta Pharmacol Toxicol* 53: 375-384. <http://dx.doi.org/10.1111/j.1600-0773.1983.tb03438.x>.
- Kjellstrand, P; Holmquist, B; Mandahl, N; Bjerkemo, M. (1983b). Effects of continuous trichloroethylene inhalation on different strains of mice. *Acta Pharmacol Toxicol* 53: 369-374. <http://dx.doi.org/10.1111/j.1600-0773.1983.tb03437.x>.
- Kjellstrand, P; Kanje, M; Bjerkemo, M. (1987). Regeneration of the sciatic nerve in mice and rats exposed to trichloroethylene. *Toxicol Lett* 38: 187-191. [http://dx.doi.org/10.1016/0378-4274\(87\)90127-5](http://dx.doi.org/10.1016/0378-4274(87)90127-5).
- Klaunig, J; Babich, M; Baetcke, K; Cook, J; Corton, J; David, R; DeLuca, J; Lai, D; McKee, R; Peters, J; Roberts, R; Fenner-Crisp, P. (2003). PPARalpha agonist-induced rodent tumors: Modes of action and human relevance. *Crit Rev Toxicol* 33: 655-780. <http://dx.doi.org/10.1080/713608372>.
- Kligerman, A; Bryant, M; Doerr, C; Erexson, G; Evansky, P; Kwanyuen, P; McGee, J. (1994). Inhalation studies of the genotoxicity of trichloroethylene to rodents. *DNA Repair* 322: 87-96.
- Kobayashi, M; Ikeda, K; Hosaka, T; Sezaki, H; Someya, T; Akuta, N; Suzuki, F; Suzuki, Y; Saitoh, S; Arase, Y; Kumada, H. (2006). Dysplastic nodules frequently develop into hepatocellular carcinoma in patients with chronic viral hepatitis and cirrhosis. *Cancer* 106: 636-647. <http://dx.doi.org/10.1002/cncr.21607>.
- Koch, R; Schlegelmilch, R; Wolf, H. (1988). Genetic effects of chlorinated ethylenes in the yeast *Saccharomyces cerevisiae*. *Mutat Res* 206: 209-216. [http://dx.doi.org/10.1016/0165-1218\(88\)90162-0](http://dx.doi.org/10.1016/0165-1218(88)90162-0).
- Kochen, W; Kohlmüller, D; De Biasi, P; Ramsay, R. (2003). The endogenous formation of highly chlorinated tetrahydro-beta-carbolines as a possible causative mechanism in idiopathic Parkinson's disease. *Adv Exp Med Biol* 527: 253-263.
- Koizumi, A. (1989). Potential of physiologically based pharmacokinetics to amalgamate kinetic data of trichloroethylene and tetrachloroethylene obtained in rats and man. *Br J Ind Med* 46: 239-249.
- Konietzko, H; Elster, I; Sayer, H; Weichardt, H. (1975). Central nervous system lesions caused by trichloroethylene <Original> Zentralnervose schaden durch trichlorathylen. 35: 240-241.
- Kopylev, L; Chen, C; White, P. (2007). Towards quantitative uncertainty assessment for cancer risks: Central estimates and probability distributions of risk in dose-response modeling. *Regul Toxicol Pharmacol* 49: 203-207. <http://dx.doi.org/10.1016/j.yrtph.2007.08.002>.

- Korrapati, M; Lock, E; Mehendale, H. (2005). Molecular mechanisms of enhanced renal cell division in protection against S-1,2-dichlorovinyl-L-cysteine-induced acute renal failure and death. Am J Physiol Renal Physiol 289: F175-F185. <http://dx.doi.org/10.1152/ajprenal.00418.2004>.
- Kozak, E and Tate, S. (1982). Glutathione-degrading enzymes of microvillus membranes. J Biol Chem 257: 6322-6327.
- Krasowski, M and Harrison, N. (2000). The actions of ether, alcohol and alkane general anaesthetics on GABA_A and glycine receptors and the effects of TM2 and TM3 mutations. Br J Pharmacol 129: 731-743. <http://dx.doi.org/10.1038/sj.bjp.0703087>.
- Kraupp-Grasl, B; Huber, W; Putz, B; Gerbracht, U; Schulte-Hermann, R. (1990). Tumor promotion by the peroxisome proliferator nafenopin involving a specific subtype of altered foci in rat liver. Cancer Res 50: 3701-3708.
- Krause, R; Lash, L; Elfarra, A. (2003). Human kidney flavin-containing monooxygenases and their potential roles in cysteine s-conjugate metabolism and nephrotoxicity. J Pharmacol Exp Ther 304: 185-191. <http://dx.doi.org/10.1124/jpet.102.042911>.
- Krejci, M; Ridgewell, R; Koechel, D. (1991). Acute effects of the D-isomer of S-(1,2-dichlorovinyl)cysteine on renal function and ultrastructure in the pentobarbital-anesthetized dog: site-specific toxicity involving the S1 and S2 cells of the proximal tubule. Toxicology 69: 151-164.
- Krishnadasan, A; Kennedy, N; Zhao, Y; Morgenstern, H; Ritz, B. (2007). Nested case-control study of occupational chemical exposures and prostate cancer in aerospace and radiation workers. Am J Ind Med 50: 383-390. <http://dx.doi.org/10.1002/ajim.20458>.
- Krishnadasan, A; Kennedy, N; Zhao, Y; Morgenstern, H; Ritz, B. (2008). Nested case-control study of occupational physical activity and prostate cancer among workers using a job exposure matrix. Cancer Causes Control 19: 107-114. <http://dx.doi.org/10.1007/s10552-007-9076-7>.
- Kroneld, R. (1989). Volatile pollutants in the environment and human tissues. Bull Environ Contam Toxicol 42: 873-877. <http://dx.doi.org/10.1007/BF01701629>.
- Kubo, Y; Mitani, H; Hino, O. (1994). Allelic loss at the predisposing gene locus in spontaneous and chemically induced renal cell carcinomas in the Eker rat. Cancer Res 54: 2633-2635.
- Kudryavtsev, B; Kudryavtseva, M; Sakuta, G; Stein, G. (1993). Human hepatocyte polyploidization kinetics in the course of life cycle. Virchows Arch B Cell Pathol Incl Mol Pathol 64: 387-393. <http://dx.doi.org/10.1007/BF02915139>.
- Kulig, B. (1987). The effects of chronic trichloroethylene exposure on neurobehavioral functioning in the rat. Neurotoxicol Teratol 9: 171-178. [http://dx.doi.org/10.1016/0892-0362\(87\)90095-X](http://dx.doi.org/10.1016/0892-0362(87)90095-X).
- Kulkarni, K; Barcak, E; El-Serag, H; Goodgame, R. (2004). The impact of immigration on the increasing incidence of hepatocellular carcinoma in the United States. Aliment Pharmacol Ther 20: 445-450. <http://dx.doi.org/10.1111/j.1365-2036.2004.02090.x>.
- Kuller, L; Ockene, J; Meilahn, E; Wentworth, D; Svendsen, K; Neaton, J. (1991). Cigarette smoking and mortality. MRFIT Research Group. Prev Med 20: 638-654. [http://dx.doi.org/10.1016/0091-7435\(91\)90060-H](http://dx.doi.org/10.1016/0091-7435(91)90060-H).
- Kumar, M; Kumar, R; Hissar, S; Saraswat, M; Sharma, B; Sahuja, P; Sarin, S. (2007). Risk factors analysis for hepatocellular carcinoma in patients with and without cirrhosis: A case-control study of 213 hepatocellular carcinoma patients from India. J Gastroenterol Hepatol 22: 1104-1111. <http://dx.doi.org/10.1111/j.1440-1746.2007.04908.x>.
- Kumar, M; Tewari, S; Sharma, P; Verma, V; Chauhan, L; Agarwal, S; Dwivedi, U; Goel, S. (2009). Study of genetic polymorphism in solvent exposed population and its correlation to in vitro effect of trichloroethylene on lymphocytes. J Environ Biol 30: 685-691.
- Kumar, P; Prasad, A; Dutta, K. (2000a). Steroidogenic alterations in testes and sera of rats exposed to trichloroethylene (TCE) by inhalation. Hum Exp Toxicol 19: 117-121.
- Kumar, P; Prasad, A; Saxena, D; Manu, U; Maji, B; Dutta, K. (2000b). Fertility and general reproduction studies in trichloroethylene exposed rats. Indian Journal of Occupational Health 43: 117-126.

- Kumar, P; Prasad, A; Maji, B; Mani, U; Dutta, K. (2001a). Hepatotoxic alterations induced by inhalation of trichlorethylene (TCE) in rats. *Biomed Environ Sci* 14: 325-332.
- Kumar, P; Prasad, A; Mani, U; Maji, B; Dutta, K. (2001b). Trichloroethylene induced testicular toxicity in rats exposed by inhalation. *Hum Exp Toxicol* 20: 585-589.
<http://dx.doi.org/10.1191/096032701718620882>.
- Kung, H-C; Hoyert, D; Xu, J; Murphy, S. (2008). Deaths: Final data for 2005. *National Vital Statistics Reports* 56.
- Kurasawa, K. (1988). Selective damage of pulmonary nonciliated bronchiolar epithelial (Clara) cells by trichloroethylene in rats. *Sangyo Eiseigaku Zasshi* 30: 121-133.
- Kvam, B; Romundstad, P; Boffetta, P; Andersen, A. (2005). Cancer in the Norwegian printing industry. *Scand J Work Environ Health* 31: 36-43.
- Kylin, B; Axell, K; Samuel, H; Lindborg, A. (1967). Effect of inhaled trichloroethylene on the CNS: As measured by optokinetic nystagmus. *Arch Environ Health* 15: 48-52.
- Kyrklund, T; Kjellstrand, P; Haglid, K. (1986). Fatty acid changes in rat brain ethanolamine phosphoglycerides during and following chronic exposure to trichloroethylene. *Toxicol Appl Pharmacol* 85: 145-153. [http://dx.doi.org/10.1016/0041-008X\(86\)90108-0](http://dx.doi.org/10.1016/0041-008X(86)90108-0).
- Kyrklund, T; Kjellstrand, P; Haglid, K. (1988). Effects of exposure to Freon 11, 1,1,1-trichloroethane or perchloroethylene on the lipid and fatty-acid composition of rat cerebral cortex. *Scand J Work Environ Health* 14: 91-94.
- Kyrklund, T and Haglid, K. (1991). Brain lipid composition in guinea pigs after intrauterine exposure to perchloroethylene. *Pharmacol Toxicol* 68: 146-148. <http://dx.doi.org/10.1111/j.1600-0773.1991.tb02054.x>.
- La Vecchia, C; Negri, E; Franceschi, S; D'Avanzo, B; Boyle, P. (1994). A case-control study of diabetes mellitus and cancer risk. *Br J Cancer* 70: 950-953.
- Lacey, JJ; Garabrant, D; Laing, T; Gillespie, B; Mayes, M; Cooper, B; Schottenfeld, D. (1999). Petroleum distillate solvents as risk factors for undifferentiated connective tissue disease (UCTD). *Am J Epidemiol* 149: 761-770.
- Lagakos, S; Wessen, B; Zelen, M. (1986). An analysis of contaminated well water and health effects in Woburn, Massachusetts. *J Am Stat Assoc* 81: 583-596.
- Laham, S. (1970). Studies on placental transfer: Trichlorethylene. *Ind Med Surg* 39: 46-49.
- Lake, B; Evans, J; Cunningham, M; Price, R. (1993). Comparison of the hepatic effects of nafenopin and WY-14,643 on peroxisome proliferation and cell replication in the rat and Syrian hamster. *Environ Health Perspect* 101: 241-247.
- Lalwani, N; Dethloff, L; Haskins, J; Robertson, D; de la Iglesia, F. (1997). Increased nuclear ploidy, not cell proliferation, is sustained in the peroxisome proliferator-treated rat liver. *Toxicol Pathol* 25: 165-176. <http://dx.doi.org/10.1177/019262339702500206>.
- Lamb, J and Hentz, K. (2006). Toxicological review of male reproductive effects and trichloroethylene exposure: assessing the relevance to human male reproductive health. *Reprod Toxicol* 22: 557-563. <http://dx.doi.org/10.1016/j.reprotox.2006.07.001>.
- Lan, Q; Zhang, L; Tang, X; Shen, M; Smith, M; Qiu, C; Ge, Y; Ji, Z; Xiong, J; He, J; Reiss, B; Hao, Z; Liu, S; Xie, Y; Guo, W; Purdue, M; Galvan, N; Xin, K; Hu, W; Beane Freeman, L; Blair, A; Li, L; Rothman, N; Vermeulen, R; Huang, H. (2010). Occupational exposure to trichloroethylene is associated with a decline in lymphocyte subsets and soluble CD27 and CD30 markers. *Carcinogenesis* 31: 1592-1596. <http://dx.doi.org/10.1093/carcin/bgq121>.
- Land, P; Owen, E; Linde, H. (1981). Morphologic changes in mouse spermatozoa after exposure to inhalational anesthetics during early spermatogenesis. *Anesthesiology* 54: 53-56.
- Landrigan, P; Stein, G; Kominsky, J; Ruhe, R; Watanabe, A. (1987). Common-source community and industrial exposure to trichloroethylene. *Arch Environ Occup Health* 42: 327-332.
<http://dx.doi.org/10.1080/00039896.1987.9934354>.

- Lane, S; Watts, R; Bentham, G; Innes, N; Scott, D. (2003). Are environmental factors important in primary systemic vasculitis? A case-control study. *Arthritis Rheum* 48: 814-823.
<http://dx.doi.org/10.1002/art.10830>.
- Langholz, B and Goldstein, L. (1996). Risk set sampling in epidemiologic cohort studies. *Stat Sci* 11: 35-53. <http://dx.doi.org/10.1214/ss/1032209663>.
- Laparé, S; Tardif, R; Brodeur, J. (1995). Effect of various exposure scenarios on the biological monitoring of organic solvents in alveolar air. II. 1,1,1-Trichloroethane and trichloroethylene. *Int Arch Occup Environ Health* 67: 375-394. <http://dx.doi.org/10.1007/BF00381051>.
- Lapis, K; Zalatnai, A; Timár, F; Thorgeirsson, U. (1995). Quantitative evaluation of lysozyme- and CD68-positive Kupffer cells in diethylnitrosamine-induced hepatocellular carcinomas in monkeys. *Carcinogenesis* 16: 3083-3085. <http://dx.doi.org/10.1093/carcin/16.12.3083>.
- Larsen, G and Stevens, J. (1986). Cysteine conjugate beta-lyase in the gastrointestinal bacterium *Eubacterium limosum*. *Mol Pharmacol* 29: 97-103.
- Larson, J and Bull, R. (1989). Effect of ethanol on the metabolism of trichloroethylene. *J Toxicol Environ Health A* 28: 395-406. <http://dx.doi.org/10.1080/15287398909531359>.
- Larson, J and Bull, R. (1992a). Species differences in the metabolism of trichloroethylene to the carcinogenic metabolites trichloroacetate and dichloroacetate. *Toxicol Appl Pharmacol* 115: 278-285. [http://dx.doi.org/10.1016/0041-008X\(92\)90333-N](http://dx.doi.org/10.1016/0041-008X(92)90333-N).
- Larson, J and Bull, R. (1992b). Metabolism and lipoperoxidative activity of trichloroacetate and dichloroacetate in rats and mice. *Toxicol Appl Pharmacol* 115: 268-277.
[http://dx.doi.org/10.1016/0041-008X\(92\)90332-M](http://dx.doi.org/10.1016/0041-008X(92)90332-M).
- Lash, L; Elfarra, A; Anders, M. (1986). Renal cysteine conjugate beta-lyase. Bioactivation of nephrotoxic cysteine S-conjugates in mitochondrial outer membrane. *J Biol Chem* 261: 5930-5935.
- Lash, L; Jones, D; Anders, M. (1988). Glutathione homeostasis and glutathione S-conjugate toxicity in kidney. In E Hodgson, JR Band & RM Philpot (Eds.), *Reviews in biochemical toxicology* (Vol. 9, pp. 29-67). New York, NY: Elsevier.
- Lash, L; Elfarra, A; Rakiewicz-Nemeth, D; Anders, M. (1990). Bioactivation mechanism of cytotoxic homocysteine S-conjugates. *Arch Biochem Biophys* 276: 322-330.
[http://dx.doi.org/10.1016/0003-9861\(90\)90727-G](http://dx.doi.org/10.1016/0003-9861(90)90727-G).
- Lash, L; Sausen, P; Duescher, R; Cooley, A; Elfarra, A. (1994). Roles of cysteine conjugate beta-lyase and S-oxidase in nephrotoxicity: Studies with S-(1,2-dichlorovinyl)-L-cysteine and S-(1,2-dichlorovinyl)-L-cysteine sulfoxide. *J Pharmacol Exp Ther* 269: 374-383.
- Lash, L; Xu, Y; Elfarra, A; Duescher, R; Parker, J. (1995). Glutathione-dependent metabolism of trichloroethylene in isolated liver and kidney cells of rats and its role in mitochondrial and cellular toxicity. *Drug Metab Dispos* 23: 846-853.
- Lash, L; Qian, W; Putt, D; Jacobs, K; Elfarra, A; Krause, R; Parker, J. (1998a). Glutathione conjugation of trichloroethylene in rats and mice: Sex-, species-, and tissue-dependent differences. *Drug Metab Dispos* 26: 12-19.
- Lash, L; Visarius, T; Sall, J; Qian, W; Tokarz, J. (1998b). Cellular and subcellular heterogeneity of glutathione metabolism and transport in rat kidney cells. *Toxicology* 130: 1-15.
[http://dx.doi.org/10.1016/S0300-483X\(98\)00093-6](http://dx.doi.org/10.1016/S0300-483X(98)00093-6).
- Lash, L; Lipscomb, J; Putt, D; Parker, J. (1999a). Glutathione conjugation of trichloroethylene in human liver and kidney: Kinetics and individual variation. *Drug Metab Dispos* 27: 351-359.
- Lash, L; Putt, D; Brashear, W; Abbas, R; Parker, J; Fisher, J. (1999b). Identification of S-(1,2-dichlorovinyl)glutathione in the blood of human volunteers exposed to trichloroethylene. *J Toxicol Environ Health A* 56: 1-21. <http://dx.doi.org/10.1080/009841099158204>.
- Lash, L; Fisher, J; Lipscomb, J; Parker, J. (2000a). Metabolism of trichloroethylene. *Environ Health Perspect* 108: 177-200.
- Lash, L; Parker, J; Scott, C. (2000b). Modes of action of trichloroethylene for kidney tumorigenesis. *Environ Health Perspect* 108: 225-240.

- Lash, L; Hueni, S; Putt, D. (2001a). Apoptosis, necrosis, and cell proliferation induced by S-(1,2-dichlorovinyl)-L-cysteine in primary cultures of human proximal tubular cells. *Toxicol Appl Pharmacol* 177: 1-16. <http://dx.doi.org/10.1006/taap.2001.9295>.
- Lash, L; Qian, W; Putt, D; Hueni, S; Elfarra, A; Krause, R; Parker, J. (2001b). Renal and hepatic toxicity of trichloroethylene and its glutathione-derived metabolites in rats and mice: Sex-, species-, and tissue-dependent differences. *J Pharmacol Exp Ther* 297: 155-164.
- Lash, L; Putt, D; Hueni, S; Krause, R; Elfarra, A. (2003). Roles of necrosis, apoptosis, and mitochondrial dysfunction in S-(1,2-dichlorovinyl)-L-cysteine sulfoxide-induced cytotoxicity in primary cultures of human renal proximal tubular cells. *J Pharmacol Exp Ther* 305: 1163-1172. <http://dx.doi.org/10.1124/jpet.102.046185>.
- Lash, L; Putt, D; Parker, J. (2006). Metabolism and tissue distribution of orally administered trichloroethylene in male and female rats: Identification of glutathione- and cytochrome P-450-derived metabolites in liver, kidney, blood, and urine. *J Toxicol Environ Health A* 69: 1285-1309. <http://dx.doi.org/10.1080/15287390500360133>.
- Lash, L; Putt, D; Huang, P; Hueni, S; Parker, J. (2007). Modulation of hepatic and renal metabolism and toxicity of trichloroethylene and perchloroethylene by alterations in status of cytochrome P450 and glutathione. *Toxicology* 235: 11-26. <http://dx.doi.org/10.1016/j.tox.2007.03.001>.
- Lash, T and Green, L. (1993). Blink reflex measurement of effects of trichloroethylene exposure on the trigeminal nerve [letter]. *Muscle Nerve* 16: 217-220.
- Laslo-Baker, D; Barrera, M; Knittel-Keren, D; Kozer, E; Wolpin, J; Khattak, S; Hackman, R; Rovet, J; Koren, G. (2004). Child neurodevelopmental outcome and maternal occupational exposure to solvents. *Arch Pediatr Adolesc Med* 158: 956-961. <http://dx.doi.org/10.1001/archpedi.158.10.956>.
- Latendresse, J and Pereira, M. (1997). Dissimilar characteristics of n-methyl-n-nitrosourea-initiated foci and tumors promoted by dichloroacetic acid or trichloroacetic acid in the liver of female B6C3F1 mice. *Toxicol Pathol* 25: 433-440. <http://dx.doi.org/10.1177/019262339702500501>.
- Lau, S; Kleiner, H; Monks, T. (1995). Metabolism as a determinant of species susceptibility to 2,3,5-(triglutathion-S-yl)hydroquinone-mediated nephrotoxicity. The role of N-acetylation and N-deacetylation. *Drug Metab Dispos* 23: 1136-1142.
- Laughter, A; Dunn, C; Swanson, C; Howroyd, P; Cattley, R; Corton, J. (2004). Role of the peroxisome proliferator-activated receptor alpha (PPAR α) in responses to trichloroethylene and metabolites, trichloroacetate and dichloroacetate in mouse liver. *Toxicology* 203: 83-98. <http://dx.doi.org/10.1016/j.tox.2004.06.014>.
- Laureno, R. (1988). Trichlorethylene neurology. *Can J Neurol Sci* 15: 354.
- Laureno, R. (1993). Trichloroethylene does not cause trigeminal neuropathy [letter]. *Muscle Nerve* 16: 217.
- Lawrence, T; Hagemann, T; Balkwill, F. (2007). Cancer. Sex, cytokines, and cancer. *Science* 317: 51-52. <http://dx.doi.org/10.1126/science.1146052>.
- Le Mesurier, S; Lykke, A; Stewart, B. (1980). Reduced yield of pulmonary surfactant: patterns of response following administration of chemicals to rats by inhalation. *Toxicol Lett* 5: 89-93.
- Leakey, J; Seng, J; Allaben, W. (2003a). Body weight considerations in the B6C3F1 mouse and the use of dietary control to standardize background tumor incidence in chronic bioassays. *Toxicol Appl Pharmacol* 193: 237-265.
- Leakey, J; Seng, J; Latendresse, J; Hussain, N; Allen, L; Allaben, W. (2003b). Dietary controlled carcinogenicity study of chloral hydrate in male B6C3F1 mice. *Toxicol Appl Pharmacol* 193: 266-280. <http://dx.doi.org/10.1016/j.taap.2003.07.007>.
- Leandri, M; Schizzi, R; Scielzo, C; Favale, E. (1995). Electrophysiological evidence of trigeminal root damage after trichloroethylene exposure. *Muscle Nerve* 18: 467-468. <http://dx.doi.org/10.1002/mus.880180416>.
- Leavitt, S; DeAngelo, A; George, M; Ross, J. (1997). Assessment of the mutagenicity of dichloroacetic acid in lacI transgenic B6C3F1 mouse liver. *Carcinogenesis* 18: 2101-2106.

- Ledda-Columbano, G; Columbano, A; Curto, M; Ennas, M; Coni, P; Sarma, D; Pani, P. (1989). Further evidence that mitogen-induced cell proliferation does not support the formation of enzyme-altered islands in rat liver by carcinogens. *Carcinogenesis* 10: 847-850.
<http://dx.doi.org/10.1093/carcin/10.5.847>.
- Ledda-Columbano, G; Coni, P; Simbula, G; Zedda, I; Columbano, A. (1993). Compensatory regeneration, mitogen-induced liver growth, and multistage chemical carcinogenesis. *Environ Health Perspect* 101: 163-168.
- Ledda-Columbano, G; Curto, M; Piga, R; Zedda, A; Menegazzi, M; Sartori, C; Shinozuka, H; Bluethmann, H; Poli, V; Ciliberto, G; Columbano, A. (1998). In vivo hepatocyte proliferation is inducible through a TNF and IL-6-independent pathway. *Oncogene* 17: 1039-1044.
<http://dx.doi.org/10.1038/sj.onc.1202018>.
- Ledda-Columbano, G; Perra, A; Concias, D; Cossu, C; Molotzu, F; Sartori, C; Shinozuka, H; Columbano, A. (2003). Different effects of the liver mitogens triiodo-thyronine and ciprofibrate on the development of rat hepatocellular carcinoma. *Toxicol Pathol* 31: 113-120.
<http://dx.doi.org/10.1080/01926230390173851>.
- Lee, D; Park, I; Kim, J; Lee, Y; Kim, D; Kang, S-K. (1998). Neurobehavioral changes in shoe manufacturing workers. *Neurotoxicol Teratol* 20: 259-263. [http://dx.doi.org/10.1016/S0892-0362\(97\)00130-X](http://dx.doi.org/10.1016/S0892-0362(97)00130-X).
- Lee, J-S; Chu, I-S; Heo, J; Calvisi, D; Sun, Z; Roskams, T; Durnez, A; Demetris, A; Thorgeirsson, S. (2004a). Classification and prediction of survival in hepatocellular carcinoma by gene expression profiling. *Hepatology* 40: 667-676. <http://dx.doi.org/10.1002/hep.20375>.
- Lee, J-S; Chu, I-S; Mikaelyan, A; Calvisi, D; Heo, J; Reddy, J; Thorgeirsson, S. (2004b). Application of comparative functional genomics to identify best-fit mouse models to study human cancer. *Nat Genet* 36: 1306-1311. <http://dx.doi.org/10.1038/ng1481>.
- Lee, K; Bruckner, J; Muralidhara, S; Gallo, J. (1996). Characterization of presystemic elimination of trichloroethylene and its nonlinear kinetics in rats. *Toxicol Appl Pharmacol* 139: 262-271.
<http://dx.doi.org/10.1006/taap.1996.0165>.
- Lee, K; Muralidhara, S; Schnellmann, R; Bruckner, J. (2000a). Contribution of direct solvent injury to the dose-dependent kinetics of trichloroethylene: Portal vein administration to rats. *Toxicol Appl Pharmacol* 164: 46-54. <http://dx.doi.org/10.1006/taap.2000.8891>.
- Lee, K; Muralidhara, S; White, C; Bruckner, J. (2000b). Mechanisms of the dose-dependent kinetics of trichloroethylene: Oral bolus dosing of rats. *Toxicol Appl Pharmacol* 164: 55-64.
<http://dx.doi.org/10.1006/taap.2000.8892>.
- Lee, L-H; Chan, C-C; Chung, C-W; Ma, Y-C; Wang, G-S; Wang, J-D. (2002). Health risk assessment on residents exposed to chlorinated hydrocarbons contaminated in groundwater of a hazardous waste site. *J Toxicol Environ Health A* 65: 219-235. <http://dx.doi.org/10.1080/15287390252800828>.
- Lee, L-H; Chung, C-W; Ma, Y-C; Wang, G-S; Chen, P-C; Hwang, Y-H; Wang, J-D. (2003). Increased mortality odds ratio of male liver cancer in a community contaminated by chlorinated hydrocarbons in groundwater. *Occup Environ Med* 60: 364-369.
<http://dx.doi.org/10.1136/oem.60.5.364>.
- Lehmann, I; Rehwagen, M; Diez, U; Seiffert, A; Rolle-Kampczyk, U; Richter, M; Wetzig, H; Borte, M; Herbarth, O. (2001). Enhanced in vivo IgE production and T cell polarization toward the type 2 phenotype in association with indoor exposure to VOC: Results of the LARS study. *Int J Hyg Environ Health* 204: 211-221. <http://dx.doi.org/10.1078/1438-4639-00100>.
- Lehmann, I; Thoelke, A; Rehwagen, M; Rolle-Kampczyk, U; Schlink, U; Schulz, R; Borte, M; Diez, U; Herbarth, O. (2002). The influence of maternal exposure to volatile organic compounds on the cytokine secretion profile of neonatal T cells. *Environ Toxicol* 17: 203-210.
<http://dx.doi.org/10.1002/tox.10055>.
- Lehnart, S; Mongillo, M; Bellinger, A; Lindegger, N; Bi-Xing, C; Hsueh, W; Reiken, S; Wronska, A; Drew, L; Ward, C; Lederer, W; Kass, R; Morley, G; Marks, A. (2008). Leaky Ca²⁺.

- release channel/ryanodine receptor 2 causes seizures and sudden cardiac death in mice. *J Clin Invest* 118: 2230-2245.
- Leigh, J and Jiang, W. (1993). Liver cirrhosis deaths within occupations and industries in the California occupational mortality study. *Addiction* 88: 767-779.
- Leighton, D, Jr and Calo, J. (1981). Distribution coefficients of chlorinated hydrocarbons in dilute air-water systems for groundwater contamination applications. *Journal of Chemical and Engineering Data* 26: 382-385. <http://dx.doi.org/10.1021/je00026a010>
- Lemke, L and McQueen, C. (1995). Acetylation and its role in the mutagenicity of the antihypertensive agent hydralazine. *Drug Metab Dispos* 23: 559-565.
- Leopardi, P; Zijno, A; Bassani, B; Pacchierotti, F. (1993). In vivo studies on chemically induced aneuploidy in mouse somatic and germinal cells. *Mutat Res* 287: 119-130. [http://dx.doi.org/10.1016/0027-5107\(93\)90151-5](http://dx.doi.org/10.1016/0027-5107(93)90151-5)
- Leuschner, J and Leuschner, F. (1991). Evaluation of the mutagenicity of chloral hydrate in vitro and in vivo. *Arzneimittelforschung* 41: 1101-1103.
- Leuschner, J and Beuscher, N. (1998). Studies on the mutagenic and carcinogenic potential of chloral hydrate. *Arzneimittelforschung* 48: 961-968.
- Levy, D; Lipton, R; Holzman, P. (1981). Smooth pursuit eye movements: Effects of alcohol and chloral hydrate. *J Psychiatr Res* 16: 1-11.
- Lewis, G; Reynolds, R; Johnson, A. (1984). Some effects of trichloroethylene on mouse lungs and livers. *Vascul Pharmacol* 15: 139-144.
- Lewitt, M; Brismar, K; Wang, J; Wivall-Helleryd, I-L; Sindelar, P; Gonzalez, F; Bergman, T; Bobek, G. (2001). Responses of insulin-like growth factor (IGF)-I and IGF-binding proteins to nutritional status in peroxisome proliferator-activated receptor-alpha knockout mice. *Growth Horm IGF Res* 11: 303-313. <http://dx.doi.org/10.1054/ghir.2001.0247>
- Li, H; Dai, Y; Huang, H; Li, L; Leng, S; Cheng, J; Niu, Y; Duan, H; Liu, Q; Zhang, X; Huang, X; Xie, J; Feng, Z; Wang, J; He, J; Zheng, Y. (2007). HLA-B*1301 as a biomarker for genetic susceptibility to hypersensitivity dermatitis induced by trichloroethylene among workers in China. *Environ Health Perspect* 115: 1553-1556. <http://dx.doi.org/10.1289/ehp.10325>.
- Li, T; Schultz, I; Keys, D; Campbell, J; Fisher, J. (2008). Quantitative evaluation of dichloroacetic acid kinetics in human--a physiologically based pharmacokinetic modeling investigation. *Toxicology* 245: 35-48. <http://dx.doi.org/10.1016/j.tox.2007.12.010>
- Liang, J and Brinkley, B. (1985). Chemical probes and possible targets for the induction of aneuploidy. *Basic Life Sci* 36: 491-505.
- Liang, J and Pacchierotti, F. (1988). Cytogenetic investigation of chemically-induced aneuploidy in mouse spermatocytes. *Mutat Res-Fundam Mol Mech Mutagen* 201: 325-335. [http://dx.doi.org/10.1016/0027-5107\(88\)90021-8](http://dx.doi.org/10.1016/0027-5107(88)90021-8).
- Liangpunsakul, S; Kolwankar, D; Pinto, A; Gorski, J; Hall, S; Chalasani, N. (2005). Activity of CYP2E1 and CYP3A enzymes in adults with moderate alcohol consumption: A comparison with nonalcoholics. *Hepatology* 41: 1144-1150. <http://dx.doi.org/10.1002/hep.20673>.
- Libbrecht, L. (2006). Hepatic progenitor cells in human liver tumor development. *World J Gastroenterol* 12: 6261-6265.
- Lide, DR (Ed.). (1998). *Handbook of chemistry and physics* (79th ed.). Boca Raton, FL: CRC Press.
- Lieber, C. (2004). The discovery of the microsomal ethanol oxidizing system and its physiologic and pathologic role. *Drug Metab Rev* 36: 511-529. <http://dx.doi.org/10.1081/DMR-200033441>.
- Ligocki, M; Leuenberger, C; Pankow, J. (1985). Trace organic-compounds in rain: 2. Gas scavenging of neutral organic-compounds. *Atmos Environ* 19: 1609-1617. [http://dx.doi.org/10.1016/0004-6981\(85\)90213-6](http://dx.doi.org/10.1016/0004-6981(85)90213-6).
- Lilly, P; Simmons, J; Pegram, R. (1994). Dose-dependent vehicle differences in the acute toxicity of bromodichloromethane. *Fundam Appl Toxicol* 23: 132-140.
- Lin, L; Lin, X; He, Y; Gao, S; Lin, X. (2003). Biological characteristics of HCC by ultrasound-guided aspiration biopsy and its clinical application. *World J Gastroenterol* 9: 941-945.

- Lin, Y; Brunt, E; Bowling, W; Hafenrichter, D; Kennedy, S; Flye, M; Ponder, K. (1995). Ras-transduced diethylnitrosamine-treated hepatocytes develop into cancers of mixed phenotype in vivo. *Cancer Res* 55: 5242-5250.
- Lindbohm, M-L; Sallmén, M; Kyrrönen, P; Kauppinen, T; Pukkala, E. (2009). Risk of liver cancer and exposure to organic solvents and gasoline vapors among Finnish workers. *Int J Cancer* 124: 2954-2959. <http://dx.doi.org/10.1002/ijc.24309>.
- Lindbohm, M; Taskinen, H; Sallmen, M; Hemminki, K. (1990). Spontaneous abortions among women exposed to organic solvents. *Am J Ind Med* 17: 449-463. <http://dx.doi.org/10.1002/ajim.4700170404>.
- Lindstrom, K. (1973). Psychological performances of workers exposed to various solvents. *Work Environ Health* 10: 151-155.
- Lindstrom, K. (1980). Changes in psychological performances of solvent-poisoned and solvent exposed workers. *Am J Ind Med* 1: 69-84.
- Lindström, K; Antti-Poika, M; Tola, S; Hyttiäinen, A. (1982). Psychological prognosis of diagnosed chronic organic solvent intoxication. *Neurobehav Toxicol Teratol* 4: 581-588.
- Lingohr, M; Thrall, B; Bull, R. (2001). Effects of dichloroacetate (DCA) on serum insulin levels and insulin-controlled signaling proteins in livers of male B6C3F1 mice. *Toxicol Sci* 59: 178-184.
- Lingohr, M; Bull, R; Kato-Weinstein, J; Thrall, B. (2002). Dichloroacetate stimulates glycogen accumulation in primary hepatocytes through an insulin-independent mechanism. *Toxicol Sci* 68: 508-515.
- Lipscomb, J; Mahle, D; Brashear, W; Barton, H. (1995). Dichloroacetic acid: Metabolism in cytosol. *Drug Metab Dispos* 23: 1202-1205.
- Lipscomb, J; Mahle, D; Brashear, W; Garrett, C. (1996). A species comparison of chloral hydrate metabolism in blood and liver. *Biochem Biophys Res Commun* 227: 340-350. <http://dx.doi.org/10.1006/bbrc.1996.1511>.
- Lipscomb, J; Garrett, C; Snawder, J. (1997). Cytochrome P450-dependent metabolism of trichloroethylene: Interindividual differences in humans. *Toxicol Appl Pharmacol* 142: 311-318. <http://dx.doi.org/10.1006/taap.1996.8040>.
- Lipscomb, J; Confer, P; Miller, M; Stamm, S; Snawder, J; Bandiera, S. (1998a). Metabolism of trichloroethylene and chloral hydrate by the Japanese medaka (*Oryzias latipes*) in vitro. *Environ Toxicol Chem* 17: 325-332.
- Lipscomb, J; Fisher, J; Confer, P; Byczkowski, J. (1998b). In vitro to in vivo extrapolation for trichloroethylene metabolism in humans. *Toxicol Appl Pharmacol* 152: 376-387. <http://dx.doi.org/10.1006/taap.1998.8485>.
- Lipscomb, J; Garrett, C; Snawder, J. (1998c). Use of kinetic and mechanistic data in species extrapolation of bioactivation: Cytochrome P-450 dependent trichloroethylene metabolism at occupationally relevant concentrations. *J Occup Health* 40: 110-117.
- Lipscomb, J; Teuschler, L; Swartout, J; Popken, D; Cox, T; Kedderis, G. (2003). The impact of cytochrome P450 2E1-dependent metabolic variance on a risk-relevant pharmacokinetic outcome in humans. *Risk Anal* 23: 1221-1238. <http://dx.doi.org/10.1111/j.0272-4332.2003.00397.x>.
- Little, J; Daisey, J; Nazaroff, W. (1992). Transport of subsurface contaminants into buildings: An exposure pathway for volatile organics. *Environ Sci Technol* 26: 2058-2066. <http://dx.doi.org/10.1021/es00035a001>.
- Liu, E; Jelinek, J; Pastore, Y; Guan, Y; Prchal, J; Prchal, J. (2003). Discrimination of polycythemias and thrombocytoses by novel, simple, accurate clonality assays and comparison with PRV-1 expression and BFU-E response to erythropoietin. *Blood* 101: 3294-3301. <http://dx.doi.org/10.1182/blood-2002-07-2287>.
- Liu, Y-T; Jin, C; Chen, Z; Cai, S-X; Yin, S-N; Li, G-L; Watanabe, T; Nakatsuka, H; Seiji, K; Inoue, O; Kawai, T; Ukai, H; Ikeda, M. (1988). Increased subjective symptom prevalence among workers exposed to trichloroethylene at sub-OEL levels. *Tohoku J Exp Med* 155: 183-195.

- Liu, Y; Bartlett, M; White, C; Muralidhara, S; Bruckner, J. (2009). Presystemic elimination of trichloroethylene in rats following environmentally relevant oral exposures. *Drug Metab Dispos* 37: 1994-1998. <http://dx.doi.org/10.1124/dmd.109.028100>.
- Lock, E; Barth, J; Argraves, S; Schnellmann, R. (2006). Changes in gene expression in human renal proximal tubule cells exposed to low concentrations of S-(1,2-dichlorovinyl)-l-cysteine, a metabolite of trichloroethylene. *Toxicol Appl Pharmacol* 216: 319-330. <http://dx.doi.org/10.1016/j.taap.2006.06.002>.
- Lock, E; Reed, C; McMillan, J; Oatis, J; Schnellmann, R. (2007). Lack of formic acid production in rat hepatocytes and human renal proximal tubule cells exposed to chloral hydrate or trichloroacetic acid. *Toxicology* 230: 234-243. <http://dx.doi.org/10.1016/j.tox.2006.11.055>.
- Lockey, J; Kelly, C; Cannon, G; Colby, T; Aldrich, V; Livingston, G. (1987). Progressive systemic sclerosis associated with exposure to trichloroethylene. *J Occup Med* 29: 493-496.
- Loeber, C; Hendrix, M; Diez De Pinos, S; Goldberg, S. (1988). Trichloroethylene: a cardiac teratogen in developing chick embryos. *Pediatr Res* 24: 740-744.
- Loeber, C and Runyan, R. (1990). A comparison of fibronectin, laminin, and galactosyltransferase adhesion mechanisms during embryonic cardiac mesenchymal cell migration in vitro. *Dev Biol* 140: 401-412.
- Lopreato, G; Phelan, R; Borghese, C; Beckstead, M; Mihic, S. (2003). Inhaled drugs of abuse enhance serotonin-3 receptor function. *Drug Alcohol Depend* 70: 11-15. [http://dx.doi.org/10.1016/S0376-8716\(02\)00330-7](http://dx.doi.org/10.1016/S0376-8716(02)00330-7).
- Lorente, C; Cordier, S; Bergeret, A; De Walle, H; Goujard, J; Aymé, S; Knill-Jones, R; Calzolari, E; Bianchi, F. (2000). Maternal occupational risk factors for oral clefts. Occupational Exposure and Congenital Malformation Working Group. *Scand J Work Environ Health* 26: 137-145.
- Lorimore, S; Coates, P; Wright, E. (2003). Radiation-induced genomic instability and bystander effects: Inter-related nontargeted effects of exposure to ionizing radiation. *Oncogene* 22: 7058-7069. <http://dx.doi.org/10.1038/sj.onc.1207044>.
- Lowengart, R; Peters, J; Cicioni, C; Buckley, J; Bernstein, L; Preston-Martin, S; Rappaport, E. (1987). Childhood leukemia and parents' occupational and home exposures. *J Natl Cancer Inst* 79: 39-46.
- Luebke, R; Chen, D; Dietert, R; Yang, Y; King, M; Luster, M. (2006). The comparative immunotoxicity of five selected compounds following developmental or adult exposure. *J Toxicol Environ Health B Crit Rev* 9: 1-26. <http://dx.doi.org/10.1080/15287390500194326>.
- Luedde, T; Beraza, N; Kotsikoris, V; van Loo, G; Nenci, A; De Vos, R; Roskams, T; Trautwein, C; Pasparakis, M. (2007). Deletion of NEMO/IKKgamma in liver parenchymal cells causes steatohepatitis and hepatocellular carcinoma. *Cancer Cell* 11: 119-132. <http://dx.doi.org/10.1016/j.ccr.2006.12.016>.
- Lumpkin, M; Bruckner, J; Campbell, J; Dallas, C; White, C; Fisher, J. (2003). Plasma binding of trichloroacetic acid in mice, rats, and humans under cancer bioassay and environmental exposure conditions. *Drug Metab Dispos* 31: 1203-1207. <http://dx.doi.org/10.1124/dmd.31.10.1203>.
- Lundberg, I; Alfredsson, L; Plato, N; Sverdrup, B; Klareskog, L; Kleinau, S. (1994). Occupation, occupational exposure to chemicals and rheumatological disease: A register based cohort study. *Scand J Rheumatol* 23: 305-310. <http://dx.doi.org/10.3109/03009749409099278>.
- Lundgren, B; Meijer, J; DePierre, J. (1987). Induction of cytosolic and microsomal epoxide hydrolases and proliferation of peroxisomes and mitochondria in mouse liver after dietary exposure to p-chlorophenoxyacetic acid, 2,4-dichlorophenoxyacetic acid and 2,4,5-trichlorophenoxyacetic acid. *Biochem Pharmacol* 36: 815-821. [http://dx.doi.org/10.1016/0006-2952\(87\)90169-9](http://dx.doi.org/10.1016/0006-2952(87)90169-9).
- Lundgren, B and DePierre, J. (1989). Proliferation of peroxisomes and induction of cytosolic and microsomal epoxide hydrolases in different strains of mice and rats after dietary treatment with clofibrate. *Xenobiotica* 19: 867-881.
- Lutz, W; Gaylor, D; Conolly, R; Lutz, R. (2005). Nonlinearity and thresholds in dose-response relationships for carcinogenicity due to sampling variation, logarithmic dose scaling, or small

- differences in individual susceptibility. *Toxicol Appl Pharmacol* 207: 565-569.
<http://dx.doi.org/10.1016/j.taap.2005.01.038>.
- Lybarger, J; Lichtveld, M; Amler, R. (1999). Biomedical testing of the kidney for persons exposed to hazardous substances in the environment. *Ren Fail* 21: 263-274.
- Lynch, A and Parry, J. (1993). The cytochalasin-B micronucleus/kinetochore assay in vitro: studies with 10 suspected aneugens. *Mutat Res* 287: 71-86. [http://dx.doi.org/10.1016/0027-5107\(93\)90146-7](http://dx.doi.org/10.1016/0027-5107(93)90146-7).
- Lynge, E; Anttila, A; Hemminki, K. (1997). Organic solvents and cancer. *Cancer Causes Control* 8: 406-419. <http://dx.doi.org/10.1023/A:1018461406120>.
- Macchiarini, P. (2006). Primary tracheal tumours. *Lancet Oncol* 7: 83-91.
[http://dx.doi.org/10.1016/S1470-2045\(05\)70541-6](http://dx.doi.org/10.1016/S1470-2045(05)70541-6).
- Macdonald, N; Holden, P; Roberts, R. (1999). Addition of peroxisome proliferator-activated receptor alpha to guinea pig hepatocytes confers increased responsiveness to peroxisome proliferators. *Cancer Res* 59: 4776-4780.
- Mackay, J; Fox, V; Griffiths, K; Fox, D; Howard, C; Coutts, C; Wyatt, I; Styles, J. (1995). Trichloroacetic acid: investigation into the mechanism of chromosomal damage in the in vitro human lymphocyte cytogenetic assay and the mouse bone marrow micronucleus test. *Carcinogenesis* 16: 1127-1133.
- Maeda, S; Kamata, H; Luo, J-L; Leffert, H; Karin, M. (2005). IKKbeta couples hepatocyte death to cytokine-driven compensatory proliferation that promotes chemical hepatocarcinogenesis. *Cell* 121: 977-990. <http://dx.doi.org/10.1016/j.cell.2005.04.014>.
- Magnusson, B and Kligman, A. (1969). The identification of contact allergens by animal assay. The guinea pig maximization test. *J Invest Dermatol* 52: 268-276.
- Maguin, K; Campo, P; Parietti-Winkler, C. (2009). Toluene can perturb the neuronal voltage-dependent Ca²⁺ channels involved in the middle-ear reflex. *Toxicol Sci* 107: 473-481.
<http://dx.doi.org/10.1093/toxsci/kfn242>.
- Mahle, D; Godfrey, R; Butler, G; Narayanan, L; Fisher, J; Taylor, A. (1999). Pharmacokinetics and metabolism of dichloroacetic acid and trichloroacetic acid administered in drinking water in rats and mice. Wright-Patterson AFB, OH: Human Effectiveness Directorate.
- Mahle, D; Gearhart, J; Grigsby, C; Mattie, D; Barton, H; Lipscomb, J; Cook, R. (2007). Age-dependent partition coefficients for a mixture of volatile organic solvents in Sprague-Dawley rats and humans. *J Toxicol Environ Health A* 70: 1745-1751.
<http://dx.doi.org/10.1080/15287390701458991>.
- Mahr, A; Neogi, T; Merkel, P. (2006). Epidemiology of Wegener's granulomatosis: Lessons from descriptive studies and analyses of genetic and environmental risk determinants. *Clin Exp Rheumatol* 24: S82-S91.
- Mailhes, J; Aardema, M; Marchetti, F. (1993). Investigation of aneuploidy induction in mouse oocytes following exposure to vinblastine-sulfate, pyrimethamine, diethylstilbestrol diphosphate, or chloral hydrate. *Environ Mol Mutagen* 22: 107-114. <http://dx.doi.org/10.1002/em.2850220209>.
- Maitre, A; Hours, M; Bonneterre, V; Arnaud, J; Arslan, M; Carpentier, P; Bergeret, A; de Gaudemaris, R. (2004). Systemic sclerosis and occupational risk factors: role of solvents and cleaning products. *J Rheumatol* 31: 2395-2401.
- Makwana, O; King, N; Ahles, L; Selmin, O; Granzier, H; Runyan, R. (2010). Exposure to low-dose trichloroethylene alters shear stress gene expression and function in the developing chick heart. *Cardiovasc Toxicol* 10: 100-107. <http://dx.doi.org/10.1007/s12012-010-9066-y>.
- Malarkey, D; Johnson, K; Ryan, L; Boorman, G; Maronpot, R. (2005). New insights into functional aspects of liver morphology. *Toxicol Pathol* 33: 27-34.
<http://dx.doi.org/10.1080/01926230590881826>.
- Malherbe, P; Alberati-Giani, D; Kohler, C; Cesura, A. (1995). Identification of a mitochondrial form of kynurene aminotransferase/glutamine transaminase K from rat brain. *FEBS Lett* 367: 141-144.
- Mallin, K. (1990). Investigation of a bladder cancer cluster in northwestern Illinois. *Am J Epidemiol* 132: S96-S106.

- Mally, A; Walker, C; Everitt, J; Dekant, W; Vamvakas, S. (2006). Analysis of renal cell transformation following exposure to trichloroethene in vivo and its metabolite S-(dichlorovinyl)-L-cysteine in vitro. *Toxicology* 224: 108-118. <http://dx.doi.org/10.1016/j.tox.2006.04.036>.
- Maloney, E and Waxman, D. (1999). trans-Activation of PPARalpha and PPARgamma by structurally diverse environmental chemicals. *Toxicol Appl Pharmacol* 161: 209-218. <http://dx.doi.org/10.1006/taap.1999.8809>.
- Maltoni, C and Cotti, G. (1986). Results of long-term carcinogenicity bioassays of tetrachloroethylene on Sprague-Dawley rats administered by ingestion. *Acta Oncol* 7: 11-26.
- Maltoni, C; Lefemine, G; Cotti, G. (1986). Experimental research on trichloroethylene carcinogenesis (Vol. 5). Princeton, NJ: Princeton Scientific Publishing.
- Maltoni, C; Lefemine, G; Cotti, G; Perino, G. (1988). Long-term carcinogenicity bioassays on trichloroethylene administered by inhalation to Sprague-Dawley rats and Swiss and B6C3F1 mice. *Ann N Y Acad Sci* 534: 316-342.
- Mandel, J; McLaughlin, J; Schlehofer, B; Mellemaaard, A; Helmert, U; Lindblad, P; McCredie, M; Adami, H-O. (1995). International renal-cell cancer study. IV. Occupation. *Int J Cancer* 61: 601-605. <http://dx.doi.org/10.1002/ijc.2910610503>.
- Mandel, J. (2001). Renal cell cancer correlated with occupational exposure to trichloroethylene. *J Cancer Res Clin Oncol* 127: 265-266. <http://dx.doi.org/10.1007/s004320000231>.
- Mandel, J; Kelsh, M; Mink, P; Alexander, D; Kalmes, R; Weingart, M; Yost, L; Goodman, M. (2006). Occupational trichloroethylene exposure and non-Hodgkin's lymphoma: A meta-analysis and review. *Occup Environ Med* 63: 597-607. <http://dx.doi.org/10.1136/oem.2005.022418>.
- Manson, J; Murphy, M; Richdale, N; Smith, M. (1984). Effects of oral exposure to trichloroethylene on female reproductive function. *Toxicology* 32: 229-242. [http://dx.doi.org/10.1016/0300-483X\(84\)90076-3](http://dx.doi.org/10.1016/0300-483X(84)90076-3).
- Marano, D; Boice, J, Jr; Fryzek, J; Morrison, J; Sadler, C; McLaughlin, J. (2000). Exposure assessment for a large epidemiological study of aircraft manufacturing workers. *Appl Occup Environ Hyg* 15: 644-656. <http://dx.doi.org/10.1080/10473220050075653>.
- Markwald, R; Runyan, R; Kitten, G; Funderburg, F; Bernanke, D; Brayer, P. (1984). Use of collagen gel cultures to study heart development: proteoglycan and glycoprotein interactions during formation of endocardial cushion tissue. In RL Trelstad (Ed.), *The role of extracellular matrix in development* (pp. 323-350): Alan R. Liss.
- Markwald, R; Eisenberg, C; Eisenberg, L; Trusk, T; Sugi, Y. (1996). Epithelial-mesenchymal transformations in early avian heart development. *Acta Anat* 156: 173-186.
- Maronpot, R; Anna, C; Devereux, T; Lucier, G; Butterworth, B; Anderson, M. (1995a). Considerations concerning the murine hepatocarcinogenicity of selected chlorinated hydrocarbons Growth factors and tumor promotion: Implications for risk assessment: Proceedings of the Seventh International Conference on Carcinogenesis and Risk Assessment, held in Austin, Texas, on December 1-4, 1993 (Vol. 391, pp. 305-323). New York, NY: Wiley-Liss.
- Maronpot, R; Fox, T; Malarkey, D; Goldsworthy, T. (1995b). Mutations in the ras proto-oncogene: clues to etiology and molecular pathogenesis of mouse liver tumors. *Toxicology* 101: 125-156. [http://dx.doi.org/10.1016/0300-483X\(95\)03112-S](http://dx.doi.org/10.1016/0300-483X(95)03112-S).
- Maronpot, R; Flake, G; Huff, J. (2004). Relevance of animal carcinogenesis findings to human cancer predictions and prevention. *Toxicol Pathol* 32: 40-48. <http://dx.doi.org/10.1080/01926230490425003>.
- Maronpot, R. (2007). Biological basis of differential susceptibility to hepatocarcinogenesis among mouse strains. Paper presented at the 3rd Workshop of the Standing Committee on Risk Analysis Issues and Reviews: Mouse liver tumors: Benefits and constraints on use in human health risk assessment, qualitative and quantitative aspects, Washington, DC.
- Marazzini, A; Betti, C; Bernacchi, F; Barrai, I; Barale, R. (1994). Micronucleus test and metaphase analyses in mice exposed to known and suspected spindle poisons. *Mutagenesis* 9: 505-515. <http://dx.doi.org/10.1093/mutage/9.6.505>.

- Marshall, E; Gensburg, L; Deres, D; Geary, N; Cayo, M. (1997). Maternal residential exposure to hazardous wastes and risk of central nervous system and musculoskeletal birth defects. *Arch Environ Health* 52: 416-425. <http://dx.doi.org/10.1080/00039899709602220>.
- Marsman, D; Cattley, R; Conway, J; Popp, J. (1988). Relationship of hepatic peroxisome proliferation and replicative DNA synthesis to the hepatocarcinogenicity of the peroxisome proliferators di(2-ethylhexyl)phthalate and [4-chloro-6-(2,3-xylidino)-2-pyrimidinylthio]acetic acid (Wy-14,643) in rats. *Cancer Res* 48: 6739-6744.
- Marsman, D; Goldsworthy, T; Popp, J. (1992). Contrasting hepatocytic peroxisome proliferation, lipofuscin accumulation and cell turnover for the hepatocarcinogens Wy-14,643 and clofibrate acid. *Carcinogenesis* 13: 1011-1017. <http://dx.doi.org/10.1093/carcin/13.6.1011>.
- Martinez-Larrañaga, M; Anadón, A; Diaz, M; Fernandez, R; Sevil, B; Fernandez-Cruz, M; Fernandez, M; Martinez, M; Anton, R. (1996). Induction of cytochrome P4501A1 and P4504A1 activities and peroxisomal proliferation by fumonisin B1. *Toxicol Appl Pharmacol* 141: 185-194. <http://dx.doi.org/10.1006/taap.1996.0275>.
- Mather, G; Exon, J; Koller, L. (1990). Subchronic 90 day toxicity of dichloroacetic and trichloroacetic acid in rats. *Toxicology* 64: 71-80. [http://dx.doi.org/10.1016/0300-483X\(90\)90100-U](http://dx.doi.org/10.1016/0300-483X(90)90100-U).
- Mattioli, S; Truffelli, D; Baldasseroni, A; Risi, A; Marchesini, B; Giacomini, C; Bacchini, P; Violante, F; Buiatti, E. (2002). Occupational risk factors for renal cell cancer: a case--control study in northern Italy. *J Occup Environ Med* 44: 1028-1036.
- Mazzullo, M; Bartoli, S; Bonora, B; Colacci, A; Lattanzi, G; Niero, A; Silingardi, P; Grilli, S. (1992). In vivo and in vitro interaction of trichloroethylene with macromolecules from various organs of rat and mouse. *Res Comm Chem Pathol Pharmacol* 76: 192-208.
- McCarthy, T and Jones, R. (1983). Industrial gassing poisonings due to trichlorethylene, perchlorethylene, and 1,1,1-trichloroethane, 1961-80. *Occup Environ Med* 40: 450-455.
- McCarver, D; Byun, R; Hines, R; Hichme, M; Wegenek, W. (1998). A genetic polymorphism in the regulatory sequences of human CYP2E1: Association with increased chlorzoxazone hydroxylation in the presence of obesity and ethanol intake. *Toxicol Appl Pharmacol* 152: 276-281. <http://dx.doi.org/10.1006/taap.1998.8532>.
- McCarver, D and Hines, R. (2002). The ontogeny of human drug-metabolizing enzymes: Phase II conjugation enzymes and regulatory mechanisms. *J Pharmacol Exp Ther* 300: 361-366. <http://dx.doi.org/10.1124/jpet.300.2.361>.
- McConnell, G; Ferguson, D; Pearson, C. (1975). Chlorinated hydrocarbons and the environment. *Endeavour* 34: 13-18.
- McCredie, M and Stewart, J. (1993). Risk factors for kidney cancer in New South Wales. IV. Occupation. *Br J Ind Med* 50: 349-354.
- McCunney, R. (1988). Diverse manifestations of trichloroethylene. *Br J Ind Med* 45: 122-126. <http://dx.doi.org/10.1136/oem.45.2.122>.
- McDaniel, T; Martin, P; Ross, N; Brown, S; Lesage, S; Pauli, B. (2004). Effects of chlorinated solvents on four species of North American amphibians. *Arch Environ Contam Toxicol* 47: 101-109.
- McEntee, G; Batts, K; Katzmann, J; Ilstrup, D; Nagorney, D. (1992). Relationship of nuclear DNA content to clinical and pathologic findings in patients with primary hepatic malignancy. *Surgery* 111: 376-379.
- McGregor, D; Reynolds, D; Zeiger, E. (1989). Conditions affecting the mutagenicity of trichloroethylene in Salmonella. *Environ Mol Mutagen* 13: 197-202.
- McGuire, V; Nelson, L; Koepsell, T; Checkoway, H; Longstreth, W, Jr. (1998). Assessment of occupational exposures in community-based case-control studies. *Annu Rev Public Health* 19: 35-53. <http://dx.doi.org/10.1146/annurev.publhealth.19.1.35>.
- McKinney, P; Alexander, F; Cartwright, R; Parker, L. (1991). Parental occupations of children with leukaemia in west Cumbria, north Humberside, and Gateshead. *Br Med J* 302: 681-687. <http://dx.doi.org/10.1136/bmj.302.6778.681>.

- McLaughlin, J; Hrubec, Z; Blot, W; Fraumeni, J. (1995). Smoking and cancer mortality among U.S. veterans: a 26-year follow-up. *Int J Cancer* 60: 190-193.
<http://dx.doi.org/10.1002/ijc.2910600210>.
- McLaughlin, J and Blot, W. (1997). A critical review of epidemiology studies of trichloroethylene and perchloroethylene and risk of renal-cell cancer. *Int Arch Occup Environ Health* 70: 222-231.
- McLaughlin, J; Lipworth, L; Tarone, R. (2006). Epidemiologic aspects of renal cell carcinoma. *Semin Oncol* 33: 527-533. <http://dx.doi.org/10.1053/j.seminoncol.2006.06.010>.
- McLean, A; Cogger, V; Chong, G; Warren, A; Markus, A; Dahlstrom, J; Le Couteur, D. (2003). Age-related pseudocapillarization of the human liver. *J Pathol* 200: 112-117.
<http://dx.doi.org/10.1002/path.1328>.
- McMichael, A; Spiertas, R; Gamble, J; Tousey, P. (1976). Mortality among rubber workers: Relationship to specific jobs. *J Occup Environ Med* 18: 178-185.
- McNutt, L-A; Wu, C; Xue, X; Hafner, J. (2003). Estimating the relative risk in cohort studies and clinical trials of common outcomes. *Am J Epidemiol* 157: 940-943.
<http://dx.doi.org/10.1093/aje/kwg074>.
- MDPH. (Massachusetts Department of Public Health), (1997a) Woburn childhood leukemia follow-up study: Information booklet. Boston, MA: Massachusetts Department of Public Health, Bureau of Environmental Health Assessment.
- MDPH. (Massachusetts Department of Public Health). (1997b). Woburn childhood leukemia follow-up study: Volume I: Analyses. Boston, MA: Massachusetts Department of Public Health, Bureau of Environmental Health Assessment.
http://www.mass.gov/Eohhs2/docs/dph/environmental/investigations/woburn_cancer_leukemia_follow_up_study_1997.pdf.
- MDPH. (Massachusetts Department of Public Health). (1997c). Woburn childhood leukemia follow-up study: Volume II: Appendix. Boston, MA: Massachusetts Department of Public Health, Bureau of Environmental Health Assessment.
- Mehendale, H. (2000). PPAR-alpha: A key to the mechanism of hepatoprotection by clofibrate. *Toxicol Sci* 57: 187-190.
- Melchiorri, C; Chieco, P; Zedda, A; Coni, P; Ledda-Columbano, G; Columbano, A. (1993). Ploidy and nuclearity of rat hepatocytes after compensatory regeneration or mitogen-induced liver growth. *Carcinogenesis* 14: 1825-1830. <http://dx.doi.org/10.1093/carcin/14.9.1825>.
- Melchiorri, C; Bolondi, L; Chieco, P; Pagnoni, M; Gramantieri, L; Barbara, L. (1994). Diagnostic and prognostic value of DNA ploidy and cell nuclearity in ultrasound-guided liver biopsies. *Cancer* 74: 1713-1719. [http://dx.doi.org/10.1002/1097-0142\(19940915\)74:6<1713::AID-CNCR2820740612>3.0.CO;2-O](http://dx.doi.org/10.1002/1097-0142(19940915)74:6<1713::AID-CNCR2820740612>3.0.CO;2-O).
- Mellemaaard, A; Engholm, G; McLaughlin, J; Olsen, J. (1994). Occupational risk factors for renal-cell carcinoma in Denmark. *Scand J Work Environ Health* 20: 160-165.
- Melnick, R; Jameson, C; Goehl, T; Maronpot, R; Collins, B; Greenwell, A; Harrington, F; Wilson, R; Tomaszewski, K; Agarwal, D. (1987). Application of microencapsulation for toxicology studies: II. Toxicity of microencapsulated trichloroethylene in Fischer 344 rats. *Fundam Appl Toxicol* 8: 432-442. [http://dx.doi.org/10.1016/0272-0590\(87\)90129-1](http://dx.doi.org/10.1016/0272-0590(87)90129-1).
- Melnick, R; Huff, J; Barrett, J; Maronpot, R; Lucier, G; Portier, C. (1993). Cell proliferation and chemical carcinogenesis: Symposium overview. *Environ Health Perspect* 101: 3-7.
- Melnick, R; Kohn, M; Portier, C. (1996). Implications for risk assessment of suggested nongenotoxic mechanisms of chemical carcinogenesis. *Environ Health Perspect* 104: 123-134.
- Melnick, R; Thayer, K; Bucker, J. (2008). Conflicting views on chemical carcinogenesis arising from the design and evaluation of rodent carcinogenicity studies. *Environ Health Perspect* 116: 130-135.
<http://dx.doi.org/10.1289/ehp.9989>.
- Menegazzi, M; Carcereri-De Prati, A; Suzuki, H; Shinozuka, H; Pibiri, M; Piga, R; Columbano, A; Ledda-Columbano, G. (1997). Liver cell proliferation induced by nafenopin and cyproterone acetate is not associated with increases in activation of transcription factors NF-KB and AP-1 or

- with expression of tumor necrosis factor α . Hepatology 25: 585-592.
<http://dx.doi.org/10.1002/hep.510250316>.
- Mensing, T; Welge, P; Voss, B; Fels, L; Fricke, H; Brüning, T; Wilhelm, M. (2002). Renal toxicity after chronic inhalation exposure of rats to trichloroethylene. Toxicol Lett 128: 243-247.
- Merdink, J; Gonzalez-Leon, A; Bull, R; Schultz, I. (1998). The extent of dichloroacetate formation from trichloroethylene, chloral hydrate, trichloroacetate, and trichloroethanol in B6C3F1 mice. Toxicol Sci 45: 33-41. <http://dx.doi.org/10.1006/toxs.1998.2500>.
- Merdink, J; Stenner, R; Stevens, D; Parker, J; Bull, R. (1999). Effect of enterohepatic circulation on the pharmacokinetics of chloral hydrate and its metabolites in F344 rats. J Toxicol Environ Health A 57: 357-368. <http://dx.doi.org/10.1080/009841099157665>.
- Merdink, J; Bull, R; Schultz, I. (2000). Trapping and identification of the dichloroacetate radical from the reductive dehalogenation of trichloroacetate by mouse and rat liver microsomes. Free Radic Biol Med 29: 125-130. [http://dx.doi.org/10.1016/S0891-5849\(00\)00330-0](http://dx.doi.org/10.1016/S0891-5849(00)00330-0).
- Merdink, J; Robison, L; Stevens, D; Hu, M; Parker, J; Bull, R. (2008). Kinetics of chloral hydrate and its metabolites in male human volunteers. Toxicology 245: 130-140.
<http://dx.doi.org/10.1016/j.tox.2007.12.018>.
- Mergler, D; Huel, G; Bowler, R; Frenette, B; Cone, J. (1991). Visual dysfunction among former microelectronics assembly workers. Arch Environ Health 46: 326-334.
- Merrick, B; Robinson, M; Condie, L. (1989). Differing hepatotoxicity and lethality after subacute trichloroethylene exposure in aqueous or corn oil gavage vehicles in B6C3F1 mice. J Appl Toxicol 9: 15-21. <http://dx.doi.org/10.1002/jat.2550090105>.
- Messing, K; Punnett, L; Bond, M; Alexanderson, K; Pyle, J; Zahm, S; Wegman, D; Stock, S; de Grosbois, S. (2003). Be the fairest of them all: challenges and recommendations for the treatment of gender in occupational health research. Am J Ind Med 43: 618-629.
<http://dx.doi.org/10.1002/ajim.10225>.
- Mester, B; Nieters, A; Deeg, E; Elsner, G; Becker, N; Seidler, A. (2006). Occupation and malignant lymphoma: A population based case control study in Germany. Occup Environ Med 63: 17-26.
<http://dx.doi.org/10.1136/oem.2005.020453>.
- Meyer, K; Lee, J; Dyck, P; Cao, W; Rao, M; Thorgeirsson, S; Reddy, J. (2003). Molecular profiling of hepatocellular carcinomas developing spontaneously in acyl-CoA oxidase deficient mice: Comparison with liver tumors induced in wild-type mice by a peroxisome proliferator and a genotoxic carcinogen. Carcinogenesis 24: 975-984. <http://dx.doi.org/10.1093/carcin/bgg040>.
- Meyers, ME. (2003). Memo regarding Former Endicott Site Indoor Air Test Data from IBM (International Business Machines Corporation) Program Manager Mitchell E. Meyers to New York State Department of Environmental Conservation, Engineering Geology Section Chief William E. Wertz.
- Mhiri, C; Choyakh, F; Ben Hmida, M; Feki, I; Ben Messaoud, M; Zouari, N. (2004). Trigeminal somatosensory evoked potentials in trichloroethylene-exposed workers. Neurosciences 9: 102-107.
- Michalik, L; Desvergne, B; Dreyer, C; Gavillet, M; Laurini, R; Wahli, W. (2002). PPAR expression and function during vertebrate development. Int J Dev Biol 46: 105-114.
- Michel, C; Desdouets, C; Slaoui, M; Isaacs, K; Roberts, R; Boitier, E. (2007). Diethylnitrosamine initiation does not alter clofibrate acid-induced hepatocarcinogenesis in the rat. Toxicol Sci 99: 58-69. <http://dx.doi.org/10.1093/toxsci/kfm168>.
- Mikol, Y; Hoover, K; Creasia, D; Poirier, L. (1983). Hepatocarcinogenesis in rats fed methyl-deficient, amino acid-defined diets. Carcinogenesis 4: 1619-1629.
- Miligi, L; Costantini, A; Benvenuti, A; Kriebel, D; Bolejack, V; Tumino, R; Ramazzotti, V; Rodella, S; Stagnaro, E; Crosignani, P; Amadori, D; Mirabelli, D; Sommani, L; Belletti, I; Troschel, L; Romeo, L; Miceli, G; Tozzi, G; Mendico, I; Vineis, P. (2006). Occupational exposure to solvents and the risk of lymphomas. Epidemiology 17: 552-561.
<http://dx.doi.org/10.1097/01.ede.0000231279.30988.4d>.

- Miller, B and Adler, I.D. (1992). Aneuploidy induction in mouse spermatocytes. *Mutagenesis* 7: 69-76.
- Miller, R and Guengerich, F. (1982). Oxidation of trichloroethylene by liver microsomal cytochrome P-450: Evidence for chlorine migration in a transition state not involving trichloroethylene oxide. *Biochemistry* 21: 1090-1097. <http://dx.doi.org/10.1021/bi00534a041>.
- Miller, R and Guengerich, F. (1983). Metabolism of trichloroethylene in isolated hepatocytes, microsomes, and reconstituted enzyme systems containing cytochrome P-450. *Cancer Res* 43: 1145-1152.
- Miller, R; Shah, R; Cattley, R; Popp, J. (1996). The peroxisome proliferations WY-14,643 and methyclofenapate induce hepatocyte ploidy alterations and ploidy-specific DNA synthesis in F344 rats. *Toxicol Appl Pharmacol* 138: 317-323.
- Mills, N and Means, A. (1972). Sorbitol dehydrogenase of rat testis: Changes of activity during development, after hypophysectomy and following gonadotrophic hormone administration. *Endocrinology* 91: 147-156.
- Minamishima, Y; Nakayama, K; Nakayama, K-I. (2002). Recovery of liver mass without proliferation of hepatocytes after partial hepatectomy in Skp2-deficient mice. *Cancer Res* 62: 995-999.
- Mirer, F. (2010). New evidence on the health hazards and control of metalworking fluids since completion of the OSHA advisory committee report. *Am J Ind Med* 53: 792-801. <http://dx.doi.org/10.1002/ajim.20853>.
- Mirsalis, J; Tyson, C; Steinmetz, K; Loh, E; Hamilton, C; Bakke, J; Spalding, J. (1989). Measurement of unscheduled DNA synthesis and S-phase synthesis in rodent hepatocytes following in vivo treatment: Testing of 24 compounds. *Environ Mol Mutagen* 14: 155-164. <http://dx.doi.org/10.1002/em.2850140305>.
- Mishima, N; Hoffman, S; Hill, E; Krug, E. (2006). Chick embryos exposed to trichloroethylene in an ex ovo culture model show selective defects in early endocardial cushion tissue formation. *Birth Defects Res A Clin Mol Teratol* 76: 517-527. <http://dx.doi.org/10.1002/bdra.20283>.
- Mitchell, A and Parsons-Smith, B. (1969). Trichloroethylene neuropathy. *Br Med J* 1: 422-423. <http://dx.doi.org/10.1136/bmj.1.5641.422>.
- Miura, Y; Anzai, K; Takahashi, S; Ozawa, T. (1997). A novel lipophilic spin probe for the measurement of radiation damage in mouse brain using in vivo electron spin resonance (ESR). *FEBS Lett* 419: 99-102. [http://dx.doi.org/10.1016/S0014-5793\(97\)01443-9](http://dx.doi.org/10.1016/S0014-5793(97)01443-9).
- Mjaatvedt, C; Lepera, R; Markwald, R. (1987). Myocardial specificity for initiating endothelial-mesenchymal cell transition in embryonic chick heart correlates with a particulate distribution of fibronectin. *Dev Biol* 119: 59-67.
- Mjaatvedt, C; Krug, E; Markwald, R. (1991). An antiserum (ES1) against a particulate form of extracellular matrix blocks the transition of cardiac endothelium into mesenchyme in culture. *Dev Biol* 145: 219-230.
- Moghaddam, A; Abbas, R; Fisher, J; Stavrou, S; Lipscomb, J. (1996). Formation of dichloroacetic acid by rat and mouse gut microflora, an in vitro study. *Biochem Biophys Res Commun* 228: 639-645. <http://dx.doi.org/10.1006/bbrc.1996.1709>.
- Moghaddam, A; Abbas, R; Fisher, J; Lipscomb, J. (1997). The role of mouse intestinal microflora in the metabolism of trichloroethylene, an in vivo study. *Hum Exp Toxicol* 16: 629-635. <http://dx.doi.org/10.1177/096032719701601101>.
- Moinzadeh, P; Breuhahn, K; Stützer, H; Schirmacher, P. (2005). Chromosome alterations in human hepatocellular carcinomas correlate with aetiology and histological grade: Results of an explorative CGH meta-analysis. *Br J Cancer* 92: 935-941. <http://dx.doi.org/10.1038/sj.bjc.6602448>.
- Monster, A; Boersma, G; Duba, W. (1976). Pharmacokinetics of trichloroethylene in volunteers, influence of workload and exposure concentration. *Int Arch Occup Environ Health* 38: 87-102. <http://dx.doi.org/10.1007/BF00378619>.

- Monster, A. (1979). Difference in uptake, elimination, and metabolism in exposure to trichloroethylene, 1,1,1-trichloroethane, and tetrachloroethylene. *Int Arch Occup Environ Health* 42: 311-317. <http://dx.doi.org/10.1007/BF00377785>.
- Monster, A; Boersma, G; Duba, W. (1979a). Kinetics of trichloroethylene in repeated exposure of volunteers. *Int Arch Occup Environ Health* 42: 283-292. <http://dx.doi.org/10.1007/BF00377782>.
- Monster, A; Boersma, G; Steenweg, H. (1979b). Kinetics of tetrachloroethylene in volunteers; influence of exposure concentration and work load. *Int Arch Occup Environ Health* 42: 303-309.
- Monster, A and Houtkooper, J. (1979). Estimation of individual uptake of trichloroethylene, 1,1,1-trichloroethane and tetrachloroethylene from biological parameters. *Int Arch Occup Environ Health* 42: 319-323. <http://dx.doi.org/10.1007/BF00377786>.
- Moore, L; Wilson, R; Campleman, S. (2005). Lifestyle factors, exposures, genetic susceptibility, and renal cell cancer risk: A review. *Cancer Invest* 23: 240-255. <http://dx.doi.org/10.1081/cnv-200055962>.
- Moore, L; Boffetta, P; Karami, S; Brennan, P; Stewart, P; Hung, R; Zaridze, D; Matveev, V; Janout, V; Kollarova, H; Bencko, V; Navratilova, M; Szeszenia-Dabrowska, N; Mates, D; Gromiec, J; Holcatova, I; Merino, M; Chanock, S; Chow, W; Rothman, N. (2010). Occupational trichloroethylene exposure and renal carcinoma risk: Evidence of genetic susceptibility by reductive metabolism gene variants. *Cancer Res* 70: 6527-6536. <http://dx.doi.org/10.1158/0008-5472.CAN-09-4167>.
- Moore, M and Harrington-Brock, K. (2000). Mutagenicity of trichloroethylene and its metabolites: Implications for the risk assessment of trichloroethylene. *Environ Health Perspect* 108: 215-223.
- Morata, T; Fiorini, A; Fischer, F; Colacioppo, S; Wallingford, K; Krieg, E; Dunn, D; Gozzoli, L; Padrão, M; Cesar, C. (1997). Toluene-induced hearing loss among rotogravure printing workers. *Scand J Work Environ Health* 23: 289-298.
- Morata, T; Johnson, A; Nylen, P; Svensson, E; Cheng, J; Krieg, E; Lindblad, A; Ernsgård, L; Franks, J. (2002). Audiometric findings in workers exposed to low levels of styrene and noise. *J Occup Environ Med* 44: 806-814. <http://dx.doi.org/10.1097/01.jom.0000026645.83602.60>.
- Morgan, J and Cassady, R. (2002). Community cancer assessment in response to long-time exposure to perchlorate and trichloroethylene in drinking water. *J Occup Environ Med* 44: 616-621.
- Morgan, R; Kelsh, M; Zhao, K; Heringer, S. (1998). Mortality of aerospace workers exposed to trichloroethylene. *Epidemiology* 9: 424-431.
- Morgan, R; Kelsh, M; Zhao, K; Heringer, S. (2000). Erratum: Mortality of aerospace workers exposed to trichloroethylene. *Epidemiology* 11: 360.
- Morgenstern, H; Froines, J; Ritz, B; Young, B. (1997). Epidemiologic study to determine possible adverse effects to Rocketdyne/Atomics International workers from exposure to ionizing radiation. Berkeley, CA: Public Health Institute. <http://www.mapcuzin.com/rocketdyne-cancer/radiation.pdf>.
- Morgenstern, H. (1998). Ecologic studies. In KJ Rothland & S Greenland (Eds.), *Modern epidemiology* (2nd ed., pp. 459-480). Philadelphia, PA: Lippincott, Williams, & Wilkins.
- Morgenstern, H; Froines, J; Ritz, B; Young, B. (1999). Epidemiologic study to determine possible adverse effects to Rocketdyne/Atomics International workers from exposure to selected chemicals. Berkeley, CA: Public Health Institute. http://www.ssflpanel.org/files/UCLA_chem.pdf.
- Morimura, K; Cheung, C; Ward, J; Reddy, J; Gonzalez, F. (2006). Differential susceptibility of mice humanized for peroxisome proliferator-activated receptor alpha to Wy-14,643-induced liver tumorigenesis. *Carcinogenesis* 27: 1074-1080. <http://dx.doi.org/10.1093/carcin/bgi329>.
- Morioka, I; Miyai, N; Yamamoto, H; Miyashita, K. (2000). Evaluation of combined effect of organic solvents and noise by the upper limit of hearing. *Ind Health* 38: 252-257. <http://dx.doi.org/10.2486/indhealth.38.252>.
- Moriya, M; Ohta, T; Watanabe, K; Miyazawa, T; Kato, K; Shirasu, Y. (1983). Further mutagenicity studies on pesticides in bacterial reversion assay systems. *Mutat Res* 116: 185-216.

- Morrow, L; Ryan, C; Goldstein, G; Hodgson, M. (1989). A distinct pattern of personality disturbance following exposure to mixtures of organic solvents. *J Occup Med* 31: 743-746.
- Morrow, L; Steinhauer, S; Hodgson, M. (1992). Delay in P300 latency in patients with organic solvent exposure. *Arch Neurol* 49: 315-320.
- Mortelmans, K; Haworth, S; Lawlor, T; Speck, W; Tainer, B; Zeiger, E. (1986). Salmonella mutagenicity tests: II. Results from the testing of 270 chemicals. *Environ Mutagen* 8: 1-119.
<http://dx.doi.org/10.1002/em.2860080702>.
- Morton, L; Youssef, A; Lloyd, E; Kiorpes, A; Goldsworthy, T; Fort, F. (2002). Evaluation of carcinogenic responses in the Eker rat following short-term exposure to selected nephrotoxins and carcinogens. *Toxicol Pathol* 30: 559-564.
- Morton, L; Holford, T; Leaderer, B; Zhang, Y; Zahm, S; Boyle, P; Flynn, S; Tallini, G; Owens, P; Zhang, B; Zheng, T. (2003). Alcohol use and risk of non-Hodgkin's lymphoma among Connecticut women (United States). *Cancer Causes Control* 14: 687-694.
- Morton, L; Hartge, P; Holford, T; Holly, E; Chiu, B; Vineis, P; Stagnaro, E; Willett, E; Franceschi, S; La Vecchia, C; Hughes, A; Cozen, W; Davis, S; Severson, R; Bernstein, L; Mayne, S; Dee, F; Cerhan, J; Zheng, T. (2005). Cigarette smoking and risk of non-Hodgkin lymphoma: A pooled analysis from the International Lymphoma Epidemiology Consortium (InterLymph). *Cancer Epidemiol Biomarkers Prev* 14: 925-933. <http://dx.doi.org/10.1158/1055-9965.EPI-04-0693>.
- Morton, L; Turner, J; Cerhan, J; Linet, M; Treseler, P; Clarke, C; Jack, A; Cozen, W; Maynadie, M; Spinelli, J; Costantini, A; Rudiger, T; Scarpa, A; Zheng, T; Weisenburger, D. (2007). Proposed classification of lymphoid neoplasms for epidemiologic research from the Pathology Working Group of the International Lymphoma Epidemiology Consortium (InterLymph). *Blood* 110: 695-708. <http://dx.doi.org/10.1182/blood-2006-11-051672>.
- Moser, V; Cheek, B; MacPhail, R. (1995). A multidisciplinary approach to toxicological screening: III. Neurobehavioral toxicity. *J Toxicol Environ Health A* 45: 173-210.
<http://dx.doi.org/10.1080/15287399509531988>.
- Moser, V; Phillips, P; McDaniel, K; MacPhail, R. (1999). Behavioral evaluation of the neurotoxicity produced by dichloroacetic acid in rats. *Neurotoxicol Teratol* 21: 719-731.
- Moser, V; MacPhail, R; Gennings, C. (2003). Neurobehavioral evaluations of mixtures of trichloroethylene, heptachlor, and di(2-ethylhexyl)phthalate in a full-factorial design. *Toxicology* 188: 125-137. [http://dx.doi.org/10.1016/S0300-483X\(03\)00083-0](http://dx.doi.org/10.1016/S0300-483X(03)00083-0).
- Moslen, M; Reynolds, E; Boor, P; Bailey, K; Szabo, S. (1977). Trichloroethylene-induced deactivation of cytochrome P-450 and loss of liver glutathione in vivo. *Res Commun Mol Pathol Pharmacol* 16: 109-120.
- Motohashi, Y and Miyazaki, Y. (1990). Time-dependent effects of trichloroethylene on motor activity in rats. *Chronobiol Int* 7: 193-201. <http://dx.doi.org/10.3109/07420529009056975>.
- Muijser, H; Lammers, J; Kullig, B. (2000). Effects of exposure to trichloroethylene and noise on hearing in rats. *Noise Health* 2: 57-66.
- Muller, G; Spassovski, M; Henschler, D. (1972). Trichloroethylene exposure and trichloroethylene metabolites in urine and blood. *Arch Toxicol* 29: 335-340.
<http://dx.doi.org/10.1007/BF00326650>.
- Muller, G; Spassovski, M; Henschler, D. (1974). Metabolism of trichloroethylene in man: II. Pharmacokinetics of metabolites. *Arch Toxicol* 32: 283-295.
<http://dx.doi.org/10.1007/BF00330110>.
- Muller, G; Spassowski, M; Henschler, D. (1975). Metabolism of trichloroethylene in man: III. Interaction of trichloroethylene and ethanol. *Arch Toxicol* 33: 173-189.
<http://dx.doi.org/10.1007/BF00311271>.
- Müller, W; Coulston, F; Korte, F. (1982). Comparative metabolism of 14C-trichloroethylene in chimpanzees, baboons, and rhesus monkeys. *Chemosphere* 11: 215-218.
[http://dx.doi.org/10.1016/0045-6535\(82\)90169-2](http://dx.doi.org/10.1016/0045-6535(82)90169-2).

- Murphy, J. (1990). Exposure to Wells G and H in Woburn, Massachusetts. Boston, MA: Massachusetts Health Research Institute, Inc. and Massachusetts Department of Public Health.
http://www.mass.gov/Eohhs2/docs/dph/environmental/investigations/woburn_exposure_wells.pdf.
- Nagaya, T; Ishikawa, N; Hata, H. (1989a). Urinary total protein and "beta"-2-microglobulin in workers exposed to trichloroethylene. Environ Res 50: 86-92.
- Nagaya, T; Ishikawa, N; Hata, H. (1989b). Sister-chromatid exchanges in lymphocytes of workers exposed to trichloroethylene. Mutat Res 222: 279-282.
- Nagaya, T; Ishikawa, N; Hata, H. (1990). No change in serum dopamine-beta-hydroxylase activity in workers exposed to trichloroethylene. Toxicol Lett 54: 221-227. [http://dx.doi.org/10.1016/0378-4274\(90\)90187-Q](http://dx.doi.org/10.1016/0378-4274(90)90187-Q).
- Nagaya, T; Ishikawa, N; Hata, H; Otobe, T. (1993). Subclinical and reversible hepatic effects of occupational exposure to trichloroethylene. Int Arch Occup Environ Health 64: 561-563. <http://dx.doi.org/10.1007/BF00517701>.
- Nakai, J; Stathopoulos, P; Campbell, G; Chu, I; Li-Muller, A; Aucoin, R. (1999). Penetration of chloroform, trichloroethylene, and tetrachloroethylene through human skin. J Toxicol Environ Health A 58: 157-170. <http://dx.doi.org/10.1080/009841099157368>.
- Nakajima, T; Koyama, Y; Sato, A. (1982). Dietary modification of metabolism and toxicity of chemical substances - with special reference to carbohydrate. Biochem Pharmacol 31: 1005-1011.
- Nakajima, T; Okino, T; Okuyama, S; Kaneko, T; Yonekura, I; Sato, A. (1988). Ethanol-induced enhancement of trichloroethylene metabolism and hepatotoxicity: Difference from the effect of phenobarbital. Toxicol Appl Pharmacol 94: 227-237. [http://dx.doi.org/10.1016/0041-008X\(88\)90264-5](http://dx.doi.org/10.1016/0041-008X(88)90264-5).
- Nakajima, T; Wang, R-S; Murayama, N; Sato, A. (1990). Three forms of trichloroethylene-metabolizing enzymes in rat liver induced by ethanol, phenobarbital, and 3-methylcholanthrene. Toxicol Appl Pharmacol 102: 546-552. [http://dx.doi.org/10.1016/0041-008X\(90\)90049-Z](http://dx.doi.org/10.1016/0041-008X(90)90049-Z).
- Nakajima, T; Wang, R; Elovaara, E; Park, S; Gelboin, H; Vainio, H. (1992a). A comparative study on the contribution of cytochrome P450 isozymes to metabolism of benzene, toluene and trichloroethylene in rat liver. Biochem Pharmacol 43: 251-257.
- Nakajima, T; Wang, R; Katakura, Y; Kishi, R; Elovaara, E; Park, S; Gelboin, H; Vainio, H. (1992b). Sex-, age- and pregnancy-induced changes in the metabolism of toluene and trichloroethylene in rat liver in relation to the regulation of cytochrome P450IIE1 and P450IIC11 content. J Pharmacol Exp Ther 261: 869-874.
- Nakajima, T; Wang, R-S; Elovaara, E; Park, S; Gelboin, H; Vainio, H. (1993). Cytochrome P450-related differences between rats and mice in the metabolism of benzene, toluene and trichloroethylene in liver microsomes. Biochem Pharmacol 45: 1079-1085. [http://dx.doi.org/10.1016/0006-2952\(93\)90252-R](http://dx.doi.org/10.1016/0006-2952(93)90252-R).
- Nakajima, T and Aoyama, T. (2000). Polymorphism of drug-metabolizing enzymes in relation to individual susceptibility to industrial chemicals. Ind Health 38: 143-152.
- Nakajima, T; Kamijo, Y; Usuda, N; Liang, Y; Fukushima, Y; Kametani, K; Gonzalez, F; Aoyama, T. (2000). Sex-dependent regulation of hepatic peroxisome proliferation in mice by trichloroethylene via peroxisome proliferator-activated receptor alpha (PPARalpha). Carcinogenesis 21: 677-682.
- Nakajima, T; Yamanoshita, O; Kamijima, M; Kishi, R; Ichihara, G. (2003). Generalized skin reactions in relation to trichloroethylene exposure: a review from the viewpoint of drug-metabolizing enzymes. J Occup Health 45: 8-14.
- Nakajima, T; Moriguchi, M; Mitsumoto, Y; Sekoguchi, S; Nishikawa, T; Takashima, H; Watanabe, T; Katagishi, T; Kimura, H; Okanoue, T; Kagawa, K. (2004). Centrosome aberration accompanied with p53 mutation can induce genetic instability in hepatocellular carcinoma. Mod Pathol 17: 722-727. <http://dx.doi.org/10.1038/modpathol.3800115>.

- Nakajima, Y; Mironov, V; Yamagishi, T; Nakamura, H; Markwald, R. (1997). Expression of smooth muscle alpha-actin in mesenchymal cells during formation of avian endocardial cushion tissue: a role for transforming growth factor beta3. *Dev Dyn* 209: 296-309.
[http://dx.doi.org/10.1002/\(SICI\)1097-0177\(199707\)209:3<296::AID-AJA5>3.0.CO;2-D](http://dx.doi.org/10.1002/(SICI)1097-0177(199707)209:3<296::AID-AJA5>3.0.CO;2-D).
- Nakanishi, S; Shiohara, E; Tsukada, M; Yamazaki, Y; Okumura, K. (1978). Acetaldehyde level in the blood and liver aldehyde dehydrogenase activities in trichloroethylene-treated rats. *Arch Toxicol* 41: 207-214.
- Narotsky, M. (2008). personal communications from Michael Narotsky, U.S. EPA, to John Fox, U.S. EPA, 19 June 2008, and to Jennifer Jinot, U.S. EPA, 10 June 2008
- Narotsky, M and Kavlock, R. (1995). A multidisciplinary approach to toxicological screening: II. Developmental toxicity. *J Toxicol Environ Health* 45: 145-171.
<http://dx.doi.org/10.1080/15287399509531987>.
- Narotsky, M; Weller, E; Chinchilli, V; Kavlock, R. (1995). Nonadditive developmental toxicity in mixtures of trichloroethylene, di(2-ethylhexyl) phthalate, and heptachlor in a 5 x 5 x 5 design. *Fundam Appl Toxicol* 27: 203-216. <http://dx.doi.org/10.1093/toxsci/27.2.203>.
- Natarajan, A; Duivenvoorden, W; Meijers, M; Zwanenburg, T. (1993). Induction of mitotic aneuploidy using Chinese hamster primary embryonic cells. Test results of 10 chemicals. *Mutat Res* 287: 47-56. [http://dx.doi.org/10.1016/0027-5107\(93\)90144-5](http://dx.doi.org/10.1016/0027-5107(93)90144-5).
- Naugler, W; Sakurai, T; Kim, S; Maeda, S; Kim, K; Elsharkawy, A; Karin, M. (2007). Gender disparity in liver cancer due to sex differences in MyD88-dependent IL-6 production. *Science* 317: 121-124. <http://dx.doi.org/10.1126/science.1140485>.
- NCI. (National Institutes of Health, National Cancer Institute). (1976). Carcinogenesis bioassay of trichloroethylene. (NCI-CG-TR-2). In Technical Report Series. Bethesda, MD: U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health.
http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr002.pdf.
- Neghab, M; Qu, S; Bai, C; Caples, J; Stacey, N. (1997). Raised concentration of serum bile acids following occupational exposure to halogenated solvents, 1,1,2-trichloro-1,2,2-trifluoroethane and trichloroethylene. *Int Arch Occup Environ Health* 70: 187-194.
<http://dx.doi.org/10.1007/s004200050205>.
- Neghab, M and Stacey, N. (1997). Toluene-induced elevation of serum bile acids: Relationship to bile acid transport. *J Toxicol Environ Health* 52: 249-268.
- Nelson, G; Swank, A; Brooks, L; Bailey, K; George, S. (2001). Metabolism, microflora effects, and genotoxicity in haloacetic acid-treated cultures of rat cecal microbiota. *Toxicol Sci* 60: 232-241. <http://dx.doi.org/10.1093/toxsci/60.2.232>.
- Nelson, H; Wiencke, J; Christiani, D; Cheng, T; Zuo, Z-F; Schwartz, B; K, LB-; Spitz, M; Wang, M; Xu, X; Kelsey, K. (1995). Ethnic differences in the prevalence of the homozygous deleted genotype of glutathione S-transferase theta. *Carcinogenesis* 16: 1243-1245.
- Nelson, J and Zenick, H. (1986). The effect of trichloroethylene on male sexual behavior: possible opioid role. *Neurobehav Toxicol Teratol* 8: 441-445.
- Nelson, L; Longstreth, W; Koepsell, T; Checkoway, H; van Belle, G. (1994). Completeness and accuracy of interview data from proxy respondents: Demographic, medical, and life-style factors. *Epidemiology* 5: 204-217.
- Nelson, M and Bull, R. (1988). Induction of strand breaks in DNA by trichloroethylene and metabolites in rat and mouse liver in vivo. *Toxicol Appl Pharmacol* 94: 45-54.
[http://dx.doi.org/10.1016/0041-008X\(88\)90335-3](http://dx.doi.org/10.1016/0041-008X(88)90335-3).
- Nelson, M; Lansing, A; Sanchez, I; Bull, R; Springer, D. (1989). Dichloroacetic acid and trichloroacetic acid-induced DNA strand breaks are independent of peroxisome proliferation. *Toxicology* 58: 239-248. [http://dx.doi.org/10.1016/0300-483X\(89\)90139-X](http://dx.doi.org/10.1016/0300-483X(89)90139-X).
- Nelson, M; Sanchez, I; Bull, R; Sylvester, S. (1990). Increased expression of c-myc and c-H-ras in dichloroacetate and trichloroacetate-induced liver tumors in B6C3F1 mice. *Toxicology* 64: 47-57.
[http://dx.doi.org/10.1016/0300-483X\(90\)90098-2](http://dx.doi.org/10.1016/0300-483X(90)90098-2).

- Nesslany, F and Marzin, D. (1999). A micromethod for the in vitro micronucleus assay. Mutagenesis 14: 403-410. <http://dx.doi.org/10.1093/mutage/14.4.403>.
- Nestmann, E; Chu, I; Kowbel, D; Matula, T. (1980). Short-lived mutagen in Salmonella produced by reaction of trichloroacetic acid and dimethyl sulphoxide. Can J Genet Cytol 22: 35-40. <http://dx.doi.org/10.1139/g80-006>.
- Ni, Y-C; Kadlubar, F; Fu, P. (1995). Formation of malondialdehyde-modified 2'-deoxyguanosinyl adduct from metabolism of chloral hydrate by mouse liver microsomes. Biochem Biophys Res Commun 216: 1110-1117. <http://dx.doi.org/10.1006/bbrc.1995.2735>.
- Ni, Y; Wong, T; Kadlubar, F; Fu, P. (1994). Hepatic metabolism of chloral hydrate to free radical(s) and induction of lipid peroxidation. Biochem Biophys Res Commun 204: 937-943. <http://dx.doi.org/10.1006/bbrc.1994.2550>.
- Ni, Y; Wong, T; Lloyd, R; Heinze, T; Shelton, S; Casciano, D; Kadlubar, F; Fu, P. (1996). Mouse liver microsomal metabolism of chloral hydrate, trichloroacetic acid, and trichloroethanol leading to induction of lipid peroxidation via a free radical mechanism. Drug Metab Dispos 24: 81-90.
- Nickerson, M; Jaeger, E; Shi, Y; Durocher, J; Mahurkar, S; Zaridze, D; Matveev, V; Janout, V; Kollarova, H; Bencko, V; Navratilova, M; Szeszenia-Dabrowska, N; Mates, D; Mukeria, A; Holcatova, I; Schmidt, L; Toro, J; Karami, S; Hung, R; Gerard, G; Linehan, W; Merino, M; Zbar, B; Boffetta, P; Brennan, P; Rothman, N; Chow, W; Waldman, F; Moore, L. (2008). Improved identification of von Hippel-Lindau gene alterations in clear cell renal tumors. Clin Cancer Res 14: 4726-4734. <http://dx.doi.org/10.1158/1078-0432.CCR-07-4921>.
- Nickoloff, B; Ben-Neriah, Y; Pikarsky, E. (2005). Inflammation and cancer: Is the link as simple as we think? J Invest Dermatol 124: x-xiv. <http://dx.doi.org/10.1111/j.0022-202X.2005.23724.x>.
- Niederlechner, B; Cairns, J; Smith, E. (1998). Modeling acute and chronic toxicity of nonpolar narcotic chemicals and mixtures to Ceriodaphnia dubia. Ecotoxicol Environ Saf 39: 136-146. <http://dx.doi.org/10.1006/eesa.1997.1621>.
- NIEHS. (National Institutes of Health, National Institute for Environmental Health Sciences). (1998). Tumor incidence in control animals by route and vehicle of administration, B6C3F1 mice. Research Triangle Park, NC.
- Nietert, P; Sutherland, S; Silver, R; Pandey, J; Knapp, R; Hoel, D; Dosemeci, M. (1998). Is occupational organic solvent exposure a risk factor for scleroderma? Arthritis Rheum 41: 1111-1118. [http://dx.doi.org/10.1002/1529-0131\(199806\)41:6<1111::AID-ART19>3.0.CO;2-J](http://dx.doi.org/10.1002/1529-0131(199806)41:6<1111::AID-ART19>3.0.CO;2-J).
- Niklasson, M; Tham, R; Larsby, B; Eriksson, B. (1993). Effects of toluene, styrene, trichloroethylene, and trichloroethane on the vestibulo- and opto-oculo motor system in rats. Neurotoxicol Teratol 15: 327-334.
- NIOSH. (National Institute for Occupational Safety and Health). (1992). Health hazard evaluation report, HETA 86-469-2189, James River Corporation, Newnan, Georgia. (HETA 86-469-2189). pp. 24. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health. <http://www.cdc.gov/niosh/hhe/reports/pdfs/1986-0469-2189.pdf>.
- NIOSH. (National Institute for Occupational Safety and Health). (1998). Criteria for a recommended standard: Occupational exposure to metalworking fluids. (98-102). pp. 242. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. <http://www.cdc.gov/niosh/98-102.html>.
- Nissen, S; Nicholls, S; Wolski, K; Howey, D; McErlean, E; Wang, M-D; Gomez, E; Russo, J. (2007). Effects of a potent and selective PPAR-alpha agonist in patients with atherogenic dyslipidemia or hypercholesterolemia: Two randomized controlled trials. JAMA 297: 1362-1373. <http://dx.doi.org/10.1001/jama.297.12.1362>.
- Nitcheva, D; Piegorsch, W; West, R. (2007). On use of the multistage dose-response model for assessing laboratory animal carcinogenicity. Regul Toxicol Pharmacol 48: 135-147. <http://dx.doi.org/10.1016/j.yrtph.2007.03.002>.

- Nöhammer, C; Brunner, F; Wölkart, G; Staber, P; Steyrer, E; Gonzalez, F; Zechner, R; Hoefler, G. (2003). Myocardial dysfunction and male mortality in peroxisome proliferator-activated receptor alpha knockout mice overexpressing lipoprotein lipase in muscle. *Lab Invest* 83: 259-269.
<http://dx.doi.org/10.1097/01.LAB.0000053916.61772.CA>.
- Noland-Gerbec, E; Pfohl, R; Taylor, D; Bull, R. (1986). 2-Deoxyglucose uptake in the developing rat brain upon pre- and postnatal exposure to trichloroethylene. *Neurotoxicology* 7: 157-164.
- Nomiyama, H and Nomiyama, K. (1979). Pathway and rate of metabolism of trichloroethylene in rats and rabbits. *Ind Health* 17: 29-37.
- Nomiyama, K and Nomiyama, H. (1971). Metabolism of trichloroethylene in human: Sex difference in urinary excretion of trichloroacetic acid and trichloroethanol. *Int Arch Occup Environ Health* 28: 37-48. <http://dx.doi.org/10.1007/BF00539728>.
- Nordlund, L; Carstensen, J; Pershagen, G. (1997). Cancer incidence in female smokers: A 26-year follow-up. *Int J Cancer* 73: 625-628. [http://dx.doi.org/10.1002/\(SICI\)1097-0215\(19971127\)73:5<625::AID-IJC2>3.0.CO;2-Z](http://dx.doi.org/10.1002/(SICI)1097-0215(19971127)73:5<625::AID-IJC2>3.0.CO;2-Z).
- Nordlund, L; Carstensen, J; Pershagen, G. (1999). Are male and female smokers at equal risk of smoking-related cancer: Evidence from a Swedish prospective study. *Scand J Public Health* 27: 56-62.
- Nordström, M; Hardell, L; Magnuson, A; Hagberg, H; Rask-Andersen, A. (1998). Occupational exposures, animal exposure and smoking as risk factors for hairy cell leukaemia evaluated in a case-control study. *Br J Cancer* 77: 2048-2052.
- NRC. (National Research Council). (1983). Risk assessment in the federal government: Managing the process. Washington, DC: National Academies Press.
http://www.nap.edu/openbook.php?record_id=366&page=R1.
- NRC. (National Research Council). (1993). Pesticides in the diets of infants and children. Washington, DC: National Academy Press.
- NRC. (National Research Council). (1994). Science and judgment in risk assessment. Washington, DC: National Academy Press.
- NRC. (National Research Council). (1997). Environmental epidemiology, volume 2: Use of the gray literature and other data in environmental epidemiology (Vol. 2). Washington, DC: National Academies Press.
- NRC. (National Research Council). (2006). Assessing the human health risks of trichloroethylene: Key scientific issues. Washington, DC: The National Academies Press.
<http://nae.edu/nae/naepcms.nsf/weblinks/MKEZ-6SSHPD?OpenDocument>.
- NRC. (National Research Council). (2009). Contaminated water supplies at Camp Lejeune: Assessing potential health effects. Washington, DC: National Academies Press.
- NTP. (National Toxicology Program). (1988). Toxicology and carcinogenesis studies of trichloroethylene (CAS No. 79-01-6) in four strains of rats (ACI, August, Marshall, Osborne-Mendel) (gavage studies). Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health.
http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr273.pdf.
- NTP. (National Toxicology Program). (1990). Carcinogenesis studies of trichloroethylene (without epichlorohydrin) (CAS No. 79-01-6) in F344/N rats and B6C3F1 mice (gavage studies). In Technical Report Series. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health.
http://ntp.niehs.nih.gov/ntp/htdocs/LT_rpts/tr243.pdf.
- Nunes, J; Ehrlich, M; Robertson, J. (2001). Toxicosis associated with dual oral exposure of rats to lead and trichloroethylene. *Toxicol Pathol* 29: 451-457.
- Nutley, E; Tcheong, A; Allen, J; Collins, B; Ma, M; Lowe, X; Bishop, J; Moore, D; Wyrobek, A. (1996). Micronuclei induced in round spermatids of mice after stem-cell treatment with chloral hydrate: evaluations with centromeric DNA probes and kinetochore antibodies. *Environ Mol Mutagen* 28: 80-89. [http://dx.doi.org/10.1002/\(SICI\)1098-2280\(1996\)28:2<80::AID-EM3>3.0.CO;2-I](http://dx.doi.org/10.1002/(SICI)1098-2280(1996)28:2<80::AID-EM3>3.0.CO;2-I).

- NYSDEC. (New York State Department of Environmental Conservation). (2008). Project information summary: Solvents, contamination and cleanup at Endicott. pp. 4. Albany, NY. <http://www.dec.ny.gov/chemical/24887.html>.
- NYSDEC. (New York State Department of Environmental Conservation). (2007). Field investigation report for groundwater and residential air sampling August 2006 through March 2007 soil vapor intrusion investigation, SCM Cortlandville site, Site #7-12-006, Cortlandville, NY. pp. 186. Albany, NY.
- NYSDOH. (New York State Department of Health). (2005). Trichloroethene air criteria document. Draft report. Troy, NY: New York State Department of Health, Center for Environmental Health.
- NYSDOH. (New York State Department of Health). (2006). Trichloroethene air criteria document: Final report. pp. 413. Troy, NY: New York State Department of Health, Center for Environmental Health, Bureau of Toxic Substance Assessment. http://www.health.ny.gov/environmental/chemicals/trichloroethylene/docs/cd_tce.pdf.
- O'Donnell, L; McLachlan, R; Wreford, N; Robertson, D. (1994). Testosterone promotes the conversion of round spermatids between stages VII and VIII of the rat spermatogenic cycle. *Endocrinology* 135: 2608-2614.
- Odum, J; Foster, J; Green, T. (1992). A mechanism for the development of Clara cell lesions in the mouse lung after exposure to trichloroethylene. *Chem Biol Interact* 83: 135-153.
- Ogata, M; Takatsuka, Y; Tomokuni, K. (1971). Excretion of organic chlorine compounds in the urine of persons exposed to vapours of trichloroethylene and tetrachloroethylene. *Occup Environ Med* 28: 386-391. <http://dx.doi.org/10.1136/oem.28.4.386>.
- Ohta, M; Saito, T; Saito, K; Kurasaki, M; Hosokawa, T. (2001). Effect of trichloroethylene on spatiotemporal pattern of LTP in mouse hippocampal slices. *Int J Neurosci* 111: 257-271.
- Okino, M; Chiu, W; Evans, M; Power, F; Lipscomb, J; Tornero-Velez, R; Dary, C; Blancato, J; Chen, C. (2005). Suitability of using in vitro and computationally estimated parameters in simplified pharmacokinetic models. *Drug Metab Rev* 37: 162.
- Okino, T; Nakajima, T; Nakano, M. (1991). Morphological and biochemical analyses of trichloroethylene hepatotoxicity: Differences in ethanol- and phenobarbital-pretreated rats. *Toxicol Appl Pharmacol* 108: 379-389. [http://dx.doi.org/10.1016/0041-008X\(91\)90084-R](http://dx.doi.org/10.1016/0041-008X(91)90084-R).
- Olivier, P; Plancke, M; Marzin, D; Clavey, V; Sauzieres, J; Fruchart, J. (1988). Effects of fenofibrate, gemfibrozil and nicotinic acid on plasma lipoprotein levels in normal and hyperlipidemic mice: A proposed model for drug screening. *Atherosclerosis* 70: 107-114. [http://dx.doi.org/10.1016/0021-9150\(88\)90104-9](http://dx.doi.org/10.1016/0021-9150(88)90104-9).
- Olshan, A; De Roos, A; Teschke, K; Neglia, J; Stram, D; Pollock, B; Castleberry, R. (1999). Neuroblastoma and parental occupation. *Cancer Causes Control* 10: 539-549. <http://dx.doi.org/10.1023/A:1008998925889>.
- Olsson, A; Skogh, T; Wingren, G. (2000). Occupational determinants for rheumatoid arthritis. *Scand J Work Environ Health* 26: 243-249.
- Olsson, A; Skogh, T; Axelson, O; Wingren, G. (2004). Occupations and exposures in the work environment as determinants for rheumatoid arthritis. *Occup Environ Med* 61: 233-238.
- Oneta, C; Lieber, C; Li, J; Rüttimann, S; Schmid, B; Lattmann, J; Rosman, A; Seitz, H. (2002). Dynamics of cytochrome P4502E1 activity in man: Induction by ethanol and disappearance during withdrawal phase. *J Hepatol* 36: 47-52. [http://dx.doi.org/10.1016/S0168-8278\(01\)00223-9](http://dx.doi.org/10.1016/S0168-8278(01)00223-9).
- Ono, Y; Somiya, I; Kawamura, M. (1991). The evaluation of genotoxicity using DNA repairing test for chemicals produced in chlorination and ozonation processes. *Water Sci Technol* 23: 329-338.
- Opdam, J. (1989). Intra and interindividual variability in the kinetics of a poorly and highly metabolising solvent. *Br J Ind Med* 46: 831-845.
- Orsi, L; Monnereau, A; Dananche, B; Berthou, C; Fenaux, P; Marit, G; Soubeyran, P; Huguet, F; Mil pied, N; Leporrier, M; Hemon, D; Troussard, X; Clavel, J. (2010). Occupational exposure to organic solvents and lymphoid neoplasms in men: Results of a French case-control study. *Occup Environ Med* 67: 664-672. <http://dx.doi.org/10.1136/oem.2009.049460>.

- Oshiro, W; Krantz, Q; Bushnell, P. (2004). A search for residual behavioral effects of trichloroethylene in rats exposed as young adults. Neurotoxicol Teratol 26: 239-251.
<http://dx.doi.org/10.1016/j.ntt.2003.12.005>.
- Ou, J; Ou, Z; McCarver, D; Hines, R; Oldham, K; Ackerman, A; Pritchard, K. (2003). Trichloroethylene decreases heat shock protein 90 interactions with endothelial nitric oxide synthase: implications for endothelial cell proliferation. Toxicol Sci 73: 90-97. <http://dx.doi.org/10.1093/toxsci/kfg062>.
- Owens, A and Marshall, E. (1955). Further studies on the metabolic rate of chloral hydrate and trichloroethanol. Johns Hopkins Med J 97: 320-326.
- Palmer, C; Hsu, M; Griffin, K; Raucy, J; Johnson, E. (1998). Peroxisome proliferator activated receptor-alpha expression in human liver. Mol Pharmacol 53: 14-22.
- Pannett, B; Coggon, D; Acheson, E. (1985). A job-exposure matrix for use in population based studies in England and Wales. Br J Ind Med 42: 777-783.
- Parent, M; Hua, Y; Siemiatycki, J. (2000a). Occupational risk factors for renal cell carcinoma in Montreal. Am J Ind Med 38: 609-618.
- Parent, M; Siemiatycki, J; Fritsch, L. (2000b). Workplace exposures and oesophageal cancer. Occup Environ Med 57: 325-334.
- Park, P; Gilmour, M; Selgrade, M. (1993). Pulmonary defenses to streptococcal infection following acute exposure to trichloroethylene (TCE) or chloroform (CHCl₃). Toxicologist 13: 107.
- Park, S; Osterloh, J; Vamvakas, S; Hashmi, M; Anders, M; Cashman, J. (1992). Flavin-containing monooxygenase-dependent stereoselective S-oxygénéation and cytotoxicity of cysteine S-conjugates and mercapturates. Chem Res Toxicol 5: 193-201.
- Parker, M; Tarlow, M; Milne Anderson, J. (1984). Glue sniffing and cerebral infarction. Arch Dis Child 59: 675-677.
- Parkin, DM; Whelan, SL; Ferlay, J; Teppo, L; Thomas, DB (Eds.). (2002). Cancer incidence in five continents: Volume VIII (Vol. 8). Lyon, France: International Agency for Research on Cancer.
- Parkinson, A; Mudra, D; Johnson, C; Dwyer, A; Carroll, K. (2004). The effects of gender, age, ethnicity, and liver cirrhosis on cytochrome P450 enzyme activity in human liver microsomes and inducibility in cultured human hepatocytes. Toxicol Appl Pharmacol 199: 193-209.
<http://dx.doi.org/10.1016/j.taap.2004.01.010>.
- Parnell, M; Koller, L; Exon, J; Arnzen, J. (1986). Trichloroacetic acid effects on rat liver peroxisomes and enzyme-altered foci. Environ Health Perspect 69: 73-79.
- Parrish, J; Austin, E; Stevens, D; Kinder, D; Bull, R. (1996). Haloacetate-induced oxidative damage to DNA in the liver of male B6C3F1 mice. Toxicology 110: 103-111.
[http://dx.doi.org/10.1016/0300-483X\(96\)03342-2](http://dx.doi.org/10.1016/0300-483X(96)03342-2).
- Parry, E; Hague, A; Parry, J. (1990). A study of mitotic division fidelity and numerical chromosome changes in ageing Syrian hamster dermal cells. Mutat Res 237: 83-93.
[http://dx.doi.org/10.1016/0921-8734\(90\)90014-I](http://dx.doi.org/10.1016/0921-8734(90)90014-I).
- Parry, J; Parry, E; Bourner, R; Doherty, A; Ellard, S; O'Donovan, J; Hoebee, B; de Stoppelaar, J; Mohn, G; Onfelt, A; Renglin, A; Schultz, N; Soderpalm-Berndes, C; Jensen, K; Kirsch-Volders, M; Elhajouji, A; Van Hummelen, P; Degrassi, F; Antoccia, A; Cimini, D; Izzo, M; Tanzarella, C; Adler, I; Kliesch, U; Schriever-Schwemmer, G; Gasser, P; Crebelli, R; Carere, A; Andreoli, C; Benigni, R; Leopardi, P; Marcon, F; Zinjo, Z; Natarajan, A; Boei, J; Kappas, A; Voutsinas, G; Zarani, F; Patrinelli, A; Pachierotti, F; Tiveron, C; Hess, P. (1996). The detection and evaluation of aneugenic chemicals. Mutat Res 353: 11-46. [http://dx.doi.org/10.1016/0027-5107\(95\)00242-1](http://dx.doi.org/10.1016/0027-5107(95)00242-1).
- Partanen, T; Heikkila, P; Hernberg, S; Kauppinen, T; Moneta, G; Ojajarvi, A. (1991). Renal cell cancer and occupational exposure to chemical agents. Scand J Work Environ Health 17: 231-239.
- Pastino, G; Yap, W; Carroquino, M. (2000). Human variability and susceptibility to trichloroethylene. Environ Health Perspect 108: 201-214.
- Paykoc, Z and Powell, J. (1945). The excretion of sodium trichloroacetate. J Pharmacol Exp Ther 85: 289-293.

- Peden-Adams. (2008). personal communication from Margie Peden-Adams, Medical University of South Carolina, to Jennifer Jinot, EPA, 26 August 2008.
- Peden-Adams, M; Eudaly, J; Heesemann, L; Smythe, J; Miller, J; Gilkeson, G; Keil, D. (2006). Developmental immunotoxicity of trichloroethylene (TCE): Studies in B6C3F1 mice. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 41: 249-271.
<http://dx.doi.org/10.1080/10934520500455289>.
- Peden-Adams, M; Eudaly, J; Lee, A; Miller, J; Keil, D; Gilkeson, G. (2008). Lifetime exposure to trichloroethylene (TCE) does not accelerate autoimmune disease in MRL +/+ mice. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 43: 1402-1409.
- Pelekis, M; Gephart, L; Lerman, S. (2001). Physiological-model-based derivation of the adult and child pharmacokinetic intraspecies uncertainty factors for volatile organic compounds. *Regul Toxicol Pharmacol* 33: 12-20. <http://dx.doi.org/10.1006/rtph.2000.1436>.
- Pellizzari, E; Hartwell, T; Harris, B, III; Waddell, R; Whitaker, D; Erickson, M. (1982). Purgeable organic compounds in mother's milk. *Bull Environ Contam Toxicol* 28: 322-328.
- Perbellini, L; Olivato, D; Zedde, A; Miglioranzi, R. (1991). Acute trichloroethylene poisoning by ingestion: Clinical and pharmacokinetic aspects. *Intensive Care Med* 17: 234-235.
<http://dx.doi.org/10.1007/BF01709883>.
- Percy, C; III, SE; Gloeckler, L. (1981). Accuracy of cancer death certificates and its effect on cancer mortality statistics. *Am J Public Health* 71: 242-250.
- Percy, C; Ries, L; Van Holten, V. (1990). The accuracy of liver cancer as the underlying cause of death on death certificates. *Public Health Rep* 105: 361-367.
- Pereira, M. (1996). Carcinogenic activity of dichloroacetic acid and trichloroacetic acid in the liver of female B6C3F1 mice. *Fundam Appl Toxicol* 31: 192-199.
<http://dx.doi.org/10.1006/faat.1996.0091>.
- Pereira, M and Phelps, J. (1996). Promotion by dichloroacetic acid and trichloroacetic acid of N-methyl-N-nitrosourea-initiated cancer in the liver of female B6C3F1 mice. *Cancer Lett* 102: 133-141.
[http://dx.doi.org/10.1016/0304-3835\(96\)04156-0](http://dx.doi.org/10.1016/0304-3835(96)04156-0).
- Pereira, M; Li, K; Kramer, P. (1997). Promotion by mixtures of dichloroacetic acid and trichloroacetic acid of N-methyl-N-nitrosourea-initiated cancer in the liver of female B6C3F1 mice. *Cancer Lett* 115: 15-23. [http://dx.doi.org/10.1016/S0304-3835\(97\)04699-5](http://dx.doi.org/10.1016/S0304-3835(97)04699-5).
- Pereira, M; Kramer, P; Conran, P; Tao, L. (2001). Effect of chloroform on dichloroacetic acid and trichloroacetic acid-induced hypomethylation and expression of the c-myc gene and on their promotion of liver and kidney tumors in mice. *Carcinogenesis* 22: 1511-1519.
- Pereira, M; Wang, W; Kramer, P; Tao, L. (2004a). Prevention by methionine of dichloroacetic acid-induced liver cancer and DNA hypomethylation in mice. *Toxicol Sci* 77: 243-248.
<http://dx.doi.org/10.1093/toxsci/kfh031>.
- Pereira, M; Wang, W; Kramer, P; Tao, L. (2004b). DNA hypomethylation induced by non-genotoxic carcinogens in mouse and rat colon. *Cancer Lett* 212: 145-151.
<http://dx.doi.org/10.1016/j.canlet.2004.03.024>.
- Perkins, N. (2004). NF-kB: tumor promoter or suppressor? *Trends Cell Biol* 14: 64-69.
<http://dx.doi.org/10.1016/j.tcb.2003.12.004>.
- Perkins, N and Gilmore, T. (2006). Good cop, bad cop: The different faces of NF-kB. *Cell Death Differ* 13: 759-772. <http://dx.doi.org/10.1038/sj.cdd.4401838>.
- Perocco, P and Prodi, G. (1981). DNA damage by haloalkanes in human lymphocytes cultured in vitro. *Cancer Lett* 13: 213-218. [http://dx.doi.org/10.1016/0304-3835\(81\)90020-3](http://dx.doi.org/10.1016/0304-3835(81)90020-3).
- Perrot, N; Nalpas, B; Yang, C; Beaune, P. (1989). Modulation of cytochrome P450 isozymes in human liver, by ethanol and drug intake. *Eur J Clin Invest* 19: 549-555. <http://dx.doi.org/10.1111/j.1365-2362.1989.tb00273.x>.
- Perry, S; Schofield, M; MacFarlane, M; Lock, E; King, L; Gibson, G; Goldfarb, P. (1993). Isolation and expression of a cDNA coding for rat kidney cytosolic cysteine conjugate beta-lyase. *Mol Pharmacol* 43: 660-665.

- Persson, B; Dahlander, A; Fredriksson, M; Brage, H; Ohlson, C; Axelson, O. (1989). Malignant lymphomas and occupational exposures. *Br J Ind Med* 46: 516-520.
- Persson, B; Fredriksson, M; Olsen, K; Boeryd, B; Axelson, O. (1993). Some occupational exposures as risk factors for malignant lymphomas. *Cancer* 72: 1773-1778.
- Persson, B and Fredriksson, M. (1999). Some risk factors for non-Hodgkin's lymphoma. *Int J Occup Med Environ Health* 12: 135-142.
- Pesch, B. (2008). personal communication from Beate Pesch, Forschungsinstitut für Arbeitsmedizin [BGFA], to Cheryl Scott, U.S. EPA, 21 February 2008
- Pesch, B; Haerting, J; Ranft, U; Klimpel, A; Oelschlägel, B; Schill, W. (2000a). Occupational risk factors for urothelial carcinoma: Agent-specific results from a case-control study in Germany. *Int J Epidemiol* 29: 238-274. <http://dx.doi.org/10.1093/ije/29.2.238>.
- Pesch, B; Haerting, J; Ranft, U; Klimpel, A; Oelschlägel, B; Schill, W. (2000b). Occupational risk factors for renal cell carcinoma: Agent-specific results from a case-control study in Germany. *Int J Epidemiol* 29: 1014-1024. <http://dx.doi.org/10.1093/ije/29.6.1014>.
- Peter, R; Böcker, R; Beaune, P; Iwasaki, M; Guengerich, F; Yang, C. (1990). Hydroxylation of chlorzoxazone as a specific probe for human liver cytochrome P-450IIE1. *Chem Res Toxicol* 3: 566-573. <http://dx.doi.org/10.1021/tx00018a012>.
- Peters, J; Preston-Martin, S; Yu, M. (1981). Brain tumors of children and occupational exposures of parents. *Science* 213: 235-237.
- Peters, J; Garabrant, D; Wright, W; Bernstein, L; Mack, T. (1985). Uses of a cancer registry in the assessment of occupational cancer risks. 69: 157-161.
- Peters, J; Cattley, R; Gonzalez, F. (1997). Role of PPAR alpha in the mechanism of action of the nongenotoxic carcinogen and peroxisome proliferator Wy-14,643. *Carcinogenesis* 18: 2029-2033.
- Pfaffenberger, C; Peoples, A; Enos, H. (1980). Distribution of volatile halogenated organic compounds between rat blood serum and adipose tissue. *Int J Environ Anal Chem* 8: 55-65. <http://dx.doi.org/10.1080/03067318008071879>.
- Phillips, T and Macdonald, R. (1971). Comparative effect of pethidine, trichloroethylene, and Entonox on fetal and neonatal acid-base and PO2. *Br Med J* 3: 558-560.
- Pikarsky, E; Porat, R; Stein, I; Abramovitch, R; Amit, S; Kasem, S; Gutkovich-Pyest, E; Urieli-Shoval, S; Galun, E; Ben-Neriah, Y. (2004). NF-kappaB functions as a tumour promoter in inflammation-associated cancer. *Nature* 431: 461-466. <http://dx.doi.org/10.1038/nature02924>.
- Plewa, M; Kargalioglu, Y; Vankerk, D; Minear, R; Wagner, E. (2002). Mammalian cell cytotoxicity and genotoxicity analysis of drinking water disinfection by-products. *Environ Mol Mutagen* 40: 134-142. <http://dx.doi.org/10.1002/em.10092>.
- Plummer, M; Best, N; Cowles, K; Vines, K. (2006). CODA: Convergence Diagnosis and Output Analysis for MCMC. *R News* 6: 7-11.
- Poet, T; Corley, R; Thrall, K; Edwards, J; Tanojo, H; Weitz, K; Hui, X; Maibach, H; Wester, R. (2000). Assessment of the percutaneous absorption of trichloroethylene in rats and humans using MS/MS real-time breath analysis and physiologically based pharmacokinetic modeling. *Toxicol Sci* 56: 61-72.
- Pogribny, I; Tryndyak, V; Woods, C; Witt, S; Rusyn, I. (2007). Epigenetic effects of the continuous exposure to peroxisome proliferator WY-14,643 in mouse liver are dependent upon peroxisome proliferator activated receptor alpha. *Mutat Res-Fundam Mol Mech Mutagen* 625: 62-71. <http://dx.doi.org/10.1016/j.mrfmmm.2007.05.004>.
- Pohl, H; Roney, N; Wilbur, S; Hansen, H; De Rosa, C. (2003). Six interaction profiles for simple mixtures. *Chemosphere* 53: 183-197. [http://dx.doi.org/10.1016/S0045-6535\(03\)00436-3](http://dx.doi.org/10.1016/S0045-6535(03)00436-3).
- Pollack, G; Li, R; Ermer, J; Shen, D. (1985). Effects of route of administration and repetitive dosing on the disposition kinetics of di(2-ethylhexyl) phthalate and its mono-de-esterified metabolite in rats. *Toxicol Appl Pharmacol* 79: 246-256. [http://dx.doi.org/10.1016/0041-008X\(85\)90346-1](http://dx.doi.org/10.1016/0041-008X(85)90346-1).
- Popper, H. (1986). Aging and the liver. In H Popper & F Schaffner (Eds.), *Progress in liver diseases*: Volume VIII (Vol. 8, pp. 659-683). Orlando, FL: Grune and Stratton.

- Porru, S; Placidi, D; Carta, A; Gelatti, U; Ribero, M; Tagger, A; Boffetta, P; Donato, F. (2001). Primary liver cancer and occupation in men: a case-control study in a high-incidence area in Northern Italy. *Int J Cancer* 94: 878-883.
- Porter, R. (1993). Follow-up of childhood leukemia incidence rate in Maricopa County standardized incidence ratio of area F to area R for 1987.
- Portier, C; Hedges, J; Hoel, D. (1986). Age-specific models of mortality and tumor onset for historical control animals in the National Toxicology Program's carcinogenicity experiments. *Cancer Res* 46: 4372-4378.
- Portier, C and Bailer, A. (1989). Testing for increased carcinogenicity using a survival-adjusted quantal response test. *Toxicol Sci* 12: 731-737.
- Poteracki, J and Walsh, K. (1998). Spontaneous neoplasms in control Wistar rats: A comparison of reviews. *Toxicol Sci* 45: 1-8. <http://dx.doi.org/10.1006/toxs.1998.2493>.
- Povey, A; Guppy, M; Wood, M; Knight, C; Black, C; Silman, A. (2001). Cytochrome P2 polymorphisms and susceptibility to scleroderma following exposure to organic solvents. *Arthritis Rheum* 44: 662-665. [http://dx.doi.org/10.1002/1529-0131\(200103\)44:3<662::AID-ANR115>3.0.CO;2-P](http://dx.doi.org/10.1002/1529-0131(200103)44:3<662::AID-ANR115>3.0.CO;2-P).
- Powell, J. (1945). Trichlorethylene: Absorption, elimination and metabolism. *Br J Ind Med* 2: 142-145.
- Prasanna, H; Hart, R; Magee, P. (1989). Effect of food restriction on the metabolism of dimethylnitrosamine (NDMA) in rats. *Anticancer Res* 9: 93-96.
- Pravecek, T; Channel, S; Schmidt, W; Kidney, J. (1996). Cytotoxicity and metabolism of dichloroacetic acid and trichloroacetic acid in B6C3F1 mouse liver tissue. *Vitro Toxicol* 9: 261-269.
- Prendergast, J; Jones, R; Jenkins, L; Siegel, J. (1967). Effects on experimental animals of long-term inhalation of trichloroethylene, carbon tetrachloride, 1,1,1-trichloroethane, dichlorodifluoromethane, and 1,1-dichloroethylene. *Toxicol Appl Pharmacol* 10: 270-289.
- Price, P; Hassett, C; Mansfield, J. (1978). Transforming activities of trichloroethylene and proposed industrial alternatives. *In Vitro* 14: 290-293.
- Price, P; Conolly, R; Chaisson, C; Gross, E; Young, J; Mathis, E; Tedder, D. (2003). Modeling interindividual variation in physiological factors used in PBPK models of humans. *Crit Rev Toxicol* 33: 469-503.
- Price, R; Taylor, S; Chivers, I; Arce-Tomas, M; Crutcher, E; Franchini, I; Slinovi, R; Cavazzini, S; Bergamaschi, E; Mutti, A; Vettori, M; Lauwers, R; Bernard, A; Kabanda, A; Roels, H; Thielemans, N; Hotz, P; De Broe, M; Elseviers, M; Nuyts, G; Gelpi, E; Hotter, G; Rosello, J; Ramis, I; Stolte, H; Fels, L; Eisenberger, U. (1996). Development and validation of new screening tests for nephrotoxic effects. *Hum Exp Toxicol* 1: S10-S19.
- Price, R; Wedeen, R; Lichtveld, M; Lybarger, J; Porter, G; Fels, L; Stolte, H; Safirstein, R; Mueller, P; Manley, S; Whiting, P; Gomez, R; Finn, W. (1999). Urinary biomarkers: Roles in risk assessment to environmental and occupational nephrotoxins: Monitoring of effects and evaluation of mechanisms of toxicity. *Ren Fail* 21: xiii-xviii.
- Prout, M; Provan, W; Green, T. (1985). Species differences in response to trichloroethylene: I pharmacokinetics in rats and mice. *Toxicol Appl Pharmacol* 79: 389-400.
- Pryor, G; Dickinson, J; Howd, R; Rebert, C. (1983). Transient cognitive deficits and high-frequency hearing loss in weanling rats exposed to toluene. *Neurobehav Toxicol Teratol* 5: 53-57.
- Pryor, G; Rebert, C; Howd, R. (1987). Hearing loss in rats caused by inhalation of mixed xylenes and styrene. *J Appl Toxicol* 7: 55-61.
- Purdue, M; Severson, R; Colt, J; Stewart, P; De Roos, A; Cerhan, J; Cozen, W; Davis, S; Hartge, P; Schenk, M; Blair, A. (2009). Degreasing and risk of non-Hodgkin lymphoma. *Occup Environ Med* 66: 557-560. <http://dx.doi.org/10.1136/oem.2008.040386>.
- Purdue, M; Bakke, B; Stewart, P; De Roos, A; Schenk, M; Lynch, C; Bernstein, L; Morton, L; Cerhan, J; Severson, R; Cozen, W; Davis, S; Rothman, N; Hartge, P; Colt, J. (2011). A case-control study of occupational exposure to trichloroethylene and non-Hodgkin lymphoma. *Environ Health Perspect* 119: 232-238. <http://dx.doi.org/10.1289/ehp.1002106>.

- Raaschou-Nielsen, O; Hansen, J; Christensen, J; Blot, W; McLaughlin, J; Olsen, J. (2001). Urinary concentrations of trichloroacetic acid in Danish workers exposed to trichloroethylene, 1947-1985. Am J Ind Med 39: 320-327.
- Raaschou-Nielsen, O; Hansen, J; Thomsen, B; Johansen, I; Lipworth, L; McLaughlin, J; Olsen, J. (2002). Exposure of Danish workers to trichloroethylene, 1947-1989. Appl Occup Environ Hyg 17: 693-703. <http://dx.doi.org/10.1080/1047322090096186>.
- Raaschou-Nielsen, O; Hansen, J; McLaughlin, J; Kolstad, H; Christensen, J; Tarone, R; Olsen, J. (2003). Cancer risk among workers at Danish companies using trichloroethylene: A cohort study. Am J Epidemiol 158: 1182-1192.
- Radican, L; Wartenberg, D; Rhoads, G; Schneider, D; Wedeen, R; Stewart, P; Blair, A. (2006). A retrospective occupational cohort study of end-stage renal disease in aircraft workers exposed to trichloroethylene and other hydrocarbons. J Occup Environ Med 48: 1-12.
- Radican, L; Blair, A; Stewart, P; Wartenberg, D. (2008). Mortality of aircraft maintenance workers exposed to trichloroethylene and other hydrocarbons and chemicals: Extended follow-up. J Occup Environ Med 50: 1306-1319. <http://dx.doi.org/10.1097/JOM.0b013e3181845f7f>.
- Raijmakers, M; Steegers, E; Peters, W. (2001). Glutathione S-transferases and thiol concentrations in embryonic and early fetal tissues. Hum Reprod 16: 2445-2450.
- Rake, J; Visser, G; Labrune, P; Leonard, J; Ullrich, K; Smit, G. (2002). Glycogen storage disease type I: Diagnosis, management, clinical course and outcome. Results of the European Study on Glycogen Storage Disease Type I (ESGSD I). Eur J Pediatr 161: S20-S34. <http://dx.doi.org/10.1007/s00431-002-0999-4>.
- Rakoff-Nahoum, S and Medzhitov, R. (2007). Regulation of spontaneous intestinal tumorigenesis through the adaptor protein MyD88. Science 317: 124-127. <http://dx.doi.org/10.1126/science.1140488>.
- Rakyan, V; Blewitt, M; Druker, R; Preis, J; Whitelaw, E. (2002). Metastable epialleles in mammals. Trends in Genetics 18: 348-351.
- Rakyan, V; Chong, S; Champ, M; Cuthbert, P; Morgan, H; Luu, K; Whitelaw, E. (2003). Transgenerational inheritance of epigenetic states at the murine Axin(Fu) allele occurs after maternal and paternal transmission. PNAS 100: 2538-2543. <http://dx.doi.org/10.1073/pnas.0436776100>.
- Ramanakumar, A; Nadon, L; Siemiatycki, J. (2008). Exposures in painting related occupations and risk of selected cancers: results from a case-control study in Montreal. Am J Ind Med 51: 419-427. <http://dx.doi.org/10.1002/ajim.20564>.
- Ramdhani, D; Kamijima, M; Yamada, N; Ito, Y; Yanagiba, Y; Nakamura, D; Okamura, A; Ichihara, G; Aoyama, T; Gonzalez, F; Nakajima, T. (2008). Molecular mechanism of trichloroethylene-induced hepatotoxicity mediated by CYP2E1. Toxicol Appl Pharmacol 231: 300-307. <http://dx.doi.org/10.1016/j.taap.2008.04.020>.
- Ramdhani, D; Kamijima, M; Wang, D; Ito, Y; Naito, H; Yanagiba, Y; Hayashi, Y; Tanaka, N; Aoyama, T; Gonzalez, F; Nakajima, T. (2010). Differential response to trichloroethylene-induced hepatosteatosis in wild-type and PPARalpha-humanized mice. Environ Health Perspect 118: 1557-1563. <http://dx.doi.org/10.1289/ehp.1001928>.
- Ramsdell, A and Markwald, R. (1997). Induction of endocardial cushion tissue in the avian heart is regulated, in part, by TGFbeta-3-mediated autocrine signaling. Dev Biol 188: 64-74. <http://dx.doi.org/10.1006/dbio.1997.8637>.
- Ramsey, J and Andersen, M. (1984). A physiologically based description of the inhalation pharmacokinetics of styrene in rats and humans. Toxicol Appl Pharmacol 73: 159-175. [http://dx.doi.org/10.1016/0041-008X\(84\)90064-4](http://dx.doi.org/10.1016/0041-008X(84)90064-4).
- Rao, M; Tatematsu, M; Subbarao, V; Ito, N; Reddy, J. (1986). Analysis of peroxisome proliferator-induced preneoplastic and neoplastic lesions of rat liver for placental form of glutathione S-transferase and gamma-glutamyltranspeptidase. Cancer Res 46: 5287-5290.
- Rapson, W; Nazar, M; Butsky, V. (1980). Mutagenicity produced by aqueous chlorination of organic compounds. Bull Environ Contam Toxicol 24: 590-596. <http://dx.doi.org/10.1007/BF01608160>.

- Rasmussen, K and Sabroe, S. (1986). Neuropsychological symptoms among metal workers exposed to halogenated hydrocarbons. Scand J Public Health 14: 161-168.
- Rasmussen, K; Sabroe, S; Wohlert, M; Ingerslev, H; Kappel, B; Nielsen, J. (1988). A genotoxic study of metal workers exposed to trichloroethylene Sperm parameters and chromosome aberrations in lymphocytes. Int Arch Occup Environ Health 60: 419-423.
- Rasmussen, K; Arlien-Søborg, P; Sabroe, S. (1993a). Clinical neurological findings among metal degreasers exposed to chlorinated solvents. Acta Neurol Scand 87: 200-204.
<http://dx.doi.org/10.1111/j.1600-0404.1993.tb04101.x>.
- Rasmussen, K; Brogren, C-H; Sabroe, S. (1993b). Subclinical affection of liver and kidney function and solvent exposure. Int Arch Occup Environ Health 64: 445-448.
<http://dx.doi.org/10.1007/BF00517951>.
- Rasmussen, K; Jeppesen, H; Sabroe, S. (1993c). Psychometric tests for assessment of brain function after solvent exposure. Am J Ind Med 24: 553-565. <http://dx.doi.org/10.1002/ajim.4700240506>.
- Rasmussen, K; Jeppesen, H; Sabroe, S. (1993d). Solvent-induced chronic toxic encephalopathy. Am J Ind Med 23: 779-792. <http://dx.doi.org/10.1002/ajim.4700230511>.
- Raunio, H; Hakkola, J; Hukkanen, J; Pelkonen, O; Edwards, R; Boobis, A; Anttila, S. (1998). Expression of xenobiotic metabolising cytochrome P450s in human pulmonary tissue. Arch Toxicol Suppl 20: 465-469.
- Ray, W; Griffin, M; Schaffner, W. (1987). Psychotropic drug use and the risk of hip fracture. 316/7 (363-369).
- Reamon-Buettner, S and Borlak, J. (2007). A new paradigm in toxicology and teratology: Altering gene activity in the absence of DNA sequence variation. Reprod Toxicol 24: 20-30.
<http://dx.doi.org/10.1016/j.reprotox.2007.05.002>.
- Rebert, C; Day, V; Matteucci, M; Pryor, G. (1991). Sensory-evoked potentials in rats chronically exposed to trichloroethylene: Predominant auditory dysfunction. Neurotoxicol Teratol 13: 83-90.
- Rebert, C; Boyes, W; Pryor, G; Svensgaard, D; Kassay, K; Gordon, G; Shinsky, N. (1993). Combined effects of solvents on the rat's auditory system: Styrene and trichloroethylene. Int J Psychophysiol 14: 49-59.
- Rebert, C; Schwartz, R; Svendsgaard, D; Pryor, G; Boyes, W. (1995). Combined effects of paired solvents on the rat's auditory system. Toxicology 105: 345-354. [http://dx.doi.org/10.1016/0300-483X\(95\)03232-5](http://dx.doi.org/10.1016/0300-483X(95)03232-5).
- Reif, J; Burch, J; Nuckols, J; Metzger, L; Ellington, D; Anger, W. (2003). Neurobehavioral effects of exposure to trichloroethylene through a municipal water supply. Environ Res 93: 248-258.
- Rennix, C; Quinn, M; Amoroso, P; Eisen, E; Wegman, D. (2005). Risk of breast cancer among enlisted Army women occupationally exposed to volatile organic compounds. Am J Ind Med 48: 157-167.
- Reznik, G; Ward, J; Hardisty, J; Russfield, A. (1979). Renal carcinogenic and nephrotoxic effects of the flame retardant tris(2,3-dibromopropyl) phosphate in F344 rats and (C57BL/6N X C3H/HeN)F1 mice. J Natl Cancer Inst 63: 205-212.
- Rhee, I; Bachman, K; Park, B; Jair, K-W; Yen, R-W; Schuebel, K; Cui, H; Feinberg, A; Lengauer, C; Kinzler, K; Baylin, S; Vogelstein, B. (2002). DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. Nature 416: 552-556. <http://dx.doi.org/10.1038/416552a>.
- Rhomberg, L. (2000). Dose-response analyses of the carcinogenic effects of trichloroethylene in experimental animals. Environ Health Perspect 108: 343-358.
- Rice, D and Barone, S, Jr. (2000). Critical periods of vulnerability for the developing nervous system: Evidence from humans and animal models. Environ Health Perspect 108: 511-533.
- Richard, A and Hunter, E. (1996). Quantitative structure-activity relationships for the developmental toxicity of haloacetic acids in mammalian whole embryo culture. Teratology 53: 352-360.
[http://dx.doi.org/10.1002/\(SICI\)1096-9926\(199606\)53:6<352::AID-TERA6>3.0.CO;2-1](http://dx.doi.org/10.1002/(SICI)1096-9926(199606)53:6<352::AID-TERA6>3.0.CO;2-1).
- Richardson, D; Terschüren, C; Hoffmann, W. (2008). Occupational risk factors for non-Hodgkin's lymphoma: a population-based case-control study in Northern Germany. Am J Ind Med 51: 258-268. <http://dx.doi.org/10.1002/ajim.20552>.

- Richardson, F; Boucheron, J; Dyroff, M; Popp, J; Swenberg, J. (1986). Biochemical and morphologic studies of heterogeneous lobe responses in hepatocarcinogenesis. *Carcinogenesis* 7: 247-251. <http://dx.doi.org/10.1093/carcin/7.2.247>.
- Richmond, R; Carter, J; Carter, H; Daniel, F; DeAngelo, A. (1995). Immunohistochemical analysis of dichloroacetic acid (DCA)-induced hepatocarcinogenesis in male Fischer (F344) rats. *Cancer Lett* 92: 67-76. [http://dx.doi.org/10.1016/0304-3835\(94\)03756-9](http://dx.doi.org/10.1016/0304-3835(94)03756-9).
- Riederer, F; Luborzewski, A; God, R; Bringmann, G; Scholz, J; Feineis, D; Moser, A. (2002). Modification of tyrosine hydroxylase activity by chloral derived beta-carbolines in vitro. *J Neurochem* 81: 814-819.
- Ries, L; Melbert, D; Krapcho, M; Stinchcomb, D; Howlader, N; Horner, M; Mariotto, A; Miller, B; Feuer, E; Altekruse, S; Lewis, D; Clegg, L; Eisner, M; Reichman, M; Edwards, B. (2008). SEER cancer statistics review, 1975-2005. pp. 907. Bethesda, MD: U.S. National Institutes of Health, National Cancer Institute. http://seer.cancer.gov/csr/1975_2005.
- Ripp, S; Overby, L; Philpot, R; Elfarra, A. (1997). Oxidation of cysteine S-conjugates by rabbit liver microsomes and cDNA-expressed flavin-containing mono-oxygenases: Studies with S-(1,2-dichlorovinyl)-L-cysteine, S-(1,2,2-trichlorovinyl)-L-cysteine, S-allyl-L-cysteine, and S-benzyl-L-cysteine. *Mol Pharmacol* 51: 507-515.
- Ripp, S; Itagaki, K; Philpot, R; Elfarra, A. (1999). Species and sex differences in expression of flavin-containing monooxygenase form 3 in liver and kidney microsomes. *Drug Metab Dispos* 27: 46-52.
- Ristoff, E and Larsson, A. (2007). Inborn errors in the metabolism of glutathione. *Orphanet J Rare Dis* 2: 16. <http://dx.doi.org/10.1186/1750-1172-2-16>.
- Ritz, B. (1999a). Cancer mortality among workers exposed to chemicals during uranium processing. *J Occup Environ Med* 41: 556-566.
- Ritz, B. (1999b). Radiation exposure and cancer mortality in uranium processing workers. *Epidemiology* 10: 531-538.
- Ritz, B; Morgenstern, H; Froines, J; Young, B. (1999). Effects of exposure to external ionizing radiation on cancer mortality in nuclear workers monitored for radiation at Rocketdyne/Atomics International. *Am J Ind Med* 35: 21-31.
- Ritz, B; Morgenstern, H; Crawford-Brown, D; Young, B. (2000). The effects of internal radiation exposure on cancer mortality in nuclear workers at Rocketdyne/Atomics International. *Environ Health Perspect* 108: 743-751.
- Robbiano, L; Mereto, E; Migliazz Morando, A; Pastore, P; Brambilla, G. (1998). Increased frequency of micronucleated kidney cells in rats exposed to halogenated anaesthetics. *Mutat Res* 413: 1-6.
- Robbiano, L; Baroni, D; Carrozzino, R; Mereto, E; Brambilla, G. (2004). DNA damage and micronuclei induced in rat and human kidney cells by six chemicals carcinogenic to the rat kidney. *Toxicology* 204: 187-195. <http://dx.doi.org/10.1016/j.tox.2004.06.057>.
- Robins, J. (1987). A graphical approach to the identification and estimation of causal parameters in mortality studies with sustained exposure periods. 40: 139S-161S. [http://dx.doi.org/10.1016/S0021-9681\(87\)80018-8](http://dx.doi.org/10.1016/S0021-9681(87)80018-8).
- Rodenbeck, S; Sanderson, L; Rene, A. (2000). Maternal exposure to trichloroethylene in drinking water and birth-weight outcomes. *Arch Environ Health* 55: 188-194.
- Rodriguez, C; Mahle, D; Gearhart, J; Mattie, D; Lipscomb, J; Cook, R; Barton, H. (2007). Predicting age-appropriate pharmacokinetics of six volatile organic compounds in the rat utilizing physiologically based pharmacokinetic modeling. *Toxicol Sci* 98: 43-56. <http://dx.doi.org/10.1093/toxsci/kfm082>.
- Roskams, T; Yang, S; Koteish, A; Duriez, A; DeVos, R; Huang, X; Achten, R; Verslype, C; Diehl, A. (2003). Oxidative stress and oval cell accumulation in mice and humans with alcoholic and nonalcoholic fatty liver disease. *Am J Pathol* 163: 1301-1311. [http://dx.doi.org/10.1016/S0002-9440\(10\)63489-X](http://dx.doi.org/10.1016/S0002-9440(10)63489-X).

- Roskams, T; Theise, N; Balabaud, C; Bhagat, G; Bhathal, P; Bioulac-Sage, P; Brunt, E; Crawford, J; Crosby, H; Desmet, V; Finegold, M; Geller, S; Gouw, A; Hytiroglou, P; Knisely, A; Kojiro, M; JHNakanuma, Y; Olynyk, J; Park, Y; Portmann, B; Saxena, R; Scheuer, P; Strain, A; Thung, S; Wanless, I; West, A. (2004). Nomenclature of the finer branches of the biliary tree: Canals, ductules, and ductular reactions in human livers. *Hepatology* 39: 1739-1745.
<http://dx.doi.org/10.1002/hep.20130>.
- Rossi, A; Migliore, L; Barale, R; Loprieno, N. (1983). In vivo and in vitro mutagenicity studies of a possible carcinogen, trichloroethylene, and its two stabilizers, epichlorohydrin and 1,2-epoxybutane. *Birth Defects Res B Dev Reprod Toxicol* 3: 75-87.
- Rothman, K. (1986). Modern epidemiology. Boston, MA: Little Brown & Co.
- Rothman, K and Greenland, S. (1998). Modern epidemiology (2nd ed.). Philadelphia, PA: Lippincott, Williams, & Wilkins.
- Ruckenstein, M. (2004). Autoimmune inner ear disease. *Curr Opin Otolaryngol Head Neck Surg* 12: 426-430.
- Rudolf, K and DePinho, R. (2001). Telomeres and telomerase in experimental liver cirrhosis. In D Shafritz (Ed.), *The liver: Biology and pathobiology* (pp. 1001–1011). Philadelphia, PA: Lippincott Williams and Wilkins.
- Rufer, E; Hacker, T; Lough, J; Smith, S. (2008). Low-dose trichloroethylene exposure during valvuloseptal morphogenesis causes ventricular septal defects in hatched chicks. *Toxicologist* 102: 314.
- Rufer, E; Hacker, T; Flentke, G; Drake, V; Brody, M; Lough, J; Smith, S. (2010). Altered cardiac function and ventricular septal defect in avian embryos exposed to low-dose trichloroethylene. *Toxicol Sci* 113: 444-452. <http://dx.doi.org/10.1093/toxsci/kfp269>.
- Ruijten, M; Verberk, M; Sallé, H. (1991). Nerve function in workers with long term exposure to trichloroethene. *Br J Ind Med* 48: 87-92.
- Russo, A; Pacchierotti, F; Metalli, P. (1984). Nondisjunction induced in mouse spermatogenesis by chloral hydrate, a metabolite of trichloroethylene. *Environ Mutagen* 6: 695-703.
<http://dx.doi.org/10.1002/em.2860060507>.
- Russo, A and Levis, A. (1992a). Further evidence for the aneuploidogenic properties of chelating agents: Induction of micronuclei in mouse male germ cells by EDTA. *Environ Mol Mutagen* 19: 125-131.
- Russo, A and Levis, A. (1992b). Detection of aneuploidy in male germ cells of mice by means of a meiotic micronucleus assay. *Mutat Res* 281: 187-191. [http://dx.doi.org/10.1016/0165-7992\(92\)90007-5](http://dx.doi.org/10.1016/0165-7992(92)90007-5).
- Russo, A; Stocco, A; Majone, F. (1992). Identification of kinetochore-containing (CREST+) micronuclei in mouse bone marrow erythrocytes. *Mutagenesis* 7: 195-197.
- Rusyn, I; Bradham, C; Cohn, L; Schoonhoven, R; Swenberg, J; Brenner, D; Thurman, R. (1999). Corn oil rapidly activates nuclear factor-kappaB in hepatic Kupffer cells by oxidant-dependent mechanisms. *Carcinogenesis* 20: 2095-2100.
- Rusyn, I; Peters, J; Cunningham, M. (2006). Modes of action and species-specific effects of di-(2-ethylhexyl)phthalate in the liver. *Crit Rev Toxicol* 36: 459-479.
<http://dx.doi.org/10.1080/10408440600779065>.
- SAB. (U.S. Environmental Protection Agency, Science Advisory Board). (2010a). [Summary minutes of the SAB quality review teleconference of an SAB panel draft report reviewing EPA's IRIS assessment for trichloroethylene]. Washington, DC.
<http://yosemite.epa.gov/sab/sabproduct.nsf/a84bfee16cc358ad85256ccd006b0b4b/a60f87e98e51f46f852577b300518230!OpenDocument&Date=2010-12-151>.
- SAB. (U.S. Environmental Protection Agency, Science Advisory Board). (2010b). Summary minutes of the U.S. Environmental Protection Agency Trichloroethylene Review Panel public teleconference. Washington, DC.

<http://yosemite.epa.gov/sab/sabproduct.nsf/a84bfee16cc358ad85256ccd006b0b4b/6cdc83d5129a174f8525772d004d77c0!OpenDocument&Date=2010-06-24>.

- SAB. (U.S. Environmental Protection Agency, Science Advisory Board). (2011). Review of EPA's Draft Assessment entitled "Toxicological Review of Trichloroethylene" (October 2009). (EPA-SAB-11-002). Washington, DC.
[http://yosemite.epa.gov/sab/sabproduct.nsf/B73D5D39A8F184BD85257817004A1988/\\$File/EPA-SAB-11-002-unsigned.pdf](http://yosemite.epa.gov/sab/sabproduct.nsf/B73D5D39A8F184BD85257817004A1988/$File/EPA-SAB-11-002-unsigned.pdf).
- Sabel, G and Clark, T. (1984). Volatile organic compounds as indicators of municipal solid waste leachate contamination. *Waste Manag Res* 2: 119-130. [http://dx.doi.org/10.1016/0734-242X\(84\)90135-6](http://dx.doi.org/10.1016/0734-242X(84)90135-6).
- Sagawa, K; Nishitani, H; Kawai, H; Kuge, Y; Ikeda, M. (1973). Transverse lesion of spinal cord after accidental exposure to trichloroethylene. *Int Arch Arbeitsmed* 31: 257-264.
- Saghir, S and Schultz, I. (2002). Low-dose pharmacokinetics and oral bioavailability of dichloroacetate in naive and GST-zeta-depleted rats. *Environ Health Perspect* 110: 757-763.
- Saihan, E; Burton, J; Heaton, K. (1978). A new syndrome with pigmentation, scleroderma, gynaecomastia, Raynaud's phenomenon and peripheral neuropathy. *Br J Dermatol* 99: 437-440.
- Saillenfait, A; Langonne, I; Sabate, J. (1995). Developmental toxicity of trichloroethylene, tetrachloroethylene and four of their metabolites in rat whole embryo culture. *Arch Toxicol* 70: 71-82. <http://dx.doi.org/10.1007/BF02733666>.
- Sakurai, T; Maeda, S; Chang, L; Karin, M. (2006). Loss of hepatic NF-kappa B activity enhances chemical hepatocarcinogenesis through sustained c-Jun N-terminal kinase 1 activation. *PNAS* 103: 10544-10551. <http://dx.doi.org/10.1073/pnas.0603499103>.
- Sallmen, M; Lindbohm, M; Anttila, A; Kyryonen, P; Taskinen, H; Nykyri, E; Hemminki, K. (1998). Time to pregnancy among the wives of men exposed to organic solvents. *Occup Environ Med* 55: 24-30.
- Sallmén, M; Lindbohm, M; Kyyrönen, P; Nykyri, E; Anttila, A; Taskinen, H; Hemminki, K. (1995). Reduced fertility among women exposed to organic solvents. *Am J Ind Med* 27: 699-713. <http://dx.doi.org/10.1002/ajim.4700270506>.
- Salvan, A. (1990) Occupational exposure and cancer mortality at an electrical manufacturing plant: A case-control study. University of California, Los Angeles, CA.
- Salvan, A; Stayner, L; Steenland, K; Smith, R. (1995). Selecting an exposure lag period. *Epidemiology* 6: 387-390.
- Salvini, M; Binaschi, S; Riva, M. (1971). Evaluation of the psychophysiological functions in humans exposed to trichloroethylene. *Br J Ind Med* 28: 293-295.
- Sanchez, I and Bull, R. (1990). Early induction of reparative hyperplasia in the liver of B6C3F1 mice treated with dichloroacetate and trichloroacetate. *Toxicology* 64: 33-46. [http://dx.doi.org/10.1016/0300-483X\(90\)90097-Z](http://dx.doi.org/10.1016/0300-483X(90)90097-Z).
- Sanders, V; Kauffmann, B; White, K; Douglas, K; Barnes, D; Sain, L; Bradshaw, T; Borzelleca, J; Munson, A. (1982a). Toxicology of chloral hydrate in the mouse. *Environ Health Perspect* 44: 137-146.
- Sanders, V; Tucker, A; White, K, Jr; Kauffmann, B; Hallett, P; Carchman, R; Borzelleca, J; Munson, A. (1982b). Humoral and cell-mediated immune status in mice exposed to trichloroethylene in the drinking water. *Toxicol Appl Pharmacol* 62: 358-368. [http://dx.doi.org/10.1016/0041-008X\(82\)90138-7](http://dx.doi.org/10.1016/0041-008X(82)90138-7).
- Santibanez, M; Vioque, J; Alguacil, J; Barber, X; de la Hera, G; Kauppinen, T. (2008). Occupational exposures and risk of oesophageal cancer by histological type: a case-control study in eastern Spain. *Occup Environ Med* 65: 774-781. <http://dx.doi.org/10.1136/oem.2007.037929>.
- Sanz, N; Díez-Fernández, C; Cascales, M. (1996). Variations of hepatic antioxidant systems and DNA ploidy in rats aged 2 to 8 months. *Biochim Biophys Acta* 1315: 123-130. [http://dx.doi.org/10.1016/0925-4439\(95\)00113-1](http://dx.doi.org/10.1016/0925-4439(95)00113-1).

- Sapkota, A; Williams, D; Buckley, T. (2005). Tollbooth workers and mobile source-related hazardous air pollutants: How protective is the indoor environment? Environ Sci Technol 39: 2936-2943.
- Sarangapani, R; Gentry, P; Covington, T; Teeguarden, J; Clewell HJ, L. (2003). Evaluation of the potential impact of age- and gender-specific lung morphology and ventilation rate on the dosimetry of vapors. Inhal Toxicol 15: 987-1016.
- Sargent, L; Zhou, X; Keck, C; Sanderson, N; Zimonjic, D; Popescu, N; Thorgeirsson, S. (1999). Nonrandom cytogenetic alterations in hepatocellular carcinoma from transgenic mice overexpressing c-Myc and transforming growth factor-alpha in the liver. Am J Pathol 154: 1047-1055.
- Sato, A; Nakajima, T; Fujiwara, Y; Murayama, N. (1977). A pharmacokinetic model to study the excretion of trichloroethylene and its metabolites after an inhalation exposure. Br J Ind Med 34: 56-63.
- Sato, A and Nakajima, T. (1978). Differences following skin or inhalation exposure in the absorption and excretion kinetics of trichloroethylene and toluene. Br J Ind Med 35: 43-49.
- Sato, A and Nakajima, T. (1979). A vial-equilibration method to evaluate the drug-metabolizing enzyme activity for volatile hydrocarbons. Toxicol Appl Pharmacol 47: 41-46.
[http://dx.doi.org/10.1016/0041-008X\(79\)90069-3](http://dx.doi.org/10.1016/0041-008X(79)90069-3).
- Sato, A; Nakajima, T; Koyama, Y. (1980). Effects of chronic ethanol consumption on hepatic metabolism of aromatic and chlorinated hydrocarbons in rats. Br J Ind Med 37: 382-386.
<http://dx.doi.org/10.1136/oem.37.4.382>.
- Sato, A; Nakajima, T; Koyama, Y. (1981). Dose-related effects of a single dose of ethanol on the metabolism in rat liver of some aromatic and chlorinated hydrocarbons. Toxicol Appl Pharmacol 60: 8-15. [http://dx.doi.org/10.1016/0041-008X\(81\)90129-0](http://dx.doi.org/10.1016/0041-008X(81)90129-0).
- Sato, A; Nakajima, T; Koyama, Y. (1983). Interaction between ethanol and carbohydrate on the metabolism in rat liver of aromatic and chlorinated hydrocarbons. Toxicol Appl Pharmacol 68: 242-249. [http://dx.doi.org/10.1016/0041-008X\(83\)90008-X](http://dx.doi.org/10.1016/0041-008X(83)90008-X).
- Sato, A and Nakajima, T. (1985). Enhanced metabolism of volatile hydrocarbons in rat liver following food deprivation, restricted carbohydrate intake and administration of ethanol, phenobarbital, polychlorinated biphenyl and 3-methylcholanthrene: A comparative study. Xenobiotica 15: 67-75. <http://dx.doi.org/10.3109/00498258509045336>.
- Sato, A; Endoh, K; Johanson, G. (1991a). The Use of Models to Investigate the Toxicokinetic Behavior of Organic Solvents. Lewis Publishers, Chelsea, Michigan: 131-152.
- Sato, A; Endoh, K; Kaneko, T; Johanson, G. (1991b). A simulation study of physiological factors affecting pharmacokinetic behaviour of organic solvent vapours. Occup Environ Med 48: 342-347.
- Sato, A. (1993). Confounding factors in biological monitoring of exposure to organic solvents. Int Arch Occup Environ Health 65: S61-S67.
- Saunders, R. (1967). A new hazard in closed environmental atmospheres. Arch Environ Health 14: 380-384.
- Sausen, P and Elfarra, A. (1990). Cysteine conjugate S-oxidase. Characterization of a novel enzymatic activity in rat hepatic and renal microsomes. J Biol Chem 265: 6139-6145.
- Sausen, P and Elfarra, A. (1991). Reactivity of cysteine S-conjugate sulfoxides: formation of S-[1-chloro-2-(S-glutathionyl)vinyl]-L-cysteine sulfoxide by the reaction of S-(1,2-dichlorovinyl)-L-cysteine sulfoxide with glutathione. Chem Res Toxicol 4: 655-660.
<http://dx.doi.org/10.1021/tx00024a010>.
- Savolainen, H; Pfäffli, P; Tengén, M; Vainio, H. (1977). Trichloroethylene and 1,1,1-trichloroethane: Effects on brain and liver after five days intermittent inhalation. Arch Toxicol 38: 229-237.
<http://dx.doi.org/10.1007/BF00293657>.
- Saxena, D; Srivastava, R; Lal, B; Chandra, S. (1987). The effect of lead exposure on the testis of growing rats. Exp Toxicol Pathol 31: 249-252.

- Saygun, M; Cakmak, A; Ekici, A; Pinar, T; Bulcun, E; Ulu, N; Ekici, M. (2007). Five annual observations of respiratory findings in gun factory workers exposed to solvents. *J Occup Environ Med* 49: 909-912. <http://dx.doi.org/10.1097/JOM.0b013e318124a8f3>.
- Sbrana, I; Di Sibio, A; Lomi, A; Scarcelli, V. (1993). C-mitosis and numerical chromosome aberration analyses in human lymphocytes: 10 known or suspected spindle poisons. *Mutat Res* 287: 57-70. [http://dx.doi.org/10.1016/0027-5107\(93\)90145-6](http://dx.doi.org/10.1016/0027-5107(93)90145-6).
- Scharf, J; Dombrowski, F; Ramadori, G. (2001). The IGF axis and hepatocarcinogenesis. *Molecular Pathology* 54: 138-144.
- Schatten, H and Chakrabarti, A. (1998). Centrosome structure and function is altered by chloral hydrate and diazepam during the first reproductive cell cycles in sea urchin eggs. *Eur J Cell Biol* 75: 9-20.
- Scheff, P and Wadden, R. (1993). Receptor modeling of volatile organic compounds. I: Emission inventory and validation. *Environ Sci Technol* 27: 617-625. <http://dx.doi.org/10.1021/es00041a005>.
- Schenk, M; Purdue, M; Colt, J; Hartge, P; Blair, A; Stewart, P; Cerhan, J; De Roos, A; Cozen, W; Severson, R. (2009). Occupation/industry and risk of non-Hodgkin's lymphoma in the United States. *Occup Environ Med* 66: 23-31. <http://dx.doi.org/10.1136/oem.2007.036723>.
- Schlehofer, B; Heuer, C; Blettner, M; Niehoff, D; Wahrendorf, J. (1995). Occupation, smoking and demographic factors, and renal cell carcinoma in Germany. *Int J Epidemiol* 24: 51-57.
- Schlesselman, J. (1982). Case control studies: Design, conduct, analysis. New York, NY: Oxford University Press.
- Schlipkötter, H; Fodor, G; Winneke, G. (1974). Zum Einfluss neurotoxischer Industriegifte auf unfallgenerierendes Fehlverhalten bei Kontroll- und Steuertätigkeiten. Wilhelmshaven, Dortmund: Wirtschaftsverlag Nordwest.
- Schmucker, D; Mooney, J; Jones, A. (1978). Stereological analysis of hepatic fine structure in the Fischer 344 rat. Influence of sublobular location and animal age. *J Cell Biol* 78: 319-337.
- Schmucker, D. (2001). Liver function and phase I drug metabolism in the elderly: A paradox. *Drugs Aging* 18: 837-851.
- Schmucker, D. (2005). Age-related changes in liver structure and function: Implications for disease? *Exp Gerontol* 40: 650-659. <http://dx.doi.org/10.1016/j.exger.2005.06.009>.
- Schraml, P; Zhaou, M; Richter, J; Brüning, T; Pommer, M; Sauter, G; Mihatsch, M; Moch, H. (1999). [Analysis of kidney tumors in trichloroethylene exposed workers by comparative genomic hybridization and DNA sequence analysis]. *Verh Dtsch Ges Pathol* 83: 218-224.
- Schulte-Hermann, R; Timmermann-Trosiener, I; Schuppler, J. (1986). Facilitated expression of adaptive responses to phenobarbital in putative pre-stages of liver cancer. *Carcinogenesis* 7: 1651-1655. <http://dx.doi.org/10.1093/carcin/7.10.1651>.
- Schultz, I; Merdink, J; Gonzalez-Leon, A; Bull, R. (1999). Comparative toxicokinetics of chlorinated and brominated haloacetates in F344 rats. *Toxicol Appl Pharmacol* 158: 103-114. <http://dx.doi.org/10.1006/taap.1999.8698>.
- Schultz, I; Merdink, J; Gonzalez-Leon, A; Bull, R. (2002). Dichloroacetate toxicokinetics and disruption of tyrosine catabolism in B6C3F1 mice: Dose-response relationships and age as a modifying factor. *Toxicology* 173: 229-247. [http://dx.doi.org/10.1016/S0300-483X\(02\)00034-3](http://dx.doi.org/10.1016/S0300-483X(02)00034-3).
- Schumucker, D; Mooney, J; Jones, A. (1977). Age-related changes in the hepatic endoplasmic reticulum: A quantitative analysis. *Science* 197: 1005-1008. <http://dx.doi.org/10.1126/science.887935>.
- Schwartz-Arad, D; Zajicek, G; Bartfeld, E. (1989). The streaming liver IV: DNA content of the hepatocyte increases with its age. *Liver Int* 9: 93-99. <http://dx.doi.org/10.1111/j.1600-0676.1989.tb00385.x>.
- Schwetz, B; Leong, B; Gehring, P. (1975). The effect of maternally inhaled trichloroethylene, perchloroethylene, methyl chloroform, and methylene chloride on embryonal and fetal development in mice and rats. *Toxicol Appl Pharmacol* 32: 84-96. [http://dx.doi.org/10.1016/0041-008X\(75\)90197-0](http://dx.doi.org/10.1016/0041-008X(75)90197-0).

- Scott, C and Chiu, W. (2006). Trichloroethylene cancer epidemiology: A consideration of select issues. Environ Health Perspect 114: 1471-1478.
- Scott, J; Forkert, P; Oulton, M; Rasmusson, M-G; Temple, S; Fraser, M; Whitefield, S. (1988). Pulmonary toxicity of trichloroethylene: induction of changes in surfactant phospholipids and phospholipase A2 activity in the mouse lung. Exp Mol Pathol 49: 141-150.
- Seelbach, A; Fissler, B; Madle, S. (1993). Further evaluation of a modified micronucleus assay with V79 cells for detection of aneugenic effects. Mutat Res 303: 163-169.
- Seidler, A; Möhner, M; Berger, J; Mester, B; Deeg, E; Elsner, G; Nieters, A; Becker, N. (2007). Solvent exposure and malignant lymphoma: A population-based case-control study in Germany. J Occup Med Toxicol 2: 2. <http://dx.doi.org/10.1186/1745-6673-2-2>.
- Seitz, C; Lin, Q; Deng, H; Khavari, P. (1998). Alterations in NF-kappaB function in transgenic epithelial tissue demonstrate a growth inhibitory role for NF-kappaB. PNAS 95: 2307-2312.
- Seitz, H and Stickel, F. (2006). Risk factors and mechanisms of hepatocarcinogenesis with special emphasis on alcohol and oxidative stress. Biol Chem 387: 349-360. <http://dx.doi.org/10.1515/bc.2006.047>.
- Selden, A; Hultberg, B; Ulander, A; Ahlborg, GJ. (1993). Trichloroethylene exposure in vapour degreasing and the urinary excretion of N-acetyl- β -D-glucosaminidase. Arch Toxicol 67: 224-226.
- Seldén, A and Ahlborg, G. (1991). Mortality and cancer morbidity after exposure to military aircraft fuel. Aviat Space Environ Med 62: 789-794.
- Selgrade, M and Gilmour, M. (2010). Suppression of pulmonary host defenses and enhanced susceptibility to respiratory bacterial infection in mice following inhalation exposure to trichloroethylene and chloroform. J Immunotoxicol 7: 350-356. <http://dx.doi.org/10.3109/1547691X.2010.520139>.
- Sellers, E; Lang, M; Koch-Weser, J; LeBlanc, E; Kalant, H. (1972). Interaction of chloral hydrate and ethanol in man. I. Metabolism. Clin Pharmacol Ther 13: 37-49.
- Sellers, E; Lang-Sellers, M; Koch-Weser, J. (1978). Comparative metabolism of chloral hydrate and triclofos. J Clin Pharmacol 18: 457-461.
- Selmin, O; Thorne, P; Caldwell, P; Taylor, M. (2008). Trichloroethylene and trichloroacetic acid regulate calcium signaling pathways in murine embryonal carcinoma cells p19. Cardiovasc Toxicol 8: 47-56. <http://dx.doi.org/10.1007/s12012-008-9014-2>.
- Seng, J; Agrawal, N; Horsley, E; Leakey, T; Scherer, E; Xia, S; Allaben, W; Leakey, J. (2003). Toxicokinetics of chloral hydrate in ad libitum-fed, dietary-controlled, and calorically restricted male B6C3F1 mice following short-term exposure. Toxicol Appl Pharmacol 193: 281-292. <http://dx.doi.org/10.1016/j.taap.2003.07.008>.
- Seo, M; Ikeda, K; Okamura, T; Kida, K; Satoh, M; Inagaki, N; Nagai, H; Nagase, H. (2008). Enhancing effect of chlorinated organic solvents on histamine release and inflammatory mediator production. Toxicology 243: 75-83. <http://dx.doi.org/10.1016/j.tox.2007.09.024>.
- Seppäläinen, A and Antti-Poika, M. (1983). Time course of electrophysiological findings for patients with solvent poisoning. A descriptive study. Scand J Work Environ Health 9: 15-24.
- Sexton, K; Adgate, J; Church, T; Ashley, D; Needham, L; Ramachandran, G; Fredrickson, A; Ryan, A. (2005). Children's exposure to volatile organic compounds as determined by longitudinal measurements in blood. Environ Health Perspect 113: 342-349. <http://dx.doi.org/10.1289/ehp.7412>.
- Sexton, K; Mongin, S; Adgate, J; Pratt, G; Ramachandran, G; Stock, T; Morandi, M. (2007). Estimating volatile organic compound concentrations in selected microenvironments using time-activity and personal exposure data. J Toxicol Environ Health A 70: 465-476. <http://dx.doi.org/10.1080/15287390600870858>.
- Shafer, T; Bushnell, P; Benignus, V; Woodward, J. (2005). Perturbation of voltage-sensitive Ca²⁺ channel function by volatile organic solvents. J Pharmacol Exp Ther 315: 1109-1118. <http://dx.doi.org/10.1124/jpet.105.090027>.

- Shah, J and Singh, H. (1988). Distribution of volatile organic chemicals in outdoor and indoor air: A national VOCs data base. Environ Sci Technol 22: 1381-1388.
<http://dx.doi.org/10.1021/es00177a001>.
- Shah, Y; Morimura, K; Yang, Q; Tanabe, T; Takagi, M; Gonzalez, F. (2007). Peroxisome proliferator-activated receptor alpha regulates a microRNA-mediated signaling cascade responsible for hepatocellular proliferation. Mol Cell Biol 27: 4238-4247. <http://dx.doi.org/10.1128/MCB.00317-07>.
- Shankar, K; Vaidya, V; Corton, J; Bucci, T; Liu, J; Waalkes, M; Mehendale, H. (2003). Activation of PPAR-alpha in streptozotocin-induced diabetes is essential for resistance against acetaminophen toxicity. J Exp Biol 17: 1748-1750. <http://dx.doi.org/10.1096/fj.02-1186fje>.
- Shannon, H; Haines, T; Bernholz, C; Julian, J; Verma, D; Jamieson, E; Walsh, C. (1988). Cancer morbidity in lamp manufacturing workers. Am J Ind Med 14: 281-290.
- Shao, J; Stapleton, P; Lin, Y; Gallagher, E. (2007). Cytochrome p450 and glutathione s-transferase mRNA expression in human fetal liver hematopoietic stem cells. Drug Metab Dispos 35: 168-175. <http://dx.doi.org/10.1124/dmd.106.012757>.
- Sharpe, C; Rochon, J; Adam, J; Suissa, S. (1989). Case-control study of hydrocarbon exposures in patients with renal cell carcinoma. CMAJ 140: 1309-1318.
- Shearer, B and Hoekstra, W. (2003). Recent advances in peroxisome proliferator-activated receptor science. Curr Med Chem 10: 267-280. <http://dx.doi.org/10.2174/1567204052930970>.
- Sheffels, P; Schroeder, J; Altuntas, T; Liggitt, H; Kharasch, E. (2004). Role of cytochrome P4503A in cysteine S-conjugates sulfoxidation and the nephrotoxicity of the sevoflurane degradation product fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether (compound A) in rats. Chem Res Toxicol 17: 1177-1189. <http://dx.doi.org/10.1021/tx049899e>.
- Sher, T; Yi, H; McBride, O; Gonzalez, F. (1993). cDNA cloning, chromosomal mapping, and functional characterization of the human peroxisome proliferator activated receptor. Biochemistry 32: 5598-5604. <http://dx.doi.org/10.1021/bi00072a015>.
- Sherins, R and Hodgen, G. (1976). Testicular gamma glutamyl-transpeptidase: An index of Sertoli cell function in man. Journal of Reproduction and Fertility 48: 191-193.
<http://dx.doi.org/10.1530/jrf.0.0480191>.
- Shiao, Y; Rice, J; Anderson, L; Diwan, B; Hard, G. (1998). von Hippel-Lindau gene mutations in N-nitrosodimethylamine-induced rat renal epithelial tumors. J Natl Cancer Inst 90: 1720-1723.
- Shiao, Y; Kamata, S; Li, L; Hooth, M; DeAngelo, A; Anderson, L; Wolf, D. (2002). Mutations in the VHL gene from potassium bromate-induced rat clear cell renal tumors. Cancer Lett 187: 207-214.
- Shih, C-L; Chen, H-H; Chiu, T. (2001). Acute exposure to trichloroethylene differentially alters the susceptibility to chemoconvulsants in mice. Toxicology 162: 35-42.
[http://dx.doi.org/10.1016/S0300-483X\(01\)00358-4](http://dx.doi.org/10.1016/S0300-483X(01)00358-4).
- Shimada, T; Swanson, A; Leber, P; Williams, G. (1985). Activities of chlorinated ethane and ethylene compounds in the Salmonella/rat microsome mutagenesis and rat hepatocyte/DNA repair assays under vapor phase exposure conditions. Cell Biol Toxicol 1: 159-179.
<http://dx.doi.org/10.1007/BF00120162>.
- Shimada, T; Yamazaki, H; Mimura, M; Inui, Y; Guengerich, F. (1994). Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: Studies with liver microsomes of 30 Japanese and 30 Caucasians. J Pharmacol Exp Ther 270: 414-423.
- Shindell, S and Ulrich, S. (1985). A cohort study of employees of a manufacturing plant using trichlorethylene. J Occup Med 27: 577-579.
- Shirasu, Y; Moriya, M; Kato, K; Furuhashi, A; Kada, T. (1976). Mutagenicity screening of pesticides in the microbial system. Mutat Res 40: 19-30. [http://dx.doi.org/10.1016/0165-1218\(76\)90018-5](http://dx.doi.org/10.1016/0165-1218(76)90018-5).
- Shlomo, M; Frenkel, A; Hager, M; Skulsky, M; Sulkis, J; Himelfarbe, M. (2002). Effects of occupational exposure to mercury or chlorinated hydrocarbons on the auditory pathway. Noise Health 4: 71-77.

- Short, B. (1993). Cell proliferation and renal carcinogenesis. Environ Health Perspect 101 Suppl 5: 115-120.
- Shroads, A; Guo, X; Dixit, V; Liu, H-P; James, M; Stacpoole, P. (2008). Age-dependent kinetics and metabolism of dichloroacetate: Possible relevance to toxicity. J Pharmacol Exp Ther 324: 1163-1171. <http://dx.doi.org/10.1124/jpet.107.134593>.
- Shu, X; Stewart, P; Wen, W-Q; Han, D; Potter, J; Buckley, J; Heineman, E; Robison, L. (1999). Parental occupational exposure to hydrocarbons and risk of acute lymphocytic leukemia in offspring. Cancer Epidemiol Biomarkers Prev 8: 783-791.
- Shu, X; Perentesis, J; Wen, W; Buckley, J; Boyle, E; Ross, J; Robison, L; Group, CsO. (2004). Parental exposure to medications and hydrocarbons and ras mutations in children with acute lymphoblastic leukemia: A report from the Children's Oncology Group. Cancer Epidemiol Biomarkers Prev 13: 1230-1235.
- Shultz, J and Weiner, H. (1979). Alteration of the enzymology of chloral hydrate reduction in the presence of ethanol. Biochem Pharmacol 28: 3379-3384. [http://dx.doi.org/10.1016/0006-2952\(79\)90076-5](http://dx.doi.org/10.1016/0006-2952(79)90076-5).
- Siegel, J and Kasmin, F. (1997). Biliary tract diseases in the elderly: Management and outcomes. Gut 41: 433-435. <http://dx.doi.org/10.1136/gut.41.4.433>.
- Siemiatycki, J; Wacholder, S; Richardson, L; Dewar, R; Gérin, M. (1987). Discovering carcinogens in the occupational environment. Methods of data collection and analysis of a large case-referent monitoring system. Scand J Work Environ Health 13: 486-492.
- Siemiatycki, J; Dewar, R; Nadon, L; Gérin, M. (1994). Occupational risk factors for bladder cancer: Results from a case-control study in Montreal, Quebec, Canada. Am J Epidemiol 140: 1061-1080.
- Siemiatycki, J; Fritschi, L; Nadon, L; Gérin, M. (1997). Reliability of an expert rating procedure for retrospective assessment of occupational exposures in community-based case-control studies. Am J Ind Med 31: 280-286.
- Siemiatycki, J (Ed.). (1991). Risk factors for cancer in the workplace. Boca Raton, FL: CRC Press.
- Sigal, S; Rajvanshi, P; Gorla, G; Sokhi, R; Saxena, R; Gebhard, D, Jr; Reid, L; Gupta, S. (1999). Partial hepatectomy-induced polyploidy attenuates hepatocyte replication and activates cell aging events. Am J Physiol 276: G1260-G1272.
- Simmon, V; Kauhanen, K; Tardiff, R. (1977). Mutagenic activity of chemicals identified in drinking water. In BBFS D Scott (Ed.), Progress in Genetic Toxicology (Vol. 2, pp. 249-268). New York; Amsterdam: Elsevier/North Holland Press.
- Simmons, J; Boyes, W; Bushnell, P; Raymer, J; Limsakun, T; McDonald, A; Sey, Y; Evans, M. (2002). A physiologically based pharmacokinetic model for trichloroethylene in the male Long-Evans rat. Toxicol Sci 69: 3-15. <http://dx.doi.org/10.1093/toxsci/69.1.3>.
- Singh, M and Sinha, U. (1976). Chloral hydrate induced haploidization in *Aspergillus nidulans*. Experientia 32: 1144-1145. <http://dx.doi.org/10.1007/BF01927592>.
- Singh, M and Sinha, U. (1979). Mitotic haploidization and growth of *Aspergillus nidulans* on media containing chloral hydrate. J Cytol Genet 14: 1-4.
- Sinks, T; Lushniak, B; Haussler, B; Snizek, J; Deng, J; Roper, P; Dill, P; Coates, R. (1992). Renal cell cancer among paperboard printing workers. Epidemiology 3: 483-489.
- Skender, L; Karacic, V; Prpic-Majic, D. (1991). A comparative study of human levels of trichloroethylene and tetrachloroethylene after occupational exposure. Arch Environ Health 46: 174-178. <http://dx.doi.org/10.1080/15459620802510866>.
- Slikker, W, Jr; Andersen, M; Bogdanffy, M; Bus, J; Cohen, S; Conolly, R; David, R; Doerrer, N; Dorman, D; Gaylor, D; Hattis, D; Rogers, J; Setzer, R; Swenberg, J; Wallace, K. (2004a). Dose-dependent transitions in mechanisms of toxicity. Toxicol Appl Pharmacol 201: 203-225. <http://dx.doi.org/10.1016/j.taap.2004.06.019>.
- Slikker, W, Jr; Andersen, M; Bogdanffy, M; Bus, J; Cohen, S; Conolly, R; David, R; Doerrer, N; Dorman, D; Gaylor, D; Hattis, D; Rogers, J; Setzer, R; Swenberg, J; Wallace, K. (2004b). Dose-

- dependent transitions in mechanisms of toxicity: Case studies. *Toxicol Appl Pharmacol* 201: 226-294. <http://dx.doi.org/10.1016/j.taap.2004.06.027>.
- Slawińska-Kowalska, M; Jedlińska, U; Rzadzińska, A; Rajkowska, E. (1999). [Hair cell regeneration in a chicken's inner ear after damage due to exposure to industrial noise]. *Otolaryngol Pol* 53: 469-477.
- Smith, G. (1970). The investigation of the mental effects of trichlorethylene. *Ergonomics* 13: 580-586.
- Smith, M; Randall, J; Read, E; Stober, J. (1989). Teratogenic activity of trichloroacetic acid in the rat. *Teratology* 40: 445-451. <http://dx.doi.org/10.1002/tera.1420400506>.
- Smith, M; Randall, J; Read, E; Stober, J. (1992). Developmental toxicity of dichloroacetate in the rat. *Teratology* 46: 217-223. <http://dx.doi.org/10.1002/tera.1420460305>.
- Snedcor, G and Cochran, W. (1980). Statistical methods (7th ed.). Ames, IA: Iowa State University Press.
- Snyder, R; Pullman, J; Carter, J; Carter, H; DeAngelo, A. (1995). In vivo administration of dichloroacetic acid suppresses spontaneous apoptosis in murine hepatocytes. *Cancer Res* 55: 3702-3705.
- Song, B; Veech, R; Saenger, P. (1990). Cytochrome P450IIE1 is elevated in lymphocytes from poorly controlled insulin-dependent diabetics. *J Clin Endocrinol Metab* 71: 1036-1040.
- Soni, M; Mangipudy, R; Mumtaz, M; Mehendale, H. (1998). Tissue repair response as a function of dose during trichloroethylene hepatotoxicity. *Toxicol Sci* 42: 158-165.
<http://dx.doi.org/10.1006/toxs.1998.2427>.
- Soni, M; Ramaiah, S; Mumtaz, M; Clewell, H; Mehendale, H. (1999). Toxicant-inflicted injury and stimulated tissue repair are opposing toxicodynamic forces in predictive toxicology. *Regul Toxicol Pharmacol* 29: 165-174. <http://dx.doi.org/10.1006/rtpb.1998.1280>.
- Sonnenfeld, N; Hertz-Pannier, I; Kaye, W. (2001). Tetrachloroethylene in drinking water and birth outcomes at the US Marine Corps Base at Camp Lejeune, North Carolina. *Am J Epidemiol* 154: 902-908.
- Sora, S and Agostini Carbone, M. (1987). Chloral hydrate, methylmercury hydroxide and ethidium bromide affect chromosomal segregation during meiosis of *Saccharomyces cerevisiae*. *Mutat Res Lett* 190: 13-17. [http://dx.doi.org/10.1016/0165-7992\(87\)90075-3](http://dx.doi.org/10.1016/0165-7992(87)90075-3).
- Soresi, M; Giannitrapani, L; D'Antona, F; Florena, A; La Spada, E; Terranova, A; Cervello, M; D'Alessandro, N; Montalto, G. (2006). Interleukin-6 and its soluble receptor in patients with liver cirrhosis and hepatocellular carcinoma. *World J Gastroenterol* 12: 2563-2568.
- Soucek, B and Vlachova, D. (1960). Excretion of trichloroethylene metabolites in human urine. *Br J Ind Med* 17: 60-64.
- Spiegelhalter, D; Best, N; Carlin, B; Van der Linde, A. (2002). Bayesian measures of model complexity and fit. *J R Stat Soc Series B Stat Methodol* 64: 583-639. <http://dx.doi.org/10.1111/1467-9868.00353>.
- Spiegelhalter, D; Thomas, A; Best, N; Lunn, D. (2003). WinBugs version 1.4 user manual. Cambridge, UK: MRC Biostatistics Unit.
- Spielberg, S. (1996). N-acetyltransferases: pharmacogenetics and clinical consequences of polymorphic drug metabolism. *J Pharmacokinet Biopharm* 24: 509-519.
- Spirtas, R; Stewart, P; Lee, J; Marano, D; Forbes, C; Grauman, D; Pettigrew, H; Blair, A; Hoover, R; Cohen, J. (1991). Retrospective cohort mortality study of workers at an aircraft maintenance facility: I. Epidemiological results. *Br J Ind Med* 48: 515-530.
<http://dx.doi.org/10.1136/oem.48.8.515>.
- Squillace, P; Moran, M; Price, C. (2004). VOCs in shallow groundwater in new residential/commercial areas of the United States. *Environ Sci Technol* 38: 5327-5338.
<http://dx.doi.org/10.1021/es0349756>.
- SRI. (SRI International). (1992). 1992 directory of chemical producers: United States of America. Menlo Park, CA: Author.
- Srivastava, S. (1991). Effect of di(2-ethylhexyl)phthalate on 17 β -hydroxysteroid dehydrogenase activity in testis of rat. *Toxicol Lett* 57: 235-239. [http://dx.doi.org/10.1016/0378-4274\(91\)90150-5](http://dx.doi.org/10.1016/0378-4274(91)90150-5).

- Staats, D; Fisher, J; Connolly, R. (1991). Gastrointestinal absorption of xenobiotics in physiologically based pharmacokinetic models: A two-compartment description. *Drug Metab Dispos* 19: 144-148.
- Stadler, J and Kennedy, G, Jr. (1996). Evaluation of the sensory irritation potential of volatile organic chemicals from carpets -- Alone and in combination. *Food Chem Toxicol* 34: 1125-1130.
- Stanley, L; Blackburn, D; Devereaux, S; Foley, J; Lord, P; Maronpot, R; Orton, T; Anderson, M. (1994). Ras mutations in methylclofenapate-induced B6C3F1 and C57BL/10J mouse liver tumours. *Carcinogenesis* 15: 1125-1131.
- Staples, R. (1974). Detection of visceral alterations in mammalian fetuses. *Teratology* 9: A37-A38.
- Stauber, A and Bull, R. (1997). Differences in phenotype and cell replicative behavior of hepatic tumors induced by dichloroacetate (DCA) and trichloroacetate (TCA). *Toxicol Appl Pharmacol* 144: 235-246. <http://dx.doi.org/10.1006/taap.1997.8159>.
- Stauber, A; Bull, R; Thrall, B. (1998). Dichloroacetate and trichloroacetate promote clonal expansion of anchorage-independent hepatocytes in vivo and in vitro. *Toxicol Appl Pharmacol* 150: 287-294. <http://dx.doi.org/10.1006/taap.1998.8417>.
- Steinberg, A. (1993). Should chloral hydrate be banned. *Pediatrics* 92: 442-446.
- Steinmaus, C; Smith, A; Jones, R; Smith, M. (2008). Meta-analysis of benzene exposure and non-Hodgkin lymphoma: Biases could mask an important association. *Occup Environ Med* 65: 371-378. <http://dx.doi.org/10.1136/oem.2007.036913>.
- Stemhagen, A; Slade, J; Altman, R; Bill, J. (1983). Occupational risk factors and liver cancer: A retrospective case-control study of primary liver cancer in New Jersey. *Am J Epidemiol* 117: 443-454.
- Stemmer, K; Ellinger-Ziegelbauer, H; Ahr, H; Dietrich, D. (2007). Carcinogen-specific gene expression profiles in short-term treated Eker and wild-type rats indicative of pathways involved in renal tumorigenesis. *Cancer Res* 67: 4052-4068. <http://dx.doi.org/10.1158/0008-5472.CAN-06-3587>.
- Stenner, R; Merdink, J; Stevens, D; Springer, D; Bull, R. (1997). Enterohepatic recirculation of trichloroethanol glucuronide as a significant source of trichloroacetic acid. *Metabolites of trichloroethylene*. *Drug Metab Dispos* 25: 529-535.
- Stephens, E; Taylor, J; Kaplan, N; Yang, C; Hsieh, L; Lucier, G; Bell, D. (1994). Ethnic variation in the CYP2E1 gene: Polymorphism analysis of 695 African-Americans, European-Americans and Taiwanese. *Pharmacogenetics* 4: 185-192.
- Stephenson, R. (1956). A modification of receptor theory. *Br J Pharmacol Chemother* 11: 379-393.
- Steup, D; Wiersma, D; McMillan, D; Sipes, I. (1991). Pretreatment with drinking water solutions containing trichloroethylene or chloroform enhances the hepatotoxicity of carbon tetrachloride in Fischer 344 rats. *Toxicol Sci* 16: 798-809. <http://dx.doi.org/10.1093/toxsci/16.4.798>.
- Steup, D; Hall, P; McMillan, D; Sipes, I. (1993). Time course of hepatic injury and recovery following coadministration of carbon tetrachloride and trichloroethylene in Fischer-344 rats. *Toxicol Pathol* 21: 327-334.
- Stevens, J and Jakoby, W. (1983). Cysteine conjugate beta-lyase. *Mol Pharmacol* 23: 761-765.
- Stevens, J. (1985). Cysteine conjugate beta-lyase activities in rat kidney cortex: Subcellular localization and relationship to the hepatic enzyme. *Biochem Biophys Res Commun* 129: 499-504. [http://dx.doi.org/10.1016/0006-291X\(85\)90179-2](http://dx.doi.org/10.1016/0006-291X(85)90179-2).
- Stevens, J; Robbins, J; Byrd, R. (1986). A purified cysteine conjugate beta-lyase from rat kidney cytosol. Requirement for an alpha-keto acid or an amino acid oxidase for activity and identity with soluble glutamine transaminase K. *J Biol Chem* 261: 15529-15537.
- Stevens, J; Ayoubi, N; Robbins, J. (1988). The role of mitochondrial matrix enzymes in the metabolism and toxicity of cysteine conjugates. *J Biol Chem* 263: 3395-3401.
- Stevens, J; Hatzinger, P; Hayden, P. (1989). Quantitation of multiple pathways for the metabolism of nephrotoxic cysteine conjugates using selective inhibitors of L-alpha-hydroxy acid oxidase (L-amino acid oxidase) and cysteine conjugate beta-lyase. *Drug Metab Dispos* 17: 297-303.

- Stewart, B; Le Mesurier, S; Lykke, A. (1979). Correlation of biochemical and morphological changes induced by chemical injury to the lung. *Chem Biol Interact* 26: 321-338.
- Stewart, P; Lee, J; Marano, D; Spirtas, R; Forbes, C; Blair, A. (1991). Retrospective cohort mortality study of workers at an aircraft maintenance facility. II. Exposures and their assessment. *Br J Ind Med* 48: 531-537.
- Stewart, P; Lees, P; Francis, M. (1996). Quantification of historical exposures in occupational cohort studies. *Scand J Work Environ Health* 22: 405-414.
- Stewart, P and Dosemeci, M. (2005). Exposure estimation in occupational epidemiology studies. Third Meeting on Assessing Human Health Risks of Trichloroethylene. Irvine, CA.
- Stewart, R; Dodd, H; Gay, H; Erley, D. (1970). Experimental human exposure to trichloroethylene. *Arch Environ Health* 20: 64-71.
- Stewart, R; Hake, C; Peterson, J. (1974a). "Degreaser's flush". *Arch Environ Health* 29: 1-5.
- Stewart, R; Hake, C; Peterson, J. (1974b). Degreasers' flush: dermal response to trichloroethylene and ethanol. *Arch Environ Occup Health* 29: 1-5.
- Stewart, W and Correa-Villaseor, A. (1991). False positive exposure errors and low exposure prevalence in community-based case-control studies. *Appl Occup Environ Hyg* 6: 534-540.
- Stopps, G and McLaughlin, M. (1967). Psychophysiological testing of human subjects exposed to solvent vapors. *Am Ind Hyg Assoc J* 28: 43-50.
- Stott, W; Quast, J; Watanabe, P. (1982). The pharmacokinetics and macromolecular interactions of trichloroethylene in mice and rats. *Toxicol Appl Pharmacol* 62: 137-151.
- Stuckhardt, J and Poppe, S. (1984). Fresh visceral examination of rat and rabbit fetuses used in teratogenicity testing. *Teratog Carcinog Mutagen* 4: 181-188.
- Styles, J; Kelly, M; Pritchard, N; Ekcombe, C. (1988). A species comparison of acute hyperplasia induced by the peroxisome proliferator methylclofenapate: Involvement of the binucleated hepatocyte. *Carcinogenesis* 9: 1647-1655. <http://dx.doi.org/10.1093/carcin/9.9.1647>.
- Styles, J; Wyatt, I; Coutts, C. (1991). Trichloroacetic acid: Studies on uptake and effects on hepatic DNA and liver growth in mouse. *Carcinogenesis* 12: 1715-1719. <http://dx.doi.org/10.1093/carcin/12.9.1715>.
- Su, Q and Bannasch, P. (2003). Relevance of hepatic preneoplasia for human hepatocarcinogenesis. *Toxicol Pathol* 31: 126-133. <http://dx.doi.org/10.1080/01926230390173905>.
- Subramoniam, A; Goel, S; Pandya, K; Seth, P. (1989). Influence of trichloroethylene treatment on phosphoinositides in rat brain. *Toxicol Lett* 49: 55-60. [http://dx.doi.org/10.1016/0378-4274\(89\)90101-X](http://dx.doi.org/10.1016/0378-4274(89)90101-X).
- Sung, T-I; Chen, P-C; Lee, L-H; Lin, Y-P; Hsieh, G-Y; Wang, J-D. (2007). Increased standardized incidence ratio of breast cancer in female electronics workers. *BMC Public Health* 7: 102. <http://dx.doi.org/10.1186/1471-2458-7-102>.
- Sung, T-I; Wang, J-D; Chen, P-C. (2008). Increased risk of cancer in the offspring of female electronics workers. *Reprod Toxicol* 25: 115-119. <http://dx.doi.org/10.1016/j.reprotox.2007.08.004>.
- Supplementary data for TCE assessment: Cancer rodents input data. (2011).
- Supplementary data for TCE assessment: Cancer rodents model selections. (2011).
- Supplementary data for TCE assessment: Cancer rodents plots. (2011).
- Supplementary data for TCE assessment: Cancer rodents results. (2011).
- Supplementary data for TCE assessment: Cancer rodents uncertainty analysis. (2011).
- Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-inhalation histograms, inhalation bioassays. (2011).
- Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-inhalation historams, oral bioassays. (2011).
- Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-oral histograms, inhalation bioassay. (2011).
- Supplementary data for TCE assessment: Cancer rodents uncertainty CSF-oral histograms, oral bioassays. (2011).

- Supplementary data for TCE assessment: Hack mouse population calibration evaluation. (2011).
- Supplementary data for TCE assessment: Hack mouse subject calibration. (2011).
- Supplementary data for TCE assessment: Human population example. (2011).
- Supplementary data for TCE assessment: Mouse population example. (2011).
- Supplementary data for TCE assessment: Non-cancer HECs altPOD plots from rodent inhalation studies. (2011).
- Supplementary data for TCE assessment: Non-cancer HECs altPOD plots from rodent oral studies. (2011).
- Supplementary data for TCE assessment: Non-cancer HECs plots from human inhalation studies. (2011).
- Supplementary data for TCE assessment: Non-cancer HECs plots from rodent inhalation studies. (2011).
- Supplementary data for TCE assessment: Non-cancer HEDs altPOD plots from rodent oral studies. (2011).
- Supplementary data for TCE assessment: Non-cancer HEDs plots from human inhalation studies. (2011).
- Supplementary data for TCE assessment: Non-cancer HEDs plots from rodent inhalation studies. (2011).
- Supplementary data for TCE assessment: Non-cancer input data contin. (2011).
- Supplementary data for TCE assessment: Non-cancer input data dichot. (2011).
- Supplementary data for TCE assessment: Non-cancer plots contin. (2011).
- Supplementary data for TCE assessment: Non-cancer plots dichot. (2011).
- Supplementary data for TCE assessment: Non-cancer results contin. (2011).
- Supplementary data for TCE assessment: Non-cancer results dichot. (2011).
- Supplementary data for TCE assessment: Rat population example. (2011).
- Supplementary data for TCE assessment: Rodents time to tumor results. (2011).
- Sverdrup, B; Kallberg, H; Bengtsson, C; Lundberg, I; Padyukov, L; Alfredsson, L; Klareskog, L; Group, EIoRAS. (2005). Association between occupational exposure to mineral oil and rheumatoid arthritis: results from the Swedish EIRA case-control study. *Arthritis Res Ther* 7: R1296-R1303. <http://dx.doi.org/10.1186/ar1824>.
- Swaen, G. (1995). Increased incidence of renal cell tumours in a cohort of cardboard workers exposed to trichloroethylene. *Arch Toxicol* 70: 127-128. <http://dx.doi.org/10.1007/BF02733674>.
- Swan, S; Shaw, G; Harris, J; Neutra, R. (1989). Congenital cardiac anomalies in relation to water contamination, Santa Clara County, California, 1981-1983. *Am J Epidemiol* 129: 885-893.
- Sweeney, C; Farrow, D; Schwartz, S; Eaton, D; Checkoway, H; Vaughan, T. (2000). Glutathione S-transferase M1, T1, and P1 polymorphisms as risk factors for renal cell carcinoma: A case-control study. *Cancer Epidemiol Biomarkers Prev* 9: 449-454.
- Sweeney, L; Kirman, C; Gargas, M; Dugard, P. (2009). Contribution of trichloroacetic acid to liver tumors observed in perchloroethylene (perc)-exposed mice. *Toxicology* 260: 77-83. <http://dx.doi.org/10.1016/j.tox.2009.03.008>.
- Sweet, C and Vermette, S. (1992). Toxic volatile organic compounds in urban air in Illinois. *Environ Sci Technol* 26: 165-173. <http://dx.doi.org/10.1021/es00025a020>.
- Szyf, M. (2007). The dynamic epigenome and its implications in toxicology. *Toxicol Sci* 100: 7-23. <http://dx.doi.org/10.1093/toxsci/kfm177>.
- Szymańska, K; Moore, L; Rothman, N; Chow, W; Waldman, F; Jaeger, E; Waterboer, T; Foretova, L; Navratilova, M; Janout, V; Kollarova, H; Zaridze, D; Matveev, V; Mates, D; Szczeszenia-Dabrowska, N; Holcatova, I; Bencko, V; Le Calvez-Kelm, F; Villar, S; Pawlita, M; Boffetta, P; Hainaut, P; Brennan, P. (2010). TP53, EGFR, and KRAS mutations in relation to VHL inactivation and lifestyle risk factors in renal-cell carcinoma from central and eastern Europe. 293: 92-98. <http://dx.doi.org/10.1016/j.canlet.2009.11.024>.
- Takashima, K; Ito, Y; Gonzalez, F; Nakajima, T. (2008). Different mechanisms of DEHP-induced hepatocellular adenoma tumorigenesis in wild-type and Ppar alpha-null mice. *J Occup Health* 50: 169-180.
- Takeda, S; Kondo, M; Kumada, T; Koshikawa, T; Ueda, R; Nishio, M; Osada, H; Suzuki, H; Nagatake, M; Washimi, O; Takagi, K; Takahashi, T; Nakao, A. (1996). Allelic-expression imbalance of the

- insulin-like growth factor 2 gene in hepatocellular carcinoma and underlying disease. Oncogene 12: 1589-1592.
- Takubo, K and Kaminishi, M. (2001). [Diseases of the digestive tract and telomere lengths: Significance and problems of telomere measurement]. Nippon Shokakibyo Gakkai Zasshi 98: 144-150.
- Tanaka, K; Smith, P; Stromberg, P; Eydelloth, R; Herold, E; Grossman, S; Frank, J; Hertzog, P; Soper, K; Keenan, K. (1992). Studies of early hepatocellular proliferation and peroxisomal proliferation in Sprague-Dawley rats treated with tumorigenic doses of clofibrate. Toxicol Appl Pharmacol 116: 71-77. [http://dx.doi.org/10.1016/0041-008X\(92\)90146-J](http://dx.doi.org/10.1016/0041-008X(92)90146-J).
- Tang, X; Li, L; Huang, J; Deng, Y. (2002). Guinea pig maximization test for trichloroethylene and its metabolites. Biomed Environ Sci 15: 113-118.
- Tang, X; Que, B; Song, X; Li, S; Yang, X; Wang, H; Huang, H; Kamijima, M; Nakajima, T; Lin, Y; Li, L. (2008). Characterization of liver injury associated with hypersensitive skin reactions induced by trichloroethylene in the guinea pig maximization test. J Occup Health 50: 114-121.
- Tanguay, R; Jorquera, R; Poudrier, J; St-Louis, M. (1996). Tyrosine and its catabolites: From disease to cancer. Acta Biochim Pol 43: 209-216.
- Tao, L; Kramer, P; Ge, R; Pereira, M. (1998). Effect of dichloroacetic acid and trichloroacetic acid on DNA methylation in liver and tumors of female B6C3F1 mice. Toxicol Sci 43: 139-144. <http://dx.doi.org/10.1093/toxsci/43.2.139>.
- Tao, L; Yang, S; Xie, M; Kramer, P; Pereira, M. (2000). Effect of trichloroethylene and its metabolites, dichloroacetic acid and trichloroacetic acid, on the methylation and expression of c-jun and c-myc protooncogenes in mouse liver: Prevention by methionine. Toxicol Sci 54: 399-407. <http://dx.doi.org/10.1093/toxsci/54.2.399>.
- Tao, L; Li, Y; Kramer, P; Wang, W; Pereira, M. (2004a). Hypomethylation of DNA and the insulin-like growth factor-II gene in dichloroacetic and trichloroacetic acid-promoted mouse liver tumors. Toxicology 196: 127-136. <http://dx.doi.org/10.1016/j.tox.2003.11.011>.
- Tao, L; Wang, W; Li, L; Kramer, P; Pereira, M. (2004b). Effect of dibromoacetic acid on DNA methylation, glycogen accumulation, and peroxisome proliferation in mouse and rat liver. Toxicol Sci 82: 62-69. <http://dx.doi.org/10.1093/toxsci/kfh266>.
- Tarsetti, F; Lenzi, R; Salvi, R; Schuler, E; Rijhsinghani, K; Lenzen, R; Tavoloni, N. (1993). Liver carcinogenesis associated with feeding of ethionine in a choline-free diet: Evidence against a role of oval cells in the emergence of hepatocellular carcinoma. Hepatology 18: 596-603. <http://dx.doi.org/10.1002/hep.1840180319>.
- Taskinen, H; Anttila, A; Lindbohm, M-L; Sallmén, M; Hemminki, K. (1989). Spontaneous abortions and congenital malformations among the wives of men occupationally exposed to organic solvents. Scand J Work Environ Health 15: 345-352.
- Taskinen, H; Kyryonen, P; Hemminki, K. (1994). Laboratory work and pregnancy outcome. J Occup Med 36: 311-319.
- Tateishi, M; Suzuki, S; Shimizu, H. (1978). Cysteine conjugate beta-lyase in rat liver. A novel enzyme catalyzing formation of thiol-containing metabolites of drugs. J Biol Chem 253: 8854-8859.
- Tateishi, T; Nakura, H; Asoh, M; Watanabe, M; Tanaka, M; Kumai, T; Takashima, S; Imaoka, S; Funae, Y; Yabusaki, Y; Kamataki, T; Kobayashi, S. (1997). A comparison of hepatic cytochrome P450 protein expression between infancy and postinfancy. Life Sci 61: 2567-2574. [http://dx.doi.org/10.1016/S0024-3205\(97\)01011-4](http://dx.doi.org/10.1016/S0024-3205(97)01011-4).
- Tatham, L; Tolbert, P; Kjeldsberg, C. (1997). Occupational risk factors for subgroups of non-Hodgkin's lymphoma. Epidemiology 8: 551-558.
- Taylor, D; Lagory, K; Zaccaro, D; Pfahl, R; Laurie, R. (1985). Effect of trichloroethylene on the exploratory and locomotor activity of rats exposed during development. Sci Total Environ 47: 415-420. [http://dx.doi.org/10.1016/0048-9697\(85\)90345-6](http://dx.doi.org/10.1016/0048-9697(85)90345-6).
- Templin, M; Parker, J; Bull, R. (1993). Relative formation of dichloroacetate and trichloroacetate from trichloroethylene in male B6C3F1 mice. Toxicol Appl Pharmacol 123: 1-8. <http://dx.doi.org/10.1006/taap.1993.1214>.

- Templin, M; Parker, J; Bull, R. (1995a). Erratum for the article "Relative formation of dichloroacetate and trichloroacetate from trichloroethylene in male B6C3F1 mice. Toxicol Appl Pharmacol 133: 177. <http://dx.doi.org/10.1006/taap.1995.1139>.
- Templin, M; Stevens, D; Stenner, R; Bonate, P; Tuman, D; Bull, R. (1995b). Factors affecting species differences in the kinetics of metabolites of trichloroethylene. J Toxicol Environ Health 44: 435-447.
- Tenant, R; Elwell, M; Spalding, J; Griesemer, R. (1991). Evidence that toxic injury is not always associated with induction of chemical carcinogenesis. Mol Carcinog 4: 420-440. <http://dx.doi.org/10.1002/mc.2940040604>.
- Terracini, B and Parker, V. (1965). A pathological study on the toxicity of s-dichlorovinyl-l-cysteine. Food Cosmet Toxicol 3: 67-74.
- Teschke, K; Olshan, A; Daniels, J; De Roos, A; Parks, C; Schulz, M; Vaughan, T. (2002). Occupational exposure assessment in case-control studies: Opportunities for improvement. Occup Environ Med 59: 575-594. <http://dx.doi.org/10.1136/oem.59.9.575>.
- Thai, S-F; Allen, J; DeAngelo, A; George, M; Fuscoe, J. (2001). Detection of early gene expression changes by differential display in the livers of mice exposed to dichloroacetic acid. Carcinogenesis 22: 1317-1322. <http://dx.doi.org/10.1093/carcin/22.8.1317>.
- Thai, S-F; Allen, J; DeAngelo, A; George, M; Fuscoe, J. (2003). Altered gene expression in mouse livers after dichloroacetic acid exposure. Mutat Res Rev Mutat Res 543: 167-180. [http://dx.doi.org/10.1016/S1383-5742\(03\)00014-0](http://dx.doi.org/10.1016/S1383-5742(03)00014-0).
- Tham, R; Larsby, B; Ödkvist, L; Norlander, B; Hydén, D; Aschan, G; Bertler, Å. (1979). The influence of trichloroethylene and related drugs on the vestibular system. Acta Pharmacol Toxicol 44: 336-342. <http://dx.doi.org/10.1111/j.1600-0773.1979.tb02340.x>.
- Tham, R; Bunnfors, I; Eriksson, B; Larsby, B; Lindgren, S; Odkvist, L. (1984). Vestibulo-ocular disturbances in rats exposed to organic solvents. Acta Pharmacol Toxicol 54: 58-63. <http://dx.doi.org/10.1111/j.1600-0773.1984.tb01895.x>.
- Tharappel, J; Cunningham, M; Spear, B; Glauert, H. (2001). Differential activation of hepatic NF- κ B in rats and hamsters by the peroxisome proliferators Wy-14,643, gemfibrozil, and dibutyl phthalate. Toxicol Sci 62: 20-27. <http://dx.doi.org/10.1093/toxsci/62.1.20>.
- Thiele, D; Eigenbrodt, E; Ware, A. (1982). Cirrhosis after repeated trichloroethylene and 1,1,1-trichloroethane exposure. Gastroenterology 83: 926-929.
- Thorgeirsson, S. (2006). Genomic decoding of hepatocellular carcinoma. Gastroenterology 131: 1344-1346. <http://dx.doi.org/10.1053/j.gastro.2006.08.060>.
- Thrall, K and Poet, T. (2000). Determination of biokinetic interactions in chemical mixtures using real-time breath analysis and physiologically based pharmacokinetic modeling. J Toxicol Environ Health A 59: 653-670. <http://dx.doi.org/10.1080/009841000156646>.
- Till, C; Koren, G; Rovet, J. (2001a). Prenatal exposure to organic solvents and child neurobehavioral performance. Neurotoxicol Teratol 23: 235-245. [http://dx.doi.org/10.1016/S0892-0362\(01\)00141-6](http://dx.doi.org/10.1016/S0892-0362(01)00141-6).
- Till, C; Westall, C; Rovet, J; Koren, G. (2001b). Effects of maternal occupational exposure to organic solvents on offspring visual functioning: A prospective controlled study. Teratology 64: 134-141. <http://dx.doi.org/10.1002/tera.1056>.
- Till, C; Westall, C; Koren, G; Nulman, I; Rovet, J. (2005). Vision abnormalities in young children exposed prenatally to organic solvents. Neurotoxicology 26: 599-613. <http://dx.doi.org/10.1016/j.neuro.2005.05.011>.
- Tola, S; Vilhunen, R; Jarvinen, E; Korkala, M-L. (1980). A cohort study on workers exposed to trichloroethylene. J Occup Environ Med 22: 737-740.
- Toma, M; Grosser, M; Herr, A; Aust, D; Meye, A; Hoefling, C; Fuessel, S; Wuttig, D; Wirth, M; Baretton, G. (2008). Loss of heterozygosity and copy number abnormality in clear cell renal cell carcinoma discovered by high-density affymetrix 10K single nucleotide polymorphism mapping array. Neoplasia 10: 634-642.

- Tomisawa, H; Suzuki, S; Ichihara, S; Fukazawa, H; Tateishi, M. (1984). Purification and characterization of C-S lyase from *Fusobacterium varium*: A C-S cleavage enzyme of cysteine conjugates and some S-containing amino acids. *J Biol Chem* 259: 2588-2593.
- Tomisawa, H; Ichihara, S; Fukazawa, H; Ichimoto, N; Tateishi, M; Yamamoto, I. (1986). Purification and characterization of human hepatic cysteine-conjugate beta-lyase. *Biochem J* 235: 569-575.
- Tong, Z; Board, P; Anders, M. (1998). Glutathione transferase zeta catalyses the oxygenation of the carcinogen dichloroacetic acid to glyoxylic acid. *Biochem J* 331: 371-374.
- Toraason, M; Clark, J; Dankovic, D; Mathias, P; Skaggs, S; Walker, C; Werren, D. (1999). Oxidative stress and DNA damage in Fischer rats following acute exposure to trichloroethylene or perchloroethylene. *Toxicology* 138: 43-53. [http://dx.doi.org/10.1016/S0300-483X\(99\)00083-9](http://dx.doi.org/10.1016/S0300-483X(99)00083-9).
- Toth, B; Hornung, D; Scholz, C; Djalali, S; Friese, K; Jeschke, U. (2007). Peroxisome proliferator-activated receptors: new players in the field of reproduction. *Am J Reprod Immunol* 58: 289-310. <http://dx.doi.org/10.1111/j.1600-0897.2007.00514.x>.
- Trevisan, A; Giraldo, M; Borella, M; Maso, S. (2001). Historical control data on urinary and renal tissue biomarkers in naive male Wistar rats. *J Appl Toxicol* 21: 409-413.
- Triebig, G; Essing, H; Schaller, K; Valentin, H. (1976). [Biochemical and psychological examinations of trichloroethylene exposed volunteers (author's transl)]. *Zentralbl Bakteriol, Parasitenkd, Infektionskrankh Hyg, Abt 1: Orig, Reihe B* 163: 383-416.
- Triebig, G; Lehrl, S; Kinzel, W; Erzigkeit, H; Galster, J; Schaller, K. (1977a). [Psychopathometrical results of follow-up studies of trichloroethylene-exposed persons (author's transl)]. *Zentralbl Bakteriol, Parasitenkd, Infektionskrankh Hyg, Abt 1: Orig, Reihe B* 164: 314-377.
- Triebig, G; Schaller, K; Erzigkeit, H; Valentin, H. (1977b). Biochemische Untersuchungen und psychologische Studien an chronisch Trichlorathylen-belasteten Personen unter Berücksichtigung expositionsfreier Intervalle [Biochemical investigations and psychological studies of persons chronically exposed to trichloroethylene with regard to non-exposure intervals]. *Int Arch Occup Environ Health* 38: 149-162.
- Triebig, G; Schaller, K; Erzigkeit, H; Valentin, L. (1977c). [Biochemical investigations and psychological studies of persons chronically exposed to trichloroethylene with regard to non-exposure intervals]. *Int Arch Occup Environ Health* 38: 149-162. <http://dx.doi.org/10.1007/BF00378353>.
- Triebig, G; Trautner, P; Weltle, D; Saure, E; Valentin, H. (1982). [Investigations on neurotoxicity of chemical substances at the workplace: III. Determination of the motor and sensory nerve conduction velocity in persons occupationally exposed to trichloroethylene]. *Int Arch Occup Environ Health* 51: 25-34. <http://dx.doi.org/10.1007/BF00378407>.
- Triebig, G; Bestler, W; Baumeister, P; Valentin, H. (1983). [Investigations on neurotoxicity of chemical substances at the workplace: IV. Determination of the motor and sensory nerve conduction velocity in persons occupationally exposed to a mixture of organic solvents]. *Int Arch Occup Environ Health* 52: 139-150. <http://dx.doi.org/10.1007/BF00405417>.
- Tröster, A and Ruff, R. (1990). Neuropsychological sequelae of exposure to the chlorinated hydrocarbon solvents trichloroethylene and trichloroethane. *Arch Clin Neuropsychol* 5: 31-47. <http://dx.doi.org/10.1093/arclin/5.1.31>.
- Truhaut, R. (1951). [Contribution to the study of toxicology of trichloroethylene; Establishment of a method of determination applicable to atmosphere and biological media]. *Ann Pharm Fr* 9: 175-187.
- Tryndyak, V; Muskhelishvili, L; Kovalchuk, O; Rodriguez-Juarez, R; Montgomery, B; Churchwell, M; Ross, S; Beland, F; Pogribny, I. (2006). Effect of long-term tamoxifen exposure on genotoxic and epigenetic changes in rat liver: implications for tamoxifen-induced hepatocarcinogenesis. *Carcinogenesis* 27: 1713-1720. <http://dx.doi.org/10.1093/carcin/bgl050>.
- Tu, A; Murray, T; Hatch, K; Sivak, A; Milman, H. (1985). In vitro transformation of BALB/c-3T3 cells by chlorinated ethanes and ethylenes. *Cancer Lett* 28: 85-92. [http://dx.doi.org/10.1016/0304-3835\(85\)90096-5](http://dx.doi.org/10.1016/0304-3835(85)90096-5).

- Tucker, A; Sanders, V; Barnes, D; Bradshaw, T; White, K, Jr; Sain, L; Borzelleca, J; Munson, A. (1982). Toxicology of trichloroethylene in the mouse. *Toxicol Appl Pharmacol* 62: 351-357.
[http://dx.doi.org/10.1016/0041-008X\(82\)90137-5](http://dx.doi.org/10.1016/0041-008X(82)90137-5).
- Tugwood, J; Aldridge, T; Lambe, K; Macdonald, N; Woodyatt, N. (1996). Peroxisome proliferator-activated receptors: Structures and function. *Ann N Y Acad Sci* 804: 252-265.
<http://dx.doi.org/10.1111/j.1749-6632.1996.tb18620.x>.
- Tzeng, H-F; Blackburn, A; Board, P; Anders, M. (2000). Polymorphism- and species-dependent inactivation of glutathione transferase zeta by dichloroacetate. *Chem Res Toxicol* 13: 231-236.
<http://dx.doi.org/10.1021/tx990175q>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1984). Health assessment document for 1,1,1-trichloroethane (methyl chloroform). (EPA-600/8-82-003F). Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office.
<http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=300016I6.txt>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1986a). Guidelines for mutagenicity risk assessment. (EPA/630/R-98/003). pp. 23. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/iris/backgrd.html>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1986b). Guidelines for the health risk assessment of chemical mixtures. (EPA/630/R-98/002). pp. 38. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum.
<http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=22567>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1988). Recommendations for and documentation of biological values for use in risk assessment. (EPA/600/6-87/008). pp. 395. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Criteria and Assessment Office.
<http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=34855>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1991). Guidelines for developmental toxicity risk assessment. (EPA/600/FR-91/001). pp. 83. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/iris/backgrd.html>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1992). A cross-species scaling factor for carcinogen risk assessment based on equivalence of mg/kg^{3/4}/day. pp. 24. Washington, DC.
- U.S. EPA. (U.S. Environmental Protection Agency). (1994a). Methods for derivation of inhalation reference concentrations and application of inhalation dosimetry. (EPA/600/8-90/066F). pp. 409. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=71993>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1994b). Interim policy for particle size and limit concentration issues in inhalation toxicity studies. Washington, DC: U.S. Environmental Protection Agency, Health Effects Division, Office of Pesticide Products.
<http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=186068>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1994c). Final Draft for the Drinking Water Criteria Document on Chlorinated Acids/Aldehydes/Ketones/Alcohols. (EPA-68-C2-0139). pp. 276. Washington, D.C.
- U.S. EPA. (U.S. Environmental Protection Agency). (1995a). The use of the benchmark dose approach in health risk assessment. (EPA/630/R-94/007). pp. 93. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/raf/publications/useof-bda-healthrisk.htm>.
- U.S. EPA. (U.S. Environmental Protection Agency). (1995b). The national public water system supervision program: Compliance report, FY 1994. (EPA 812-R-95-001). Washington, DC: U.S. Environmental Protection Agency, Office Enforcement and Compliance Assurance, Office of Ground Water and Drinking Water.
<http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=20001TKR.txt>.

- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (1996). Guidelines for reproductive toxicity risk assessment. (EPA/630/R-96/009). pp. 143. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/raf/publications/pdfs/REPRO51.PDF>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (1997). Exposure factors handbook (final report). (EPA/600/P-95/002Fa-c). pp. 1193. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=12464>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (1998a). Guidelines for neurotoxicity risk assessment. (EPA/630/R-95/001F). pp. 89. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/raf/publications/pdfs/NEUROTOX.PDF>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (1998b). EPA data file: Office of groundwater and drinking water. Washington, DC.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2000a). Science policy council handbook: Risk characterization. (EPA 100-B-00-002). pp. 189. Washington, D.C.: U.S. Environmental Protection Agency, Office of Research and Development, Office of Science Policy. <http://www.epa.gov/osa/spc/pdfs/rchandbk.pdf>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2000b). Benchmark dose technical guidance document [external review draft]. (EPA/630/R-00/001). pp. 96. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/raf/publications/benchmark-dose-doc-draft.htm>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2000c). Supplementary guidance for conducting health risk assessment of chemical mixtures. (EPA/630/R-00/002). pp. 209. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=20533>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2001). Sources, emission and exposure for trichloroethylene (TCE) and related chemicals. (EPA/600/R-00/099). pp. 138. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. <http://www.epa.gov/nceaww1/pdfs/tce/TCEMarch2001.pdf>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2002a). The occurrence of disinfection by-products (DBPs) of health concern in drinking water: Results of a nationwide DBP occurrence study. (EPA/600/R-02/068). Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development, National Exposure Research Laboratory. http://epa.gov/athens/publications/reports/EPA_600_R02_068.pdf.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2002b). A review of the reference dose and reference concentration processes. (EPA/630/P-02/0002F). pp. 192. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=51717>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2002c). OSWER draft guidance for evaluating the vapor intrusion to indoor air pathway from groundwater and soils (subsurface vapor intrusion guidance). (EPA 530-D-02-004). Washington, DC. <http://epa.gov/osw/hazard/correctiveaction/eis/vapor/guidance.pdf>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2002d). Health assessment of 1,3-butadiene. (EPA/600/P-98/001F). pp. 435. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=54499>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2003a). Framework for cumulative risk assessment. (EPA/630/P-02/001F). pp. 129. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=30004TJH.txt>.
- [U.S. EPA.](#) (U.S. Environmental Protection Agency). (2003b). Toxicological review of dichloroacetic acid (CAS No. 79-43-6). (EPA 635/R-03/007). pp. 192. Washington, DC. <http://www.epa.gov/iris/toxreviews/0654tr.pdf>.

- [U.S. EPA.](http://water.epa.gov/lawsregs/rulesregs/regulatingcontaminants/sixyearreview/first_review/upload/support_6yr_occurancemethods_final.pdf) (U.S. Environmental Protection Agency). (2003c). Occurrence estimation methodology and occurrence findings report of the six-year review of existing national primary drinking water regulations. (EPA-815/R-03-006). pp. 874. Washington, DC: U.S. Environmental Protection Agency, Office of Water.
http://water.epa.gov/lawsregs/rulesregs/regulatingcontaminants/sixyearreview/first_review/upload/support_6yr_occurancemethods_final.pdf.
- [U.S. EPA.](http://water.epa.gov/action/advisories/drinking/upload/2005_05_06_criteria_drinking_percapita_2004.pdf) (U.S. Environmental Protection Agency). (2004). Estimated per capita water ingestion and body weight in the United States: An update. (EPA-822/R-00-001). pp. 521. Washington, DC: U.S. Environmental Protection Agency, Office of Water, Office of Science and Technology.
http://water.epa.gov/action/advisories/drinking/upload/2005_05_06_criteria_drinking_percapita_2004.pdf.
- [U.S. EPA.](http://www.epa.gov/airtoxics/nata1999/ted/hapem5guide.pdf) (U.S. Environmental Protection Agency). (2005a). The HAPEM5 user's guide: Hazardous air pollutant exposure model, version 5. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards.
<http://www.epa.gov/airtoxics/nata1999/ted/hapem5guide.pdf>.
- [U.S. EPA.](http://www.epa.gov/cancerguidelines/) (U.S. Environmental Protection Agency). (2005b). Guidelines for carcinogen risk assessment. (EPA/630/P-03/001F). pp. 166. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/cancerguidelines/>.
- [U.S. EPA.](http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm) (U.S. Environmental Protection Agency). (2005c). Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants (final). (EPA/630/P-03/003F). pp. 50. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. <http://www.epa.gov/raf/publications/guidance-on-selecting-age-groups.htm>.
- [U.S. EPA.](http://www.epa.gov/Region2/waste/ibm_e725.pdf) (U.S. Environmental Protection Agency). (2005d). Documentation of environmental indicator determination: RCRA corrective action. IBM-Endicott. Albany, NY.
http://www.epa.gov/Region2/waste/ibm_e725.pdf.
- [U.S. EPA.](http://www.epa.gov/cancerguidelines/guidelines-carcinogen-supplement.htm) (U.S. Environmental Protection Agency). (2005e). Supplemental guidance for assessing susceptibility from early-life exposure to carcinogens. (EPA/630/R-03/003F). pp. 126. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum.
<http://www.epa.gov/cancerguidelines/guidelines-carcinogen-supplement.htm>.
- [U.S. EPA.](http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=158363) (U.S. Environmental Protection Agency). (2006a). A framework for assessing health risk of environmental exposures to children. (EPA/600/R-05/093F). pp. 145. Washington, DC: U.S. Environmental Protection Agency, National Center for Environmental Assessment.
<http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=158363>.
- [U.S. EPA.](http://www.epa.gov/peerreview/pdfs/peer_review_handbook_2006.pdf) (U.S. Environmental Protection Agency). (2006b). Peer review handbook (3rd edition). (EPA/100/B-06/002). pp. 190. Washington, DC: U.S. Environmental Protection Agency, Science Policy Council. http://www.epa.gov/peerreview/pdfs/peer_review_handbook_2006.pdf.
- [U.S. EPA.](http://www.epa.gov/ttn/atw/nata1999/index.html) (U.S. Environmental Protection Agency). (2007a). Technology transfer network 1999 national-scale air toxics assessment, from <http://www.epa.gov/ttn/atw/nata1999/index.html>
- [U.S. EPA.](http://www.epa.gov/airdata/) (U.S. Environmental Protection Agency). (2007b). AirData: Access to Air Pollution Data (Version 2007): Office of Air Quality Planning and Standards.
- [U.S. EPA.](http://www.epa.gov/ogwdw000/databases/pdfs/data_factoids_2008.pdf) (U.S. Environmental Protection Agency). (2008a). Factoids: Drinking water and ground water statistics for 2008. (EPA-816/K-08-004). pp. 16. Washington, DC: U.S. Environmental Protection Agency, Office of Water.
http://www.epa.gov/ogwdw000/databases/pdfs/data_factoids_2008.pdf.
- [U.S. EPA.](http://www.epa.gov/IRIS/subst/0123.htm) (U.S. Environmental Protection Agency). (2008b). 1,1,2-Trichloro-1,2,2-trifluoroethane. Washington, D.C. <http://www.epa.gov/IRIS/subst/0123.htm>.
- [U.S. EPA.](http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=199243) (U.S. Environmental Protection Agency). (2008c). Child-specific exposure factors handbook. (EPA/600/R-06/096F). pp. 687. Washington, DC: U.S. Environmental Protection Agency, National Center for Environmental Assessment.
<http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=199243>.

- [U.S. EPA](#). (U.S. Environmental Protection Agency). (2011a). Recommended use of body weight 3/4 as the default method in derivation of the oral reference dose. (EPA/100/R11/0001). pp. 50. Washington, DC. <http://www.epa.gov/raf/publications/interspecies-extrapolation.htm>.
- [U.S. EPA](#). (U.S. Environmental Protection Agency). (2011b). Supplementary data for TCE assessment: Human posteriors by subject.
- [U.S. EPA](#). (U.S. Environmental Protection Agency). (2011c). Supplementary data for TCE assessment: Mouse posterior by subject.
- [U.S. EPA](#). (U.S. Environmental Protection Agency). (2011d). Supplementary data for TCE assessment: Rat posterior by subject.
- [U.S. EPA](#). (U.S. Environmental Protection Agency). (2011e). ADAF sample calculator. Washington, DC: Author.
- [U.S. EPA](#). (U.S. Environmental Protection Agency). (2011f). Age Dependent Adjustment Factor (ADAF) application: Office of Water Policy Document. pp. 21. Washington, DC.
- [U.S. GAO](#). (U.S. Government Accountability Office). (2007a). Testimony before the subcommittee on oversight and investigations, committee on energy and commerce, house of representatives: Defense health care: Issues related to past drinking water contamination at marine corps base Camp Lejeune. Washington, DC.
- [U.S. GAO](#). (U.S. Government Accountability Office). (2007b). Report to congressional committees: Defense health care: Activities related to past drinking water contamination at marine corps base Camp Lejeune. (GAO-07-276). pp. 81. Washington, DC.
- [Ulrich, R.](#) (2003). The toxicogenomics of nuclear receptor agonists. *Curr Opin Chem Biol* 7: 505-510. [http://dx.doi.org/10.1016/S1367-5931\(03\)00080-2](http://dx.doi.org/10.1016/S1367-5931(03)00080-2).
- [Ulrich, R; Rockett, J; Gibson, G; Pettit, S.](#) (2004). Overview of an interlaboratory collaboration on evaluating the effects of model hepatotoxicants on hepatic gene expression. *Environ Health Perspect* 112: 423-427.
- [Umezawa, T; Yonemoto, J; Soma, Y; Miura, T.](#) (1997). Behavioral effects of trichloroethylene and tetrachloroethylene in mice. *Pharmacol Biochem Behav* 58: 665-671. [http://dx.doi.org/10.1016/S0091-3057\(97\)00046-4](http://dx.doi.org/10.1016/S0091-3057(97)00046-4).
- [Urabe, Y; Nouso, K; Higashi, T; Nakatsukasa, H; Hino, N; Ashida, K; Kinugasa, N; Yoshida, K; Uematsu, S; Tsuji, T.](#) (1996). Telomere length in human liver diseases. *Liver Int* 16: 293-297. <http://dx.doi.org/10.1111/j.1600-0676.1996.tb00748.x>.
- [US EPA](#). (1980). Ambient water quality criteria for trichloroethylene. (EPA/440/5-80/077). pp. 66. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards, Criteria and Standards Division. <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=2000M4KP.txt>.
- [US EPA](#). (1985). Health assessment document for trichloroethylene: Final report. (EPA/600/8-82/006F). pp. 377. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office. <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=2000BFCV.txt>.
- [USAF](#). (U.S. Air Force). (2004). Determination of partition coefficients for a mixture of volatile organic compounds in rats and humans at different life stages. (AFRL-HE-WP-TR-2005-0012). Wright-Patterson Air Force Base, OH: U.S. Air Force Research Laboratory, Human Effectiveness Directorate, Applied Biotechnology Branch. <http://dodreports.com/pdf/ada436299.pdf>.
- [USGS](#). (U.S. Geological Survey). (2006). Volatile organic compounds in the nation's ground water and drinking-water supply wells. Reston, VA: U.S. Department of the Interior, U.S. Geological Survey. <http://pubs.usgs.gov/circ/circ1292/pdf/circular1292.pdf>.
- [Vagnarelli, P; De Sario, A; De Carli, L.](#) (1990). Aneuploidy induced by chloral hydrate detected in human lymphocytes with the Y97 probe. *Mutagenesis* 5: 591-592. <http://dx.doi.org/10.1093/mutage/5.6.591>.

- Vaidya, V; Shankar, K; Lock, E; Bucci, T; Mehendale, H. (2003a). Role of tissue repair in survival from s-(1,2-dichlorovinyl)-L-cysteine-induced acute renal tubular necrosis in the mouse. *Toxicol Sci* 74: 215-227. <http://dx.doi.org/10.1093/toxsci/kfg111>.
- Vaidya, V; Shankar, K; Lock, E; Bucci, T; Mehendale, H. (2003b). Renal injury and repair following S-1, 2 dichlorovinyl-L-cysteine administration to mice. *Toxicol Appl Pharmacol* 188: 110-121. [http://dx.doi.org/10.1016/S0041-008X\(02\)00080-7](http://dx.doi.org/10.1016/S0041-008X(02)00080-7).
- Vaidya, V; Shankar, K; Lock, E; Dixon, D; Mehendale, H. (2003c). Molecular mechanisms of renal tissue repair in survival from acute renal tubule necrosis: role of ERK1/2 pathway. *Toxicol Pathol* 31: 604-618.
- Vainio, H; Nickels, J; Linnainmaa, K. (1982). Phenoxy acid herbicides cause peroxisome proliferation in Chinese hamsters. *Scand J Work Environ Health* 8: 70-73.
- Vainio, H; Linnainmaa, K; Kähönen, M; Nickels, J; Hietanen, E; Marniemi, J; Peltonen, P. (1983). Hypolipidemia and peroxisome proliferation induced by phenoxyacetic acid herbicides in rats. *Biochem Pharmacol* 32: 2775-2779.
- Valic, E; Waldhör, T; Konnaris, C; Michitsch, A; Wolf, C. (1997). Acquired dyschromatopsia in combined exposure to solvents and alcohol. *Int Arch Occup Environ Health* 70: 403-406.
- Vamvakas, S; Dekant, W; Berthold, K; Schmidt, S; Wild, D; Henschler, D. (1987). Enzymatic transformation of mercapturic acids derived from halogenated alkenes to reactive and mutagenic intermediates. *Biochem Pharmacol* 36: 2741-2748. [http://dx.doi.org/10.1016/0006-2952\(87\)90258-9](http://dx.doi.org/10.1016/0006-2952(87)90258-9).
- Vamvakas, S; Dekant, W; Schiffmann, D; Henschler, D. (1988a). Induction of unscheduled DNA synthesis and micronucleus formation in Syrian hamster embryo fibroblasts treated with cysteine S-conjugates of chlorinated hydrocarbons. *Cell Biol Toxicol* 4: 393-403. <http://dx.doi.org/10.1007/BF00117768>.
- Vamvakas, S; Elfarra, A; Dekant, W; Henschler, D; Anders, M. (1988b). Mutagenicity of amino acid and glutathione S-conjugates in the Ames test. *Mutat Res* 206: 83-90.
- Vamvakas, S; Dekant, W; Henschler, D. (1989). Assessment of unscheduled DNA synthesis in a cultured line of renal epithelial cells exposed to cysteine S-conjugates of haloalkenes and haloalkanes. *Mutat Res* 222: 329-335. [http://dx.doi.org/10.1016/0165-1218\(89\)90108-0](http://dx.doi.org/10.1016/0165-1218(89)90108-0).
- Vamvakas, S; Bittner, D; Köster, U. (1993). Enhanced expression of the protooncogenes c-myc and c-fos in normal and malignant renal growth. *Toxicol Lett* 67: 161-172.
- Vamvakas, S; Richter, H; Bittner, D. (1996). Induction of dedifferentiated clones of LLC-PK1 cells upon long-term exposure to dichlorovinylcysteine. *Toxicology* 106: 65-74.
- Vamvakas, S; Brüning, T; Thomasson, B; Lammert, M; Baumüller, A; Bolt, H; Dekant, W; Birner, G; Henschler, D; Ulm, K. (1998). Renal cell cancer correlated with occupational exposure to trichloroethene. *J Cancer Res Clin Oncol* 124: 374-382. <http://dx.doi.org/10.1007/s004320050186>.
- van Bree, L; Commandeur, J; Lamberts, B; Cornelissen, M; van Roon, M; Laterveer, H; de Vries, J. (1990). Induction of drug metabolism enzymes by dihalogenated biphenyls. *J Biochem Toxicol* 5: 57-63. <http://dx.doi.org/10.1002/jbt.2570050109>.
- Van Hummelen, P and Kirsch-Volders, M. (1992). Analysis of eight known or suspected aneugens by the in vitro human lymphocyte micronucleus test. *Mutagenesis* 7: 447-455. <http://dx.doi.org/10.1093/mutage/7.6.447>.
- van Lieshout, E; Knapen, M; Lange, W; Steegers, E; Peters, W. (1998). Localization of glutathione S-transferases alpha and pi in human embryonic tissues at 8 weeks gestational age. *Hum Reprod* 13: 1380-1386.
- van Lieshout, E and Peters, W. (1998). Age and gender dependent levels of glutathione and glutathione S-transferases in human lymphocytes. *Carcinogenesis* 19: 1873-1875.
- Van Duuren, B; Goldschmidt, B; Loewengart, G; Smith, A; Melchionne, S; Seidman, I; Roth, D. (1979). Carcinogenicity of halogenated olefinic and aliphatic hydrocarbons in mice. *J Natl Cancer Inst* 63: 1433-1439.

- Vartiainen, T; Pukkala, E; Rienoja, T; Strandman, T; Kaksonen, K. (1993). Population exposure to tri- and tetrachloroethene and cancer risk: Two cases of drinking water pollution. Chemosphere 27: 1171-1181. [http://dx.doi.org/10.1016/0045-6535\(93\)90165-2](http://dx.doi.org/10.1016/0045-6535(93)90165-2).
- Veeramachaneni, D; Palmer, J; Amann, R. (2001). Long-term effects on male reproduction of early exposure to common chemical contaminants in drinking water. Hum Reprod 16: 979-987.
- Verma, Y and Rana, S. (2003). Gender differences in the metabolism of benzene, toluene and trichloroethylene in rat with special reference to certain biochemical parameters. J Environ Biol 24: 135-140.
- Vernon, R and Ferguson, R. (1969). Effects of trichloroethylene on visual-motor performance. Arch Environ Occup Health 18: 894-900.
- Vesterberg, O and Astrand, I. (1976). Exposure to trichloroethylene monitored by analysis of metabolites in blood and urine. J Occup Med 18: 224-226.
- Vesterberg, O; Gorczak, J; Krasts, M. (1976). Exposure to trichloroethylene II: Metabolites in blood and urine. Scand J Work Environ Health 2: 212-219.
- Vickers, A and Lucier, G. (1996). Estrogen receptor levels and occupancy in hepatic sinusoidal endothelial and Kupffer cells are enhanced by initiation with diethylnitrosamine and promotion with 17alpha-ethinylestradiol in rats. Carcinogenesis 17: 1235-1242.
- Vieira, I; Sonnier, M; Cresteil, T. (1996). Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. Eur J Biochem 238: 476-483. <http://dx.doi.org/10.1111/j.1432-1033.1996.0476z.x>.
- Vielhauer, V; Sarafoff, M; Gais, P; Rabes, H. (2001). Cell type-specific induction of O6-alkylguanine-DNA alkyltransferase mRNA expression in rat liver during regeneration, inflammation and preneoplasia. J Cancer Res Clin Oncol 127: 591-602.
- Vig, P; Russo, F; Edwards, R; Tadrous, P; Wright, N; Thomas, H; Alison, M; Forbes, S. (2006). The sources of parenchymal regeneration after chronic hepatocellular liver injury in mice. Hepatology 43: 316-324. <http://dx.doi.org/10.1002/hep.21018>.
- Villaschi, S; Giovanetti, A; Lombardi, C; Nicolai, G; Garbati, M; Andreozzi, U. (1991). Damage and repair of mouse bronchial epithelium following acute inhalation of trichloroethylene. Exp Lung Res 17: 601-614.
- Vineis, P; Miligi, L; Costantini, A. (2007). Exposure to solvents and risk of non-Hodgkin lymphoma: clues on putative mechanisms. Cancer Epidemiol Biomarkers Prev 16: 381-384. <http://dx.doi.org/10.1158/1055-9965.EPI-07-0124>.
- Vlaanderen, J; Lan, Q; Kromhout, H; Rothman, N; Vermeulen, R. (2011). Occupational benzene exposure and the risk of lymphoma subtypes: a meta-analysis of cohort studies incorporating three study quality dimensions. Environ Health Perspect 119: 159-167. <http://dx.doi.org/10.1289/ehp.1002318>.
- Vogelstein, B and Kinzler, K. (2004). Cancer genes and the pathways they control. Nat Med 10: 789-799. <http://dx.doi.org/10.1038/nm1087>.
- Vollmar, B; Pradarutti, S; Richter, S; Menger, M. (2002). In vivo quantification of ageing changes in the rat liver from early juvenile to senescent life. Liver Int 22: 330-341. <http://dx.doi.org/10.1034/j.1600-0676.2001.01631.x>.
- Von Tuneln, L; Yi, P; Bucci, T; Samokyszyn, V; Chou, M; Kadlubar, F; Fu, P. (2002). Tumorigenicity of chloral hydrate, trichloroacetic acid, trichloroethanol, malondialdehyde, 4-hydroxy-2-nonenal, crotonaldehyde, and acrolein in the B6C3F1 neonatal mouse. Cancer Lett 185: 13-19. [http://dx.doi.org/10.1016/S0304-3835\(02\)00231-8](http://dx.doi.org/10.1016/S0304-3835(02)00231-8).
- Voss, C; Zerban, H; Bannasch, P; Berger, M. (2005). Lifelong exposure to di-(2-ethylhexyl)-phthalate induces tumors in liver and testes of Sprague-Dawley rats. Toxicology 206: 359-371. <http://dx.doi.org/10.1016/j.tox.2004.07.016>.
- Voss, K; Riley, R; Dunn, C; Christopher Corton, J. (2006). The role of tumor necrosis factor alpha and the peroxisome proliferator-activated receptor alpha in modulating the effects of fumonisins in mouse liver. Toxicology 222: 165-174. <http://dx.doi.org/10.1016/j.tox.2006.02.012>.

- Wada, N; Marsman, D; Popp, J. (1992). Dose-related effects of the hepatocarcinogen, Wy-14,643, on peroxisomes and cell replication. *Fundam Appl Toxicol* 18: 149-154.
- Wainfan, E and Poirier, L. (1992). Methyl groups in carcinogenesis: effects on DNA methylation and gene expression. *Cancer Res* 52: 2071s-2077s.
- Walgren, J; Kurtz, D; McMillan, J. (2000a). The effect of the trichloroethylene metabolites trichloroacetate and dichloroacetate on peroxisome proliferation and DNA synthesis in cultured human hepatocytes. *Cell Biol Toxicol* 16: 257-273. <http://dx.doi.org/10.1023/A:1007638227821>.
- Walgren, J; Kurtz, D; McMillan, J. (2000b). Expression of PPAR(alpha) in human hepatocytes and activation by trichloroacetate and dichloroacetate. *Res Commun Mol Pathol Pharmacol* 108: 116-132.
- Wallace, LA. (1987). The total exposure assessment methodology (TEAM) study: Summary and analysis: Volume I. Washington, DC: U.S. Environmental Protection Agency.
- Wallace, L; Pellizzari, E; Hartwell, T; Rosenzweig, M; Erickson, M; Sparacino, C; Zelon, H. (1984). Personal exposure to volatile organic compounds: 1. Direct measurements in breathing-zone air, drinking water, food and exhaled breath. *Environ Res* 35: 293-319.
- Walles, S. (1986). Induction of single-strand breaks in dna of mice by trichloroethylene and tetrachloroethylene. *Toxicol Lett* 31: 31-35. [http://dx.doi.org/10.1016/0378-4274\(86\)90191-8](http://dx.doi.org/10.1016/0378-4274(86)90191-8).
- Wang, G and Stacey, N. (1990). Elevation of individual serum bile acids on exposure to trichloroethylene or alpha-naphthylisothiocyanate. *Toxicol Appl Pharmacol* 105: 209-215. [http://dx.doi.org/10.1016/0041-008X\(90\)90182-T](http://dx.doi.org/10.1016/0041-008X(90)90182-T).
- Wang, G; Takano, T; Tomita, K; Nakata, K; Nakamura, K. (1993). [Interaction of trichloroethylene and isopropyl alcohol in the perfused rat liver]. *Nippon Eiseigaku Zasshi* 48: 1000-1005.
- Wang, G; Ansari, G; Khan, M. (2007a). Involvement of lipid peroxidation-derived aldehyde-protein adducts in autoimmunity mediated by trichloroethene. *J Toxicol Environ Health A* 70: 1977-1985. <http://dx.doi.org/10.1080/15287390701550888>.
- Wang, G; Cai, P; Ansari, G; Khan, M. (2007b). Oxidative and nitrosative stress in trichloroethene-mediated autoimmune response. *Toxicology* 229: 186-193. <http://dx.doi.org/10.1016/j.tox.2006.10.014>.
- Wang, G; König, R; Ansari, G; Khan, M. (2008). Lipid peroxidation-derived aldehyde-protein adducts contribute to trichloroethene-mediated autoimmunity via activation of CD4+ T cells. *Free Radic Biol Med* 44: 1475-1482. <http://dx.doi.org/10.1016/j.freeradbiomed.2008.01.012>.
- Wang, J-D. (2004). Increased mortality odds ratio of male liver cancer in a community contaminated by chlorinated hydrocarbons in groundwater. Symposium on New Scientific Research Related to the Health Effects of Trichloroethylene. Washington, DC.
- Wang, J-L; Chen, W-L; Tsai, S-Y; Sung, P-Y; Huang, R-N. (2001). An in vitro model for evaluation of vaporous toxicity of trichloroethylene and tetrachloroethylene to CHO-K1 cells. *Chem Biol Interact* 137: 139-154. [http://dx.doi.org/10.1016/S0009-2797\(01\)00226-5](http://dx.doi.org/10.1016/S0009-2797(01)00226-5).
- Wang, R-S; Nakajima, T; Honma, T. (1999). Trichloroethylene inhibits aldehyde dehydrogenase only for aliphatic aldehydes of short chains in rats. *Toxicology* 132: 9-18. [http://dx.doi.org/10.1016/S0300-483X\(98\)00132-2](http://dx.doi.org/10.1016/S0300-483X(98)00132-2).
- Wang, R; Zhang, Y; Lan, Q; Holford, T; Leaderer, B; Zahm, S; Boyle, P; Dosemeci, M; Rothman, N; Zhu, Y; Qin, Q; Zheng, T. (2009). Occupational exposure to solvents and risk of non-Hodgkin lymphoma in Connecticut women. *Am J Epidemiol* 169: 176-185. <http://dx.doi.org/10.1093/aje/kwn300>.
- Wang, Z; Hall, S; Maya, J; Li, L; Asghar, A; Gorski, J. (2003). Diabetes mellitus increases the in vivo activity of cytochrome P450 2E1 in humans. *Br J Clin Pharmacol* 55: 77-85.
- Wanson, J-C; Bernaert, D; Penasse, W; Mosselmans, R; Bannasch, P. (1980). Separation in distinct subpopulations by elutriation of liver cells following exposure of rats to N-nitrosomorpholine. *Cancer Res* 40: 459-471.
- Ward, E; Thun, M; Hannan, L; Jemal, A. (2006). Interpreting cancer trends. *Ann N Y Acad Sci* 1076: 29-53. <http://dx.doi.org/10.1196/annals.1371.048>.

- Ward, J; Peters, J; Perella, C; Gonzalez, E. (1998). Receptor and nonreceptor-mediated organ-specific toxicity of di(2-ethylhexyl)phthalate (DEHP) in peroxisome proliferator-activated receptor alpha-null mice. *Toxicol Pathol* 26: 240-246.
- Warr, T; Parry, E; Parry, J. (1993). A comparison of two in vitro mammalian cell cytogenetic assays for the detection of mitotic aneuploidy using 10 known or suspected aneugens. *Mutat Res-Fundam Mol Mech Mutagen* 287: 29-46. [http://dx.doi.org/10.1016/0027-5107\(93\)90143-4](http://dx.doi.org/10.1016/0027-5107(93)90143-4).
- Warren, D; Graeter, L; Channel, S; Eggers, J; Goodyear, C; Macmahon, K; Sudberry, G; Latendresse, J; Fisher, J; Baker, W. (2006). Trichloroethylene, trichloroacetic acid, and dichloroacetic acid: do they affect eye development in the Sprague-Dawley rat. *Int J Toxicol* 25: 279-284. <http://dx.doi.org/10.1080/10915810600745975>.
- Wartenberg, D; Reyner, D; Scott, C. (2000). Trichloroethylene and cancer: Epidemiologic evidence. *Environ Health Perspect* 108: 161-176.
- Waseem, M; Ali, M; Dogra, S; Dutta, K; Kaw, J. (2001). Toxicity of trichloroethylene following inhalation and drinking contaminated water. *J Appl Toxicol* 21: 441-444.
- Waskell, L. (1978). A study of the mutagenicity of anesthetics and their metabolites. *Mutat Res-Fundam Mol Mech Mutagen* 57: 141-153. [http://dx.doi.org/10.1016/0027-5107\(78\)90261-0](http://dx.doi.org/10.1016/0027-5107(78)90261-0).
- Watanabe, S and Fukui, T. (2000). Suppressive effect of curcumin on trichloroethylene-induced oxidative stress. *J Nutr Sci Vitaminol* 46: 230-234.
- Watanabe, T; Shimada, H; Tanaka, Y. (1978). Human hepatocytes and aging: A cytophotometrical analysis in 35 sudden-death cases. *Virchows Arch B Cell Pathol Incl Mol Pathol* 27: 307-316.
- Watson, R. (2002). Effects of Phenobarbital on DNA Methylation in GC-Rich Regions of Hepatic DNA from Mice That Exhibit Different Levels of Susceptibility to Liver Tumorigenesis. *Toxicol Sci* 68: 51-58. <http://dx.doi.org/10.1093/toxsci/68.1.51>.
- Watson, R; Jacobson, C; Williams, A; Howard, W; DeSesso, J. (2006). Trichloroethylene-contaminated drinking water and congenital heart defects: a critical analysis of the literature. *Reprod Toxicol* 21: 117-147. <http://dx.doi.org/10.1016/j.reprotox.2005.07.013>.
- Waxman, S and Wurmbach, E. (2007). De-regulation of common housekeeping genes in hepatocellular carcinoma. *BMC Genomics* 8: 243. <http://dx.doi.org/10.1186/1471-2164-8-243>.
- Weiderpass, E; Pukkala, E; Kauppinen, T; Mutanen, P; Paakkulainen, H; Vasama-Neuvonen, K; Boffetta, P; Partanen, T. (1999). Breast cancer and occupational exposures in women in Finland. *Am J Ind Med* 36: 48-53. [http://dx.doi.org/10.1002/\(SICI\)1097-0274\(199907\)36:1<48::AID-AJIM7>3.0.CO;2-2](http://dx.doi.org/10.1002/(SICI)1097-0274(199907)36:1<48::AID-AJIM7>3.0.CO;2-2).
- Weiderpass, E; Vainio, H; Kauppinen, T; Vasama-Neuvonen, K; Partanen, T; Pukkala, E. (2003). Occupational exposures and gastrointestinal cancers among Finnish women. *J Occup Environ Med* 45: 305-315.
- Weidman, J; Dolinoy, D; Murphy, S; Jirtle, R. (2007). Cancer susceptibility: Epigenetic manifestation of environmental exposures. *Cancer J* 13: 9-16. <http://dx.doi.org/10.1097/PPO.0b013e31803c71f2>.
- Weisel, C and Jo, W-K. (1996). Ingestion, inhalation, and dermal exposures to chloroform and trichloroethene from tap water. *Environ Health Perspect* 104: 48-51.
- Weisel, C; Kim, H; Haltmeier, P; Klotz, J. (1999). Exposure estimates to disinfection by-products of chlorinated drinking water. *Environ Health Perspect* 107: 103-110.
- Weisenburger, D. (1992). Pathological classification of non-Hodgkin's lymphoma for epidemiological studies. *Cancer Res* 52: 5456s-5462s.
- Weiss, J; Moysich, K; Swede, H. (2005). Epidemiology of male breast cancer. *Cancer Epidemiol Biomarkers Prev* 14: 20-26.
- Weiss, P. (1971). A cell is not an island entire of itself. *Perspect Biol Med* 14: 182-205.
- Weistenhofer, W; Blaszkewicz, M; Bolt, H; Golka, K. (2008). N-acetyltransferase-2 and medical history in bladder cancer cases with a suspected occupational disease (BK 1301) in Germany. *J Toxicol Environ Health A* 71: 906-910. <http://dx.doi.org/10.1080/15287390801988681>.
- Wells, G; Schroth, W; Brauch, H; Ross, E. (2009). Bilateral renal-cell carcinoma associated with an acquired VHL mutation and long-term trichloroethylene exposure. *Clin Nephrol* 71: 708-713.

- Wendling, O; Chambon, P; Mark, M. (1999). Retinoid X receptors are essential for early mouse development and placentogenesis. PNAS 96: 547-551.
- Werner, H and Le Roith, D. (2000). New concepts in regulation and function of the insulin-like growth factors: Implications for understanding normal growth and neoplasia. Cell Mol Life Sci 57: 932-942.
- Werner, M; Birner, G; Dekant, W. (1995a). The role of cytochrome P4503A1/2 in the sex-specific sulfoxidation of the hexachlorobutadiene metabolite, N-acetyl-S-(pentachlorobutadienyl)-L-cysteine in rats. Drug Metab Dispos 23: 861-868.
- Werner, M; Guo, Z; Birner, G; Dekant, W; Guengerich, F. (1995b). The sulfoxidation of the hexachlorobutadiene metabolite N-acetyl-S-(1,2,3,4,4-pentachlorobutadienyl)-L-cysteine is catalyzed by human cytochrome P450 3A enzymes. Chem Res Toxicol 8: 917-923.
- Werner, M; Birner, G; Dekant, W. (1996). Sulfoxidation of mercapturic acids derived from tri- and tetrachloroethene by cytochromes P450 3A: A bioactivation reaction in addition to deacetylation and cysteine conjugate beta-lyase mediated cleavage. Chem Res Toxicol 9: 41-49.
<http://dx.doi.org/10.1021/tx950075u>.
- West, G; Woodruff, W; Brown, J. (2002). Allometric scaling of metabolic rate from molecules and mitochondria to cells and mammals. PNAS 99: 2473-2478.
<http://dx.doi.org/10.1073/pnas.012579799>.
- Westergren, I; Kjellstrand, P; Linder, L; Johansson, B. (1984). Reduction of brain specific gravity in mice prenatally exposed to trichloroethylene. Toxicol Lett 23: 223-226.
- Wheeler, C and Guenthner, T. (1990). Spectroscopic quantitation of cytochrome P-450 in human lung microsomes. J Biochem Toxicol 5: 269-272. <http://dx.doi.org/10.1002/jbt.2570050410>.
- Wheeler, C; Wrighton, S; Guenthner, T. (1992). Detection of human lung cytochromes P450 that are immunochemically related to cytochrome P450IIE1 and cytochrome P-450IIIA. Biochem Pharmacol 44: 183-187.
- White, A; Takehisa, S; Eger EI, I; Wolff, S; Stevens, W. (1979). Sister chromatid exchanges induced by inhaled anesthetics. 50: 426-430.
- White, J and Carlson, G. (1981a). Epinephrine-induced cardiac arrhythmias in rabbits exposed to trichloroethylene: role of trichloroethylene metabolites. Toxicol Appl Pharmacol 60: 458-465.
- White, J and Carlson, G. (1981b). Epinephrine-induced cardiac arrhythmias in rabbits exposed to trichloroethylene: potentiation by ethanol. Toxicol Appl Pharmacol 60: 466-471.
- White, K; David, D; Butterworth, L; Klykken, P. (2000). Assessment of autoimmunity-inducing potential using the brown Norway rat challenge model. Toxicol Lett 112-113: 443-451.
- White, R; Feldman, R; Eviator, I; Jabre, J; Niles, C. (1997). Hazardous waste and neurobehavioral effects: A developmental perspective. Environ Res 73: 113-124.
<http://dx.doi.org/10.1006/enrs.1997.3699>.
- WHO. (World Health Organization). (1977). Manual of the international statistical classification of diseases, injuries, and causes of death. Geneva, Switzerland.
<http://www.worldcat.org/oclc/3540720>.
- WHO. (World Health Organization). (1985). Environmental health criteria: Trichloroethylene. Geneva, Switzerland.
- Wideroff, L; Gridley, G; Chow, W-H; Linet, M; Mellemkjaer, L; Olsen, J; Keehn, S; Borch-Johnsen, K. (1997). Cancer incidence in a population-based cohort of patients hospitalized with diabetes mellitus in Denmark. J Natl Cancer Inst 89: 1360-1365.
<http://dx.doi.org/10.1093/jnci/89.18.1360>.
- Wiesenbüttner, B; Selinski, S; Golka, K; Brüning, T; Bolt, H. (2007). Re-assessment of the influence of polymorphisms of phase-II metabolic enzymes on renal cell cancer risk of trichloroethylene-exposed workers. Int Arch Occup Environ Health 81: 247-251. <http://dx.doi.org/10.1007/s00420-007-0200-5>.

- Wigle, D; Turner, M; Gomes, J; Parent, M. (2008). Role of hormonal and other factors in human prostate cancer. *J Toxicol Environ Health B Crit Rev* 11: 242-259.
<http://dx.doi.org/10.1080/10937400701873548>.
- Wilcosky, T; Checkoway, H; Marshall, E; Tyroler, H. (1984). Cancer mortality and solvent exposures in the rubber industry. *Am Ind Hyg Assoc J* 45: 809-811.
- Wilkens, L; Flemming, P; Gebel, M; Bleck, J; Terkamp, C; Wingen, L; Kreipe, H; Schlegelberger, B. (2004). Induction of aneuploidy by increasing chromosomal instability during dedifferentiation of hepatocellular carcinoma. *PNAS* 101: 1309-1314. <http://dx.doi.org/10.1073/pnas.0305817101>.
- Williams, A and DeSesso, J. (2008). Trichloroethylene and ocular malformations: analysis of extant literature. *Int J Toxicol* 27: 81-95. <http://dx.doi.org/10.1080/10915810701876760>.
- Williams, F; Meenagh, A; Darke, C; Acosta, A; Daar, A; Gorodezky, C; Hammond, M; Nascimento, E; Middleton, D. (2001). Analysis of the distribution of HLA-B alleles in populations from five continents. *Hum Immunol* 62: 645-650.
- Williams, P; Benton, L; Warmerdam, J; Sheehans, P. (2002). Comparative risk analysis of six volatile organic compounds in California drinking water. *Environ Sci Technol* 36: 4721-4728.
- Wilson, J. (1965). Methods for administering agents and detecting malformations in experimental animals. In J Warkany (Ed.), *Teratology: Principles and techniques* (8 ed., pp. 262-267). Chicago, IL: University of Chicago Press.
- Wilson, R; Donahue, M; Gridley, G; Adami, J; El Ghormli, L; Dosemeci, M. (2008). Shared occupational risks for transitional cell cancer of the bladder and renal pelvis among men and women in Sweden. *Am J Ind Med* 51: 83-99. <http://dx.doi.org/10.1002/ajim.20522>.
- Windemuller, F and Ettema, J. (1978). Effects of combined exposure to trichloroethylene and alcohol on mental capacity. *Int Arch Occup Environ Health* 41: 77-85.
<http://dx.doi.org/10.1007/BF00381792>.
- Windham, G; Shusterman, D; Swan, S; Fenster, L; Eskenazi, B. (1991). Exposure to organic solvents and adverse pregnancy outcome. *Am J Ind Med* 20: 241-259.
<http://dx.doi.org/10.1002/ajim.4700200210>.
- Windham, G; Zhang, L; Gunier, R; Croen, L; Grether, J. (2006). Autism spectrum disorders in relation to distribution of hazardous air pollutants in the San Francisco Bay area. *Environ Health Perspect* 114: 1438-1444.
- Winneke, G; Kramer, U; Kastka, J. (1976). Zur Beeinflussung psychomotor Leistungen Durch Alkohol und durch verschiedene Lösungsmitteldämpfe. In M Horvath (Ed.), *Adverse effects of environmental chemicals and psychotropic drugs* (Vol. 2, pp. 99-110). Amsterdam, The Netherlands: Elsevier.
- Winneke, G; Fodor, G; Schlipkoter, H. (1978). Carbon monoxide, trichloroethylene, and alcohol: Reliability and validity of neurobehavioral effects. In DA Otto (Ed.), *Multidisciplinary perspectives in event-related brain potential research: Proceedings of the Fourth International Congress on Event-Related Slow Potentials of the Brain (EPIC IV)*, Hendersonville, North Carolina, April 4-10, 1976 (pp. 461-469). Hendersonville, NC: U.S. Environmental Protection Agency.
- Winneke, G. (1982). Acute behavioral effects of exposure to some organic solvents - psychophysiological aspects. *Acta Neurol Scand Suppl* 92: 117-129.
- Withey, J; Collins, B; Collins, P. (1983). Effect of vehicle on the pharmacokinetics and uptake of four halogenated hydrocarbons from the gastrointestinal tract of the rat. *J Appl Toxicol* 3: 249-253.
<http://dx.doi.org/10.1002/jat.2550030506>.
- Withey, J and Karpinski, K. (1985). The fetal distribution of some aliphatic chlorinated hydrocarbons in the rat after vapor phase exposure. *Biol Res Pregnancy Perinatol* 6: 79-88.
- Wolf, D; Goldsworthy, T; Janszen, D; Harden, R; Donner, E; David, C; Everitt, J. (2000). Promotion by sodium barbital induces early development but does not increase the multiplicity of hereditary renal tumors in Eker rats. *Carcinogenesis* 21: 1553-1558.

- Wolff, D and Siegmund, R. (1978). [The circadian dependent effect of trichloroethylene on spontaneous locomotor activity and of tetrachloroethane on mortality in mice]. Biologisches Zentralblatt 97: 345-351.
- Wolfgang, G; Gandolfi, A; Stevens, J; Brendel, K. (1989a). N-acetyl S-(1,2-dichlorovinyl)-L-cysteine produces a similar toxicity to S-(1,2-dichlorovinyl)-L-cysteine in rabbit renal slices: Differential transport and metabolism. Toxicol Appl Pharmacol 101: 205-219.
[http://dx.doi.org/10.1016/0041-008X\(89\)90270-6](http://dx.doi.org/10.1016/0041-008X(89)90270-6).
- Wolfgang, G; Gandolfi, A; Stevens, J; Brendel, K. (1989b). In vitro and in vivo nephrotoxicity of the L and D isomers of S-(1,2-dichlorovinyl)-cysteine. Toxicology 58: 33-42.
- Wood, L; Parsons, D; Jones, S; Lin, J; Sjoblom, T; Leary, R; Shen, D; Boca, S; Barber, T; Ptak, J; Silliman, N; Szabo, S; Dezso, Z; Ustyanksky, V; Nikolskaya, T; Nikolsky, Y; Karchin, R; Wilson, P; Kaminker, J; Zhang, Z; Croshaw, R; Willis, J; Dawson, D; Shipitsin, M; Willson, J; Sukumar, S; Polyak, K; Park, B; Pethiyagoda, C; Pant, P; Ballinger, D; Sparks, A; Hartigan, J; Smith, D; Suh, E; Papadopoulos, N; Buckhaults, P; Markowitz, S; Parmigiani, G; Kinzler, K; Velculescu, V; Vogelstein, B. (2007). The genomic landscapes of human breast and colorectal cancers. Science 318: 1108-1113. <http://dx.doi.org/10.1126/science.1145720>.
- Woodruff, T; Axelrad, D; Caldwell, J; Morello-Frosch, R; Rosenbaum, A. (1998). Public health implications of 1990 air toxics concentrations across the United States. Environ Health Perspect 106: 245-251.
- Woods, C; Burns, A; Bradford, B; Ross, P; Kosyk, O; Swenberg, J; Cunningham, M; Rusyn, I. (2007a). WY-14,643-induced cell proliferation and oxidative stress in mouse liver are independent of NADPH oxidase. Toxicol Sci 98: 366-374. <http://dx.doi.org/10.1093/toxsci/kfm104>.
- Woods CG, Burns AM, Maki A, Bradford BU, Cunningham ML, Connor HD, Kadiiska MB, Mason RP, Peters JM, Rusyn I. 2006. Sustained formation of alpha-(4-pyridyl-1-oxide)-N-tert-butylnitron radical adducts in mouse liver by peroxisome proliferators is dependent upon peroxisome proliferator activated receptor-alpha, but not NADPH oxidase. Free Radic Biol Med. 42(3):335-42. <http://dx.doi.org/10.1016/j.freeradbiomed.2006.10.053>.
- Woods, C; Heuvel, J; Rusyn, I. (2007b). Genomic profiling in nuclear receptor-mediated toxicity. Toxicol Pathol 35: 474-494. <http://dx.doi.org/10.1080/01926230701311351>.
- Woolf, B. (1955). On estimating the relation between blood group and disease. Ann Hum Genet 19: 251-253. <http://dx.doi.org/10.1111/j.1469-1809.1955.tb01348.x>.
- Woolhiser, M; Krieger, S; Thomas, J; Hotchkiss, J. (2006). Trichloroethylene (TCE): Immunotoxicity potential in CD rats following a 4-week vapor inhalation exposure. Midland, MI: Unpublished.
- Wright, P; Thomas, W; Stacey, N. (1991). Effects of trichloroethylene on hepatic and splenic lymphocytotoxic activities in rodents. Toxicology 70: 231-242.
- Wu, C and Schaum, J. (2000). Exposure assessment of trichloroethylene. Environ Health Perspect 108: 359-363.
- Wu, C and Schaum, J. (2001). Sources, emissions and exposures for trichloroethylene (TCE) and related chemicals. (EPA/600/R-00/099). Washington, DC: U.S. Environmental Protection Agency.
- Wu, K and Berger, T. (2007). Trichloroethylene metabolism in the rat ovary reduces oocyte fertilizability. Chem Biol Interact 170: 20-30. <http://dx.doi.org/10.1016/j.cbi.2007.06.038>.
- Wu, K and Berger, T. (2008). Ovarian gene expression is stable after exposure to trichloroethylene. Toxicol Lett 177: 59-65. <http://dx.doi.org/10.1016/j.toxlet.2007.12.008>.
- Wu, X-Z and Chen, D. (2006). Origin of hepatocellular carcinoma: Role of stem cells. J Gastroenterol Hepatol 21: 1093-1098 <http://dx.doi.org/10.1111/j.1440-1746.2006.04485.x>.
- Xu, H; Tanphaichitr, N; Forkert, P-G; Anupriwan, A; Weerachatyanukul, W; Vincent, R; Leader, A; Wade, M. (2004). Exposure to trichloroethylene and its metabolites causes impairment of sperm fertilizing ability in mice. Toxicol Sci 82: 590-597. <http://dx.doi.org/10.1093/toxsci/kfh277>.
- Xu, W and Adler, I. (1990). Clastogenic effects of known and suspect spindle poisons studied by chromosome analysis in mouse bone marrow cells. Mutagenesis 5: 371-374.

- Xu, X; Yang, R; Wu, N; Zhong, P; Ke, Y; Zhou, L; Yuan, J; Li, G; Huang, H; Wu, B. (2009). Severe hypersensitivity dermatitis and liver dysfunction induced by occupational exposure to trichloroethylene. *Ind Health* 47: 107-112.
- Yamamura, K; Ikeda, T; Sadmoto, T; Maehara, N; Harabuchi, I; Takashima, H; Kiyosawa, H. (1983). Effects of trichloroethylene exposure on hearing: An investigation of cochlear microphonics and action potential of the guinea pig. *Eur J Appl Physiol* 52: 47-50.
<http://dx.doi.org/10.1007/BF00429024>.
- Yang, Q; Ito, S; Gonzalez, F. (2007). Hepatocyte-restricted constitutive activation of PPAR alpha induces hepatoproliferation but not hepatocarcinogenesis. *Carcinogenesis* 28: 1171-1177.
<http://dx.doi.org/10.1093/carcin/bgm046>.
- Yano, M; Yamamoto, T; Kobayashi, S; Ikeda, Y; Matsuzaki, M. (2008). Defective Ca²⁺ cycling as a key pathogenic mechanism of heart failure. *Circ J* 72 Suppl A: A22-A30.
- Yauck, J; Malloy, M; Blair, K; Simpson, P; McCarver, D. (2004). Proximity of residence to trichloroethylene-emitting sites and increased risk of offspring congenital heart defects among older women. *Birth Defects Res A Clin Mol Teratol* 70: 808-814.
<http://dx.doi.org/10.1002/bdra.20060>.
- Yeh, M; Larson, A; Campbell, J; Fausto, N; Rulyak, S; Swanson, P. (2007). The expression of transforming growth factor-alpha in cirrhosis, dysplastic nodules, and hepatocellular carcinoma: An immunohistochemical study of 70 cases. *Am J Surg Pathol* 31: 681-689.
<http://dx.doi.org/10.1097/PAS.0b013e31802ff7aa>.
- Yeldandi, A; Milano, M; Subbarao, V; Reddy, J; Rao, M. (1989). Evaluation of liver cell proliferation during ciprofibrate-induced hepatocarcinogenesis. *Cancer Lett* 47: 21-27.
[http://dx.doi.org/10.1016/0304-3835\(89\)90172-9](http://dx.doi.org/10.1016/0304-3835(89)90172-9).
- Yeung, R; Xiao, G; Everitt, J; Jin, F; Walker, C. (1995). Allelic loss at the tuberous sclerosis 2 locus in spontaneous tumors in the Eker rat. *Mol Carcinog* 14: 28-36.
- Yoon, H; Monks, T; Walker, C; Lau, S. (2001). Transformation of kidney epithelial cells by a quinol thioether via inactivation of the tuberous sclerosis-2 tumor suppressor gene. *Mol Carcinog* 31: 37-45.
- Yoon, J; Mason, J; Valencia, R; Woodruff, R; Zimmering, S. (1985). Chemical mutagenesis testing in Drosophila. IV. Results of 45 coded compounds tested for the National Toxicology Program. *Environ Mutagen* 7: 349-367. <http://dx.doi.org/10.1002/em.2860070310>.
- Yoon, M; Madden, M; Barton, H. (2007). Extrahepatic metabolism by CYP2E1 in PBPK modeling of lipophilic volatile organic chemicals: Impacts on metabolic parameter estimation and prediction of dose metrics. *J Toxicol Environ Health A* 70: 1527-1541.
- Yoshida, M; Fukabori, S; Hara, K; Yuasa, H; Nakaaki, K; Yamamura, Y; Yoshida, K. (1996). Concentrations of trichloroethylene and its metabolites in blood and urine after acute poisoning by ingestion. *Hum Exp Toxicol* 15: 254-258. <http://dx.doi.org/10.1177/096032719601500312>.
- Yoshioka, T; Krauser, J; Guengerich, F. (2002). Microsomal oxidation of tribromoethylene and reactions of tribromoethylene oxide. *Chem Res Toxicol* 15: 1414-1420.
- Yu, K; Barton, H; Mahle, D; Frazier, J. (2000). In vivo kinetics of trichloroacetate in male Fischer 344 rats. *Toxicol Sci* 54: 302-311.
- Yu, M; Garabrant, D; Peters, J; Mack, T. (1988). Tobacco, alcohol, diet, occupation, and carcinoma of the esophagus. *Cancer Res* 48: 3843-3848.
- Yu, S; Rao, S; Reddy, J. (2003). Peroxisome proliferator-activated receptors, fatty acid oxidation, steatohepatitis and hepatocarcinogenesis. *Curr Mol Med* 3: 561-572.
- Yuan, J. (1995). Effects of drinking pattern on the peak/trough blood concentrations in drinking water studies. *Food Chem Toxicol* 33: 565-571.
- Yuan, J; Castelao, J; Gago-Dominguez, M; Yu, M; Ross, R. (1998). Tobacco use in relation to renal cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 7: 429-433.

- Yue, F; Huang, H; Huang, J; Liang, L; Huang, Z; Wei, Q; Chen, X. (2007). HLA-DM polymorphism and risk of trichloroethylene induced medicamentosa-like dermatitis. *Biomed Environ Sci* 20: 506-511.
- Zack, M; Cannon, S; Loyd, D; Heath, C, Jr; Falletta, J; Jones, B; Housworth, J; Crowley, S. (1980). Cancer in children of parents exposed to hydrocarbon-related industries and occupations. *Am J Epidemiol* 111: 329-336.
- Zahm, S. (1992). Computerized occupational referent population system (CORPS): Study documentation. Rockville, MD: National Cancer Institute & National Institute for Occupational Safety and Health.
- Zajicek, G; Schwartz-Arad, D; Bartfeld, E. (1989). The streaming liver V: Time and age-dependent changes of hepatocyte DNA content, following partial hepatectomy. *Liver* 9: 164-171. <http://dx.doi.org/10.1111/j.1600-0676.1989.tb00394.x>
- Zajicek, G and Schwartz-Arad, D. (1990). Streaming liver VII: DNA turnover in acinus zone-3. *Liver* 10: 137-140. <http://dx.doi.org/10.1111/j.1600-0676.1990.tb00448.x>
- Zajicek, G; Arber, N; Schwartz-Arad, D. (1991). Streaming liver VIII: Cell production rates following partial hepatectomy. *Liver* 11: 347-351. <http://dx.doi.org/10.1111/j.1600-0676.1991.tb00541.x>
- Zandi-Nejad, K; Eddy, A; Glasscock, R; Brenner, B. (2004). Why is proteinuria an ominous biomarker of progressive kidney disease. *Kidney Int Suppl* S76-S89. <http://dx.doi.org/10.1111/j.1523-1755.2004.09220.x>
- Zanelli, U; Puccini, P; Acerbi, D; Ventura, P; Gervasi, P. (1996). Induction of peroxisomal beta-oxidation and P-450 4A-dependent activities by pivalic and trichloroacetic acid in rat liver and kidney. *Arch Toxicol* 70: 145-149.
- Zeise, L; Wilson, R; Crouch, E. (1987). Dose-response relationships for carcinogens: A review. *Environ Health Perspect* 73: 259-306.
- Zenick, H; Blackburn, K; Hope, E; Richdale, N; Smith, M. (1984). Effects of trichloroethylene exposure on male reproductive function in rats. *Toxicology* 31: 237-250. [http://dx.doi.org/10.1016/0300-483X\(84\)90105-7](http://dx.doi.org/10.1016/0300-483X(84)90105-7)
- Zeppa, P; Benincasa, G; Troncone, G; Lucariello, A; Zabatta, A; Cochand-Priollet, B; Fulciniti, F; Vetrani, A; De Rosa, G; Palombini, L. (1998). Retrospective evaluation of DNA ploidy of hepatocarcinoma on cytologic samples. *Diagn Cytopathol* 19: 323-329. [http://dx.doi.org/10.1002/\(SICI\)1097-0339\(199811\)19:5<323::AID-DC2>3.0.CO;2-Y](http://dx.doi.org/10.1002/(SICI)1097-0339(199811)19:5<323::AID-DC2>3.0.CO;2-Y)
- Zhang, B; Pan, X; Cobb, G; Anderson, T. (2007). microRNAs as oncogenes and tumor suppressors. *Dev Biol* 302: 1-12. <http://dx.doi.org/10.1016/j.ydbio.2006.08.028>
- Zhang, G and Stevens, J. (1989). Transport and activation of S-(1,2-dichlorovinyl)-L-cysteine and N-acetyl-S-(1,2-dichlorovinyl)-L-cysteine in rat kidney proximal tubules. *Toxicol Appl Pharmacol* 100: 51-61. [http://dx.doi.org/10.1016/0041-008X\(89\)90091-4](http://dx.doi.org/10.1016/0041-008X(89)90091-4)
- Zhang, J and Yu, K. (1998). What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. *JAMA* 280: 1690-1691.
- Zhang, L; Steinmaus, C; Eastmond, D; Xin, X; Smith, M. (2009). Formaldehyde exposure and leukemia: A new meta-analysis and potential mechanisms. *Mutat Res Rev Mutat Res* 681: 150-168. <http://dx.doi.org/10.1016/j.mrrev.2008.07.002>
- Zhang, S; Miao, D; Liu, A; Zhang, L; Wei, W; Xie, H; Lu, W. (2010). Assessment of the cytotoxicity and genotoxicity of haloacetic acids using microplate-based cytotoxicity test and CHO/HGPRT gene mutation assay. *Mutat Res* 703: 174-179. <http://dx.doi.org/10.1016/j.mrgentox.2010.08.014>
- Zhang, Y; Cantor, K; Lynch, C; Zheng, T. (2004). A population-based case-control study of occupation and renal cell carcinoma risk in Iowa. *J Occup Environ Med* 46: 235-240.
- Zhao, Y; Krishnadasan, A; Kennedy, N; Morgenstern, H; Ritz, B. (2005). Estimated effects of solvents and mineral oils on cancer incidence and mortality in a cohort of aerospace workers. *Am J Ind Med* 48: 249-258. <http://dx.doi.org/10.1002/ajim.20216>

- Zhou, H; Randers-Pehrson, G; Geard, C; Brenner, D; Hall, E; Hei, T. (2003). Interaction between radiation-induced adaptive response and bystander mutagenesis in mammalian cells. *Radiat Res* 160: 512-516.
- Zhou, Y and Waxman, D. (1998). Activation of peroxisome proliferator-activated receptors by chlorinated hydrocarbons and endogenous steroids. *Environ Health Perspect* 106 Suppl 4: 983-988.
- Zhu, J; Newhook, R; Marro, L; Chan, C. (2005). Selected volatile organic compounds in residential air in the city of Ottawa, Canada. *Environ Sci Technol* 39: 3964-3971.
<http://dx.doi.org/10.1021/es050173u>.
- Zielinska-Psuja, B; Orlowski, J; Kowalowka, J; Zieba-Proc, B. (1998). [Metabolic implications of benzene and trichloroethylene: Part I: The effect trichloroethylene on phenol elimination in rats]. *Bromatol Chem Toksykol* 31: 13-20.
- Zielinski, A. (1973). General health state of women professionally exposed to trichloroethylene vapours. *Med Pr* 24: 263-271.
- Zordan, M; Osti, M; Pesce, M; Costa, R. (1994). Chloral hydrate is recombinogenic in the wing spot test in *Drosophila melanogaster*. *Mutat Res* 322: 111-116. [http://dx.doi.org/10.1016/0165-1218\(94\)00017-4](http://dx.doi.org/10.1016/0165-1218(94)00017-4).